

Heme Oxygenase-1 in Tumor Biology and Therapy

Halina Was, Jozef Dulak and Alicja Jozkowicz*

Department of Medical Biotechnology, Faculty of Biochemistry, Biophysics and Biotechnology, Jagiellonian University, Krakow, Poland

Abstract: Heme oxygenase-1 (HO-1) degrades heme to carbon monoxide (CO), biliverdin, and ferrous iron. As HO-1 expression is highly increased by stressful conditions, the major role of the enzyme is the protection against oxidative injury. Additionally, it regulates cell proliferation, modulates inflammatory response and facilitates angiogenesis. Beneficial activities of HO-1 have been recognized in many pathological states e.g. atherosclerosis, diabetes, ischemia/reperfusion injury or organ transplantation. Interestingly HO-1 expression is very often boosted in tumor tissues and could be further elevated in response to radio-, chemo-, or photodynamic therapy. A growing body of evidence suggests that HO-1 may play a role in tumor induction and can potentially improve the growth and spread of tumors. This review discusses the implications of HO-1 properties for tumor proliferation and cell death, differentiation, angiogenesis and metastasis, and tumor-related inflammation. Finally, it suggests that pharmacological agents that regulate HO activity or HO-1 gene silencing may become powerful tools for preventing the onset or progression of various cancers and sensitize them to anticancer therapies.

Keywords: Heme oxygenase-1, carcinogenesis, oxidative stress, cytoprotection, angiogenesis, metastasis.

CARCINOGENESIS

Transition from normal to pre-cancer and cancer cells is a result of multi-step accumulation of genetic and epigenetic modifications. In many types of cancers the aneuploidy is detected and pre-malignant lesions with aneuploidy can transform into malignancy more frequently than that with normal DNA content [1]. Growing body of evidence suggests that aberration in the number of chromosomes is a cause rather than a consequence of malignant transformation [2]. Also the loss of heterozygosity, defined as a loss of genomic material in one of a pair of chromosomes in regions that include tumor suppressor genes, is an early predictor of transformation of pre-cancerous lesions [3]. Finally, the telomerase activity, necessary for maintenance or extension of telomeres, is generally not found in normal tissues, but occurs in low levels in pre-cancerous lesions and in cancers [4].

Deregulations or mutations of genes responsible for cellular growth and differentiation, such as growth factors, receptor or cytoplasmatic tyrosine kinases, serine/threonine kinases, regulatory GTPases, and transcription factors, play a chief role in tumor development. Oncogenes encoded by mutated genes do not possess important regulatory elements and their production and activation is uncontrolled [5]. Cancer-related genes include also tumor suppressors. Their protein products either diminish the cell cycle or favor apoptosis. It has been shown that tumor suppressor genes are important targets in sensing and responding to DNA damage in the multi-step process of carcinogenesis [6, 7].

One of the critical processes correlated with malignancy is angiogenesis. It is a crucial event not only for growth of

primary tumors but also for metastasis. Tumor-associated angiogenic factors, such as basic fibroblast growth factor (bFGF) and vascular endothelial growth factor (VEGF), are generated both by tumor cells and by infiltrating leukocytes [8]. Recently, a role of tumor vasculogenesis was also indicated [9].

An inflammatory microenvironment is an essential component of all tumors [10]. Inflammation influences different stages of tumorigenesis, including initiation, promotion, malignant conversion, invasion, and metastasis. It also affects immune surveillance and responses to therapy [11]. Bacterial and viral infections [12], tobacco smoke [13], obesity [14, 15], patient age [16], and cell senescence [17] are postulated to be the tumor promoters that act through inflammatory mechanisms.

Final important aspect of tumorigenesis is an oxidative stress that can modulate all steps of the process. It is induced by reactive oxygen species (ROS), including not only the oxygen radicals (superoxide anion radical, hydroxyl radicals, etc.) but also some non-radical derivatives (hydrogen peroxide, singlet oxygen, alkyl peroxide etc.) [18]. Strong oxidative stress leads to cytotoxicity, inhibits cell proliferation and results in apoptotic/necrotic cell death, whereas low or intermediate amounts of ROS cause DNA damage and mutations, and induce inflammatory reaction, leading to augmented cell proliferation and increasing risk of carcinogenesis [19]. ROS levels are precisely regulated by endogenous defense systems, including intracellular superoxide dismutase (SOD), catalase, glutathione peroxidase, and heme oxygenase-1 (HO-1) [20].

EXPRESSION AND ACTIVITY OF HEME OXYGENASES

Heme oxygenases, the rate-limiting enzymes in heme catabolism, catalyze the stereospecific degradation of heme to biliverdin, with the concurrent release of ferrous iron

*Address correspondence to this author at the Department of Medical Biotechnology, Faculty of Biochemistry, Biophysics and Biotechnology, Jagiellonian University, Gronostajowa 7, 30-387 Krakow, Poland; Tel: +48-12-664- 6411; Fax: +48-12-664-6918; E-mail: alicja.jozkowicz@uj.edu.pl

ions and carbon monoxide (CO) [21]. In addition to their role in regulating the cellular levels of heme, HO is responsible for the recycling of iron from senescent red blood cells. Released iron induces the expression of the iron-sequestering ferritin and activates Fe-ATPase, an iron transporter, which reduce intracellular Fe²⁺ content [22]. The second end-product of HO activity, CO, is exhaled from the organisms through the lung [23]. Finally, biliverdin is converted to bilirubin by the cytosolic enzyme biliverdin reductase (BvR). Bilirubin is subsequently oxidized by cytochrome P450 enzymes or glucuronidated by UDP-glucuronyl transferase and then excreted as bilirubin glucuronides into the bile [24, 25] (Fig. 1).




Fig. (1). Schematic demonstration of HO-1 pathway.

Heme oxygenase was first purified to homogeneity from rat liver and then from porcine and bovine spleen [26-28]. Subsequently it was detected in human tissues [29]. There are two isoforms of HO, which are encoded by two distinct genes, but share approximately 40% of amino acid sequence homology [30, 31]. The originally isolated enzyme, with molecular mass of 32 kDa, is designated as HO-1, whereas the second protein, with molecular mass of 36 kDa, is designated as HO-2 [32, 33]. HO-1 is the product of one transcript, but HO-2 is encoded by two transcripts of the single gene which differ in the use of the polyadenylation signal [34].

HO-2 is a constitutively expressed protein, detected at high levels in the brain and, at lower levels, also in testes, endothelium, distal nephron segments, liver, and gut myenteric plexus [32, 33]. In contrast, HO-1 can be strongly induced in many tissues in response to cellular stress caused by a wide spectrum of stimuli, including heme [35], heavy metals [35], UV irradiation, ROS [36], nitric oxide (NO) [37], inflammatory cytokines [38], heat shock [39], ethanol [40], and prostaglandins [41] (Fig. 1). It can also be activated by hypoxia, but this effect is tissue- and species-dependent [42, 43]. On the other hand, HO-1 seems to be constitutively expressed in renal inner medullary cells [44], Kupffer cells

in the liver [45], Purkinje cells in the cerebellum [46], and CD4⁺/CD25⁺ regulatory T lymphocytes [47]. Enzymatic activity of HO-1 reduces oxidative stress, diminishes inflammatory response, and lowers the rate of apoptosis. Namely, it enables the removal of heme, a potentially toxic prooxidant molecule, allows decreasing a lipid-soluble transmissible form of iron, and leads to the generation of biologically active products.

Heme is an essential iron complex responsible for oxygen and electron transport. It can be released from the heme proteins and, together with the released iron, may catalyze free radical reactions, being harmful to the cells. The major source of heme are erythrocytes, therefore the blood vessels are at the greatest risk of exposure [48]. The hydrophobicity of heme analogues (ferriporphyrins) is decisive for entry into cells and necessary for oxidative injury. It has been shown that ferrihemoglobin can readily be oxidized to heme-releasing ferrihemoglobin in the presence of inflammatory cell-derived oxidants [49-51]. Additionally, heme originating from hemoglobin can interact with NO to enhance heme uptake [52]. Importantly, heme induces neutrophil activation [53], augments endothelial cell adhesion molecule expression and subsequently triggers robust inflammatory reactions [54, 55]. The uptake of heme and resulting augmentation of cellular oxidant vulnerability are both blocked by the heme-binding proteins, such as hemopexin [56, 57], albumin, and haptoglobin [58].

Degradation of heme by HO is the major source of CO, previously regarded as a toxic air pollutant, but currently recognized as an important mediator which, similarly to NO, plays a role in neurotransmission and muscle relaxation. It appears that HO-1-derived CO and bilirubin exert a vasorelaxant effect not only *via* a cGMP-dependent pathway but also *via* a cGMP-independent activation of voltage-gated K⁺ channels [59] or indirectly through an increase in adiponectin levels [60]. Due to induction of soluble guanylyl cyclase (sGC), CO also inhibits platelet aggregation, decreases leukocyte adhesion, and reduces endothelial cell apoptosis. Finally, anti-inflammatory properties of CO are related to reduced production of proinflammatory cytokines such as tumor necrosis factor (TNF), interleukin-1 β (IL-1 β), or macrophage inflammatory protein-1 β (MIP-1 β), and concomitant upregulation of anti-inflammatory IL-10 [61].

Ferrous iron ions, the second product of heme degradation, can be dangerous, taking a part in the Fenton reaction to generate toxic hydroxyl radicals, which can oxidize lipids and damage the DNA and proteins. However, iron can be sequestered by ferritin or removed by ATP-dependent pumps, preventing oxidative stress [62]. At the end, both biliverdin and bilirubin, which have been long considered as toxic end-products of heme metabolism, turned out to be the potent antioxidants and inhibitors of complement cascade. Thus, unconjugated bilirubin is capable of scavenging the singlet oxygen and can act as a reducing agent for certain peroxidases, including horseradish peroxidase and prostaglandin H synthase [24].

EXPRESSION OF HO-1 IN TUMORS

Expression of HO-1 is usually higher in cancer cells than in surrounding healthy tissues, as shown for lymphosarcoma

[63], prostate cancers [64-66], brain tumors [67], adenocarcinoma [68], hepatoma [69, 70], squamous carcinoma [71, 72], glioblastoma [73], melanoma [74], Kaposi sarcoma [75] or pancreatic carcinoma [76]. It can be induced by some oncogenes, such as viral G protein-coupled receptor (vGPCR) encoded by HHV-8, a Kaposi sarcoma-associated herpes virus [77]. Also the BCR/ABL fusion kinase, typical for chronic myeloid leukemia, was demonstrated to upregulate HO-1 [78]. Concomitantly, chronic arsenic-induced malignant transformation of rat liver epithelial cells increases the levels of oncogenes, namely α -fetoprotein (AFP), Wilm's tumor protein-1 (WT-1), c-jun, c-myc, and H-ras, which is accompanied by elevation of HO-1 [79]. Of importance, co-expression of c-Myc and TGF α in the mouse liver, that accelerates hepatocarcinogenesis, was related to upregulation of HO-1. The effect was abolished by supplementation with vitamin E, suggesting the role of ROS [80] (Table 1). Noteworthy, tumors showing strong neovascularization and massive hemorrhages contain significant amounts of heme-releasing oxidized hemoglobin [81]. Heme can directly induce HO-1 expression, what is a protective mechanism against brain [82], intestine [83] or hepatic oxidative injury [84]. Finally, expression of HO-1 in cancer cells can be further elevated in response to chemotherapy, irradiation [76, 85], or photodynamic therapy [86-89].

Table 1. Elevated Expression of HO-1 in Tumors

Type of Cancer	References
Lymposarcoma	[63]
Prostate cancer	[64-66]
Brain tumor	[67]
Adenocarcinoma	[68]
Hepatoma	[69, 70]
Squamous carcinoma	[71, 72]
Glioblastoma	[73]
Melanoma	[74]
Kaposi sarcoma	[75]
Pancreatic cancer	[76]
Chronic myeloid leukemia	[78]

The exact location of HO-1 in transformed tissue may depend on type of tumor. As found in human melanomas [74] or in rat and human gliomas [73] HO-1 was almost exclusively expressed in macrophages, which accumulated in perinecrotic areas [73]. Interestingly, macrophages of non-small cell lung cancer exhibited decreased level of HO-1 in comparison to tumor-free lung macrophages [90]. On the other hand, in rat hepatoma HO-1 was found only in tumor cells [69]. Furthermore, expression of HO-1 in hormone-refractory prostate cancer (HRPCA) was much higher than in localized prostate cancer (PCA) or benign prostate tissues, while there were no such differences in stromal cells surrounding the tumors [66]. Finally, human pancreatic carcinoma displayed marked HO-1 immunoreactivity both in cancer cells and in immunocytes [76].

Recently, a nuclear localization of HO-1 has been shown in human prostate cancer [65, 91], especially in cells treated with hemin [65]. For the first time, it was observed in murine fibroblast and human hepatoma cell lines, where HO-1 translocated to the nucleus leading to activation of oxidant-responsive transcription factors, including activator protein-1 (AP-1). Thus, it was postulated that the nuclear form of HO-1, although enzymatically inactive, may upregulate genes that promote cytoprotection against oxidative stress [92]. Additionally, the nuclear localization of HO-1 was demonstrated in astroglia and implicated in brain development or neurodegenerative diseases [93]. It was also found in rat fetal lung cells exposed to hyperoxia, where it could act as a chaperone or nuclear messenger [94], and in brown adipocytes, where it was suggested as a transcription factor playing a role in adipogenesis [95].

ROLE OF HO-1 IN TUMOR CELL PROLIFERATION

HO-1 is highly upregulated in fast proliferating cells such as epithelium within the wounded skin or psoriatic lesions [96] and cancer cells in the growing tumors [64]. It also seems to affect the cell cycle progression, but in a cell-type specific manner. Thus, HO-1 stimulated the proliferation of keratinocytes [97], vascular endothelium [77, 98], or regulatory CD4⁺CD25⁺ T lymphocytes [99, 100], and these effects were reversed by its pharmacological inhibitor (tin protoporphyrin-IX, SnPPIX) or by siRNA [97]. In contrast, HO-1 attenuated divisions of fibroblasts [101], smooth muscle cells [102-104], epithelial cells [105], effector T lymphocytes [106], and astroglia [107]. Again, this effect could be reversed by HO-1 inhibition, confirming the HO-1 specificity [102].

Similarly in tumors, the influence of HO-1 on proliferation depends much on the cell type. Its direct pro-mitogenic action has been shown in pancreatic cancer, where siRNA-mediated downregulation of HO-1 was coupled with a statistically significant inhibition of growth [76], and in murine or human melanoma, where overexpression of HO-1 led to increased cell proliferation [108]. Also, enhancement in divisions of transformed endothelial cells infected with HHV-8 resulted from induction of HO-1, as it was abolished by HO-1 inhibitor (chromium mesoporphyrin-IX, CrMPPIX) [75]. Pharmacological or genetic inhibition of HO-1 reduced also the proliferation of prostate carcinoma [66], hepatocellular carcinoma [70], and neoplastic mast cells [109]. Finally, knockdown of HO-1 caused marked growth inhibition in urothelial cancer cell line [88].

In turn, the antiproliferative actions of HO-1 have been demonstrated for lung adenocarcinoma, breast cancer and prostate cancer cells. Namely, HO-1 inhibition with SnPPIX reversed the growth arrest observed in A549 lung adenocarcinoma overexpressing HO-1 [110]. Similarly, application of SnPPIX resulted in a small but significant increase in proliferation of rat and human breast cancer cell lines, whereas induction of HO-1 with cobalt protoporphyrin-IX (CoPPIX) or heme, as well as transduction of cells with lentiviral vectors coding for HO-1 led to growth reduction [111]. Finally, a significant decrease in cell proliferation was observed in hemin-treated or HO-1 overexpressing prostate carcinoma [91]. It must be stressed, however, that employ-

ment of complementary methods, such as siRNA application that allow for specific HO-1 inhibition, is necessary to get the conclusive results, because pharmacological modulators of HO-1 activity may exert many HO-1 independent effects [112].

Of importance, proproliferative effects of HO-1 have been confirmed *in vivo*. Most experiments supported the permissive role of HO-1 in tumor growth, although in some cases, for instance in some prostate cancers, HO-1 overexpression might reduce the mitotic index [91]. Usually, pharmacological or genetic induction of HO-1 facilitated tumor progression, as manifested by bigger volumes of nodules or by more numerous cancer cells. Such a correlation has been demonstrated in angioma [77], and melanoma [108]. In accordance, reduced HO-1 expression resulted in decreased growth of hepatoma, sarcoma [69], pancreatic cancer [76, 113], angioma [77], lung cancer [114], hepatocellular carcinoma [70], and some prostate cancer lines [66].

Exact mechanisms responsible for modulation of cell cycle by HO-1 still require explanation. Nevertheless, microarray analysis of murine melanoma transcriptome showed that augmented proliferation of HO-1 overexpressing cells can be associated with lower levels of p21, the major negative regulator of cell cycle, and with downregulation of Mdm2, the inhibitor of p53 [108]. Apart from other known functions, the p53 plays a role in regulation of oxidative stress, through activation of numerous genes involved in production of ROS. Concurrently, ROS can further activate p53, which may lead to upregulation of cytoprotective and DNA repairing genes [115]. One of such antioxidative responders is HO-1, which contains the p53 responsive element in the promoter sequence [115].

Low levels of ROS can facilitate the reaction to mitogens [116], and HO-1 has been identified as a mediator of ROS-augmented proliferation in the breast cancer cells [117]. Moreover, silencing the HO-1 gene in HRPcA cells, using siRNA or pharmacological inhibitor OB-24, decreased oxidative stress, reduced MAPK (ERK1/2 and p38) signaling, and led to inhibition of prostate cancer proliferation both *in vitro* and *in vivo* [66]. Interestingly, HO-1 was stimulated in a p53-dependent manner in irradiated mice, and irradiation dose required to its induction was higher than that for upregulation of p21. Furthermore, HO-1 response occurred later than the stimulation of p21. It may suggest the existence of a negative feedback loop between the activation of p53 by ROS and the p53-dependent expression of HO-1: increased HO-1 would reduce intracellular oxidative stress leading to diminution of p53 activity and consequently to a decrease in p53 target genes, including p21 [115].

The cyclin-dependent kinase (Cdk) inhibitor p21 is often downregulated in tumors, and its binding to Cdk2 blocks the progression of cell cycle [118]. Importance of p21 attenuation has been already revealed in clinical studies, e.g. in cutaneous malignant melanoma [119]. HO-1 activation attenuates p21 expression in different cell types, including endothelial cells, melanoma, and colon carcinoma [108, 120, 121]. However, there are some studies showing that overexpression of HO-1 results in a significant increase in the levels of p21 in tumors [122, 123]. Important aspect in tumor biology is also a subcellular localization of p21. Recently, 4-methylnitrosamino-1-(3-pyridyl)-1-butanone (NNK), the

most potent carcinogen of cigarette smoke, has been shown to up-regulate the HO-1 expression and simultaneously induce proliferation of lung cancer cells. Noteworthy, such a treatment promoted the nuclear localization of p21. In accordance, troglitazone (activator of peroxisome proliferator-activated receptor- γ , PPAR γ) that is known to arrest the growth of lung cancer, increased the cytosolic fraction of p21 [120]. Nuclear-localization of p21 was reported to correlate with progression of tumors and worsen survival of patients [121]. Importantly, p21 nuclear translocation in cells treated with NNK was completely blocked by HO-1 inhibitor, zinc protoporphyrin-IX (ZnPPiX) [120].

Finally, it was observed that ZnPPiX may reduce activity of transcriptional factors, Sp1 (*Stimulatory protein-1*) and Egr1 (*Early growth response factor-1*), and thereby inhibit the transcription from the cyclin-D1 promoter, suggesting that HO-1 may upregulate the cyclin-D1 generation [124]. This supposition has been confirmed recently in fibroblasts isolated from HO-1 deficient mice, where cyclin-D1 expression was significantly lower than in their wild-type counterparts [Was *et al.*, paper under revision]. Thus, attenuation of D1-dependent activities can be also one of mechanisms underlying the HO-1-dependent regulation of cell proliferation.

There is no single mediator of the proproliferative action of HO-1. In endothelial cells the most important seems the generation of CO and downregulation of p21 [98, 125]. Confusingly, in vascular smooth muscle cells (VSMC) p21 was identified as a protein mediating an HO-1-dependent inhibition of proliferation, and antiproliferative potential of HO-1 was significantly reduced in VSMC obtained from p21 null mice [102]. Additionally, treatment of wild-type VSMC with hemin downregulated the expression of cyclin-D1 and upregulated that of p21, the effects opposite to those observed in endothelium and many cancer cells [126]. Similarly as in VSMC, the positive correlation of HO-1 and p21 was demonstrated in epithelial cells [44], papillary thyroid carcinoma [122], gastric cancer cell line [123], pancreatic [127], and hepatic stellate cells [128]. Again, the postulated effector molecule in antiproliferative action of HO-1 was CO which stimulated p38-MAPK through activation of sGC and elevation of cGMP. Constitutive activity or overexpression of p38-MAPK can result in permanent cell-cycle arrest and premature cell senescence [129].

One can also hypothesize that CO may influence cell proliferation through caveolin-1 (cav-1) [130], the main structural component of caveolae [131]. Increased expression of cav-1 is characteristic for aged animals and for fully differentiated or senescent cells [132], while its generation is reduced in tumors [133]. In VSMC and fibroblasts the CO activates p38 β -MAPK isoform, which leads to upregulation of cav-1 and inhibition of proliferation [130]. Interestingly, HO-1 has been shown to localize in caveolae, and cav-1 seemed to negatively regulate the enzymatic activity of HO-1 in pulmonary endothelium [134].

The second effector product in antiproliferative activity of HO-1 can be biliverdin. This compound is very effectively converted to bilirubin, which in turn can inhibit expression of cyclin A, D1, and E, as well as cdk-2, and block hyperphosphorylation of retinoblastoma protein (Rb) [103]. Additionally, bilirubin inhibits phosphorylation of p38-MAPK

[129] and of ERK1/2, the kinases involved in regulation of cell proliferation [135]. It should be kept in mind, however, that p38-MAPK may be both a permissive [100] and inhibitory regulator of cell cycle progression, depending on the cell context and/or p38-MAPK isoform active [136].

Finally, reduction in cellular heme content and resulting iron shortage in cells with a very high levels of HO-1 may also lead to inhibition of tumor growth [137]. Iron is essential for proliferation, especially in rapidly dividing cancer cells [138]. Iron chelators such as desferrioxamine (DFO) already showed anti-tumoral potential in many tumors *in vitro* and in limited number of clinical trials [139]. Thus, DFO inhibited growth of human myeloid leukemia and neuroblastoma cell lines [140], and displayed antitumor activity in neuroblastoma patients [141]. Finally, DFO significantly improved effectiveness of gemcitabine treatment in mice bearing the pancreatic cancer [142].

Hitherto, data on role of HO-1 end-products in proliferation of cancer cells are very limited and still confusing. Thus, CO exposure led to a tendency towards increased division of pancreatic cancer cells [142], whereas it blocked proliferation of prostate cancer cell lines, and the latter effect was dependent on inhibition of Rb phosphorylation [143]. In turn, application of both biliverdin and ferrous histidinate resulted in increased pancreatic cancer cell proliferation [142]. The well-documented, but conflicting reports regarding effects of HO-1 on cell cycle suggest that the final output does not depend on HO-1 alone, but is rather determined by the reciprocal balance of several players, including HO-1, p21, p53, ferritin, caveolin-1, and possibly other, still not recognized factors (Fig. 2).




Fig. (2). Schematic demonstration of the role of HO-1 in regulation of cell proliferation in tumors. N - nuclear, C - cytoplasmic.

ROLE OF HO-1 IN TUMOR DIFFERENTIATION

The role of HO-1 in cell differentiation is not well recognized. Nevertheless, there are some data showing the

influence of HO-1 on maturation of osteoclasts, osteoblasts, dendritic cells, and adipocytes. It was demonstrated that activation of HO-1 prevents differentiation of osteoclasts [144], and inhibits the maturation of primary osteoblasts, as illustrated by attenuated mineralized bone nodule formation, reduced alkaline phosphatase activity and decreased expression of several differentiation markers such as alkaline phosphatase, osteocalcin, and RUNX2 [145]. Similarly, HO-1/ferritin system prevented inorganic phosphate-mediated calcification and osteoblastic differentiation of human smooth muscle cells [146], and depressed the maturation of antigen-presenting cells into monocyte-derived dendritic cells [147]. On the other hand, upregulation of HO-1 shifted the balance of mesenchymal stem cells (MSC) differentiation in favor of osteoblast lineage, whereas HO-1 inhibition drove them towards adipogenesis [148].

The role of HO-1 in cancer cell differentiation is even less established. However, changes in HO-1 expression were shown in erythroleukemia, in which the induction of erythroid maturation by dimethyl sulfoxide (DMSO) was coupled with a rapid decline in mRNAs for HO-1, heat-shock protein-70 (Hsp-70), and nonspecific δ -aminolevulinate synthase (ALAS) [137]. Changes in HO-1 expression were also found in the differentiating myeloid leukemia cell line, which can develop bidirectionally i.e., to erythrocytes after treatment with hemin and to monocytes in response to 12-O-tetradecanoylphorbol 13-acetate (TPA). Interestingly, inhibition of HO-1 activity by SnPPiX suppressed only the TPA-induced maturation to monocytes. Therefore, HO-1 could be suggested as a directional switch and permissive enzyme in monocytic differentiation program of myeloid leukemia [149].

Effects of HO-1 on cell differentiation seem to be associated with redox signaling. Recent studies suggest that production of ROS can be essential for neuronal maturation. Elevation of ROS in haematopoietic progenitors was also found to trigger differentiation into mature blood cell types, through a signaling pathway that involves JNK and FoxO activation [150]. Similarly, hemin-induced ROS may activate AKT and ERK signaling pathways, thereby promoting a significant increase in neovessel formation and expression of endothelial markers in endothelial progenitor cells (EPC) [151]. It can also potentiate pro-differentiating potential of BML-210, a histone deacetylase inhibitor, in human leukemia cell lines [152]. Increased activity of HO-1, through degradation of hemin and decrease in ROS generation, might attenuate these effects. It appears that HO-1 might protect cells from oxidative stress-related injury during differentiation but, if expressed at a high level, the enzyme by effective removal of ROS might inhibit cell differentiation. Nevertheless, such a postulate is still mostly speculative and needs to be supported by strong experimental data (Fig. 3).

ROLE OF HO-1 IN TUMOR CYTOPROTECTION AND APOPTOSIS

HO-1 is activated by many stress-inducing factors and this response may afford protection from oxidative damage and reduce the rate of apoptosis in different cells [153-155]. The most convincing illustration of cytoprotective properties




Fig. (3). Schematic demonstration of the role of HO-1 in regulation of cell differentiation in tumors.

of HO-1 come from the studies on HO-1 knockout mice, where HO-1 deficiency leads to severe oxidative stress, increased lipid peroxidation, enhanced oxidative damage in cardiovascular system, and progressive chronic inflammation in the kidney and liver. Moreover, fibroblasts isolated from HO-1^{-/-} individuals demonstrate augmented production of ROS and reduced viability when exposed to oxidants such as hemin or H₂O₂ [156]. Of importance, cytoprotective effects of HO-1 have been confirmed in various diseases, including atherosclerosis, ischemia-reperfusion and acute renal injury, toxic nephropathy, cisplatin nephrotoxicity or endotoxic shock [157].

Cytoprotective and antiapoptotic properties of HO-1 were also demonstrated in cancer cells. Accordingly, pharmacological or genetic activation of HO-1 significantly improved survival of hepatoma [69, 70], thyroid carcinoma [122], chronic myelogenous leukemia [78], gastric carcinoma [123], colon carcinoma [158], melanoma [108], lung carcinoma [159, 160], and glioma [161], whereas HO-1 inhibition reduced viability of colon carcinoma [158], acute myeloid leukemia [162], and hormone-refractory prostate cancer [66].

Of importance, expression of HO-1 is additionally elevated in response to anticancer treatments [76, 85-87, 89], thereby augmenting the resistance of cancer cells [78, 86, 163]. On the other hand, the contribution of HO-1 in tumor cytoprotection allows for increasing the sensitivity of cancer cells to therapeutic procedures by inhibition of HO-1 expression and/or activity. The feasibility of such an approach has been revealed in cultured tumor cells of pancreatic cancer subjected to radiotherapy and chemotherapy [76], in colon carcinoma [164], lung carcinoma [160], and chronic myelogenous leukemia [78, 165] treated with chemotherapy, or in T-cell leukemia [166], melanoma [167], and urothelial carcinoma [88] exposed to photodynamic therapy. Results of *in vitro* experiments have been confirmed in animal models. Particularly, administration of ZnPPIX or ZnPPIX-poly-

ethylene glycol (ZnPPIX-PEG) led to enhanced apoptosis in rat hepatoma and sarcoma or in murine lung tumors [168], while HO-1 siRNA treatment significantly increased the rate of apoptosis in murine hepatocarcinomas [70]. Finally, it was reported that most nasopharyngeal carcinoma patients with undetectable expression of HO-1 were responsive to radiotherapy, whereas almost 60% of those with tumors positive for HO-1 showed high resistance to the treatment [169]. These findings may indicate the HO-1 as a helpful marker in classifying the patients to radiotherapy, and may suggest the inhibition of HO-1 as a new sensitizing strategy for nasopharyngeal carcinoma.

Mechanisms underlying the cytoprotective effects of HO-1 in tumor cells are still not fully understood. One possible pathway is removal of free heme, a prooxidant and important modulator of signal transduction [170]. Heme can be harmful to the cells, but in the same time it directly regulates the activity of Nrf2 and Bach1 transcription factors [171, 172], inducing the expression of not only HO-1 but the whole set of cytoprotective and detoxifying genes [173, 174]. Because of such a complex response it is not trivial to assess the real contribution of HO-1 in cell protection. For example, apoptosis induced by imatinib (ABL tyrosine kinase inhibitor) in leukemia cell lines was suppressed following the treatment with hemin or with δ -aminolevulinic acid (δ -ALA), the heme precursor [134]. Furthermore, hemin selectively counteracted the imatinib-induced repression of antiapoptotic Bcl-2a and Bcl-2b genes [175], and decreased the sensitivity of cells to anthracyclins [163]. The cytoprotective effects were reduced by Nrf2 siRNA, indicating the significance of this transcription factor, but without explanation the role of HO-1 [163]. Nrf2 was also activated in leukemic cells in response to TNF, what augmented the cell resistance to TNF-induced apoptosis. Here, the application of HO-1 siRNA, which restored cell sensitivity, evidenced the role of HO-1 in Nrf2-mediated cytoprotection [162]. Nrf2 signaling pathway was also responsible for induction of HO-1 and protection against NO-induced damage in colon carcinoma cells [174]. Similarly, Nrf2 mediated the HO-1 upregulation and resulting cytoprotection in monocytes treated with epigallocatechin 3-gallate (EGCG), the anticancerogenic polyphenol found in green tea [176, 177], and in neuronal cells exposed to NO [178]. Importantly, Nrf2 is not only an upstream inducer of HO-1, but can be also upregulated in response to HO-1 activation, the effect mediated by CO and PI3K pathway, and this positive feedback loop might be one of potential mechanism of HO-1-induced cytoprotection [179].

Another possible explanation for cytoprotective and antiapoptotic activity of HO-1 is the increase in cellular biliverdin and bilirubin levels. Both compounds are known to inhibit hepatocyte apoptosis *in vitro* [180] and *in vivo* [181]. Bilirubin can also protect keratinocytes against nicotine-induced cytotoxicity [182], and cardiomyocytes against doxorubicin-induced death by increasing Bcl-2 and decreasing Bax expressions [183]. In accordance, supplementation of cultured cells with biliverdin or bilirubin mimicked the cytoprotective action of HO-1, as shown for hepatoma [168] and colon carcinoma [158]. Finally, the effect of HO-1 inhibition on ROS elevation in colon carcinoma was completely abrogated, if cells were supplemented with bilirubin [184]. It must be kept in mind, however, that

unconjugated bilirubin can be toxic, as demonstrated in cervical adenocarcinoma [185] or colorectal cancer [186].

Cytoprotection can be also mediated by HO-1-derived CO [168], which act through activation of both p38-MAPK [130] and sGC [187]. Importantly, CO was not only shown to block the release of mitochondrial cytochrome c, the crucial event in induction of apoptosis, but also to inhibit expression of proapoptotic p53 [164]. Exogenous CO may increase the viability of different cell types, including fibroblasts [101], endothelium [188], hepatocytes [189], and pancreatic β -cells [190]. Like biliverdin, CO can inhibit the doxorubicin-induced apoptosis in cardiomyocytes [183, 191]. However, these effects appear to be cell-type specific, as no CO-exerted protection was found in colon carcinoma, gastric cancer cells, and chronic myelogenous leukemia [78, 158].

Finally ferritin, induced by many HO-1 activators, and then by iron released from heme, can contribute to HO-1 mediated cytoprotection. Indeed, in malignant oral keratinocytes exposed to NO, HO-1 reduced cell mortality by upregulation of iron regulatory proteins (IRP1 and IRP2), transferrin receptor (TfR), and ferritin [192]. Moreover, susceptibility of tumor cells to oxidants was inversely correlated with ferritin protein levels [193]. For example, reduced expression of ferritin resulted in enhanced sensitivity to oxidative stress and apoptosis in melanoma [194] and T-cell lymphoma [195]. Accordingly, increased expression of ferritin was suggested as a mechanism responsible for the protective effect of hemin in leukemia cells subjected to photodynamic therapy [196]. Evaluations of ferritin levels in tumor tissues generated, however, much more complex and perhaps cell-type specific picture: in some tumors, such as colon or breast cancers, ferritin was upregulated when compared with healthy tissues, in others, such as liver cancer, it was decreased [197]. Interestingly, immunohistochemical analysis of human melanoma showed that ferritin expression in metastatic lesions was significantly higher than in primary nodules [194].

It must be kept in mind, however, that induction of HO-1 is not always adequate to protect the cells. As demonstrated in breast carcinoma and B-lymphoblasts HO-1 failed to defend the cells from chemotherapy-induced apoptosis [198]. Moreover in some experiments, overexpression of HO-1 enhanced oxidative damage to mitochondria and led to augmented apoptosis in breast cancer cells, astroglia [107], and VSMC [199]. Thus, although HO-1 up-regulation can generally confer antioxidant protection, under certain conditions it may promote ROS accumulation within the mitochondria and other cellular compartments [66]. Nevertheless, the vast majority of studies performed in different tumor cells, such as hepatoma [168], colon adenocarcinoma [87, 158, 168], thyroid carcinoma [122], gastric carcinoma [123], myelogenous leukemia cells [78], and melanoma [108] strongly suggest that HO-1 is a powerful cytoprotective and antiapoptotic enzyme, which improves survival of cancer cells subjected to different therapeutic treatments (Fig. 4).

ROLE OF HO-1 IN TUMOR ANGIOGENESIS

Angiogenesis, the formation of new blood vessels from preexisting ones, plays a crucial role in growth and spreading

of tumors. The notion that tumor progression can be accompanied by increased vascularity was first described in




Fig. (4). Schematic demonstration of the role of HO-1 in regulation of cell death in tumors.

the beginning of the 20th century by the German pathologist Rudolf Virchow [200], and then by others [201, 202]. In 1971, a theoretical paper by Judah Folkman postulated that the growth of tumors depends on the induction of new blood vessels, and that inhibitors of angiogenesis might be useful for treating human cancers [203]. Of importance, apart from its cytoprotective actions, HO-1 has been recognized as a proangiogenic enzyme. Thus, one can speculate that these properties may further support the role of HO-1 in tumor progression.

The role of HO-1 in angiogenesis has been shown in different experimental settings. So, endothelial cells isolated from HO-1-deficient mice produced less VEGF than did their wild-type counterparts, and their response to exogenous VEGF and bFGF was weaker [204]. On the other hand, genetic overexpression of HO-1 in endothelial cells increased production of VEGF and facilitated VEGF-induced activities, namely proliferation, migration, formation of capillary-like tubular structures in a Matrigel matrix, and outgrowth of capillaries from endothelial spheroids in a collagen gel [98]. In accordance, HO-1 has been demonstrated to stimulate expression of VEGF also in vascular smooth muscle cells [102] and keratinocytes [205]. At least in microvascular endothelium and vascular smooth muscle cells, this effect appeared to be mediated by CO [98, 125,

206]. Importantly, the results of experiments performed *in vitro* have been confirmed *in vivo* in the rat ischemic hindlimb model or in skin wounds of diabetic mice, where HO-1 gene transfer increased angiogenesis [207].

The influence of HO-1 on tumor vascularization was suggested first by observations that HO-1 expression in tumor infiltrating macrophages correlates with increased vascular density, as shown in human gliomas and vertical growth melanomas [74]. Then it was reported that HO-1 accelerates pancreatic cancer growth by promoting tumor angiogenesis in mice [113]. Significance of HO-1 has been confirmed in experiments where targeted knockdown of HO-1 by siRNA or inhibition of HO-1 activity by SnPPIX impaired the cell survival and VEGF expression, as shown in endothelioma [77], lung carcinoma [114], and in tumors formed by transformed fibroblasts [208]. Moreover, HO-1 inhibition interfered with tumor neoangiogenesis in hepatocellular carcinoma *in vivo* [70]. In concordance, activation of HO-1 in glioma cells with hemin or in breast carcinoma with 15-deoxy-delta-12,14-prostaglandin J₂ (15d-PGJ₂) was accompanied by elevation of VEGF, the effect blocked by pretreatment of cells with HO-1 inhibitor [82]. Finally, in 15d-PGJ₂-stimulated breast cancer cells the expression of matrix metalloproteinase-1 (MMP-1), crucial for growth of capillaries, was HO-1 dependent and facilitated by iron released from HO-1-catalyzed heme degradation [209].

A proangiogenic role of HO-1 was directly evidenced in murine melanoma [108]. Interestingly, HO-1 overexpression in the melanoma cell line did not lead to increased production of VEGF *in vitro*, and VEGF did not contribute significantly to the increased angiogenic potential of media harvested from such cells. However, HO-1 overexpressing melanoma cells inoculated subcutaneously into mice, gave more vascularized tumors with higher content of VEGF, what may suggest an indirect influence of HO-1 present in cancer cells on production of VEGF in stroma, especially in macrophages. This effect could be possibly mediated by diffused CO [108].

It seems that HO-1 might contribute to proangiogenic effects of thymidine phosphorylase (TP), the enzyme that is often elevated in the blood of cancer patients, and has been implicated in pathophysiological angiogenesis. TP stimulates migration of endothelial cells [210] and formation of capillaries both *in vitro* and *in vivo* [74, 211, 212]. It directly upregulates HO-1, and coexpression of these two proteins was detected in advanced melanoma tumors [74], and bladder carcinoma [211]. TP has been shown to induce oxidative stress in tumor cells, and subsequently the release of angiogenic factors, such as VEGF, IL-8 and MMP-1 [213]. Interestingly, some effects of TP, such as regulation of VSMC proliferation, were blocked both by inhibitors of TP and HO-1 [214].

Similarly, it was found that hepatocyte growth factor (HGF), a multifunctional cytokine of mesenchymal origin [215], promotes hepatocarcinogenesis and stimulates angiogenesis directly or indirectly through upregulation of VEGF [216]. In hepatoma cells it can activate hypoxia-inducible factor-1 α (HIF-1 α), major proangiogenic transcription factor, *via* stimulation of ROS generation, which induce in turn the HO-1 expression [217]. The major proangiogenic and anti-

apoptotic effects of HGF are similar to that of HO-1. Interrelation of these two pathways may potentially play an important role in tumor progression. This hypothetical supposition needs, however, an experimental verification (Fig. 5).




Fig. (5). Schematic demonstration of the role of HO-1 in regulation of angiogenesis in tumors.

Tumor vascularization is a complex process, relying not only on sprouting angiogenesis, but also on adult vasculogenesis, in which bone marrow-derived EPC home to the tumor site, differentiate into mature endothelial cells and incorporate into growing tumor vessels, or stimulate local angiogenesis through production of proangiogenic growth factors [218-220]. One of the key factors governing the EPC homing is VEGF, whose expression and activity is influenced by HO-1. The second very important agent is stroma cell-derived growth factor-1 (SDF-1), a chemokine that plays a major role in migration, recruitment, and retention of EPC to ischemic or injured tissues and contributes to neovascularization.

Analyses of intracranial murine gliomas evidenced the increased production of SDF-1 in the vasculogenic tumors. Furthermore, enforced expression of SDF-1 augmented, whereas blocking the SDF-1 receptor (C-X-C motif receptor-4, CXCR4) reduced homing and engraftment of vascular progenitors [221]. Moreover, SDF-1 may promote the retainment of BM-derived pericytes in close association with perfused, functional tumor vessels. It was convincingly illustrated in experiments, where SDF-1 cDNA was delivered using adenoviral vectors to the tumors in which VEGF was inhibited. SDF-1 overexpression led to augmented tumor growth, increased number of large, lumen-bearing vascular structures, and enhanced vessel pericyte coverage [222]. These observations confirm its role in tumor vascularization.

No data are available on potential contribution of HO-1 in SDF-1-induced activities in tumors. Such an influence, however, is highly possible, in light of recent experiments demonstrating an important role of HO-1 in proper response of endothelial cells or EPC to SDF-1. Thus, SDF-1 activates HO-1 expression both in mature endothelium and in EPC. Pharmacological inhibition or genetic ablation of HO-1 precludes the SDF-1-mediated migration, endothelial tube formation *in vitro* and sprouting of capillaries from aortic rings *ex vivo*, effects that can be restored by CO. The requirement of HO-1 for proper SDF-1 action was confirmed *in vivo* in Matrigel-plug, wound-healing, and retinal ischemia models of angiogenesis and vasculogenesis [223].

Additionally, protection of ischemic myocardium resulting from AAV overexpression of HO-1 correlated with higher production of VEGF and SDF-1, enhanced vascularization, and increased number of progenitor cells recruited to the infarcted area. In accordance, inhibition of VEGF and SDF-1 by neutralizing antibodies significantly attenuated HO-1-mediated neovascularization and protection, evidencing that VEGF and SDF-1 are the downstream effectors of HO-1 activity [224, 225]. Finally, systemic HO-1 induction led to elevated serum levels of VEGF and SDF-1, increased number of circulating EPCs, and improved re-endothelialization of denuded vessels [112]. Therefore, understanding the role of HO-1 in SDF-1-mediated tumor vascularization is urgently needed (Fig. 5).

ROLE OF HO-1 IN METASTASIS

Angiogenesis is the key event not only in tumor growth but also in metastasis, a multistep process that includes modulation of cell adherence, degradation of surrounding extracellular matrix, migration, proliferation at a secondary site, and neovascularization of newly formed nodules. Metastases spread in three ways - by local extension from the tumor to the surrounding tissues, through the bloodstream to distant sites, or through the lymphatic system to neighboring or distant lymph nodes. The progression of tumor to invasive and metastatic form correlates with poor clinical outcome [226].

Because of proangiogenic properties of HO-1, one can expect that it also would facilitate metastasis. Such a supposition has been confirmed in several models. Namely, melanoma or pancreatic cancer cells engineered to overexpress HO-1 formed much more nodules in the lungs of mice after intravenous injection than their wild-type counterparts [108, 113], whereas pharmacological inhibition of HO-1 completely inhibited metastasis [113]. Accordingly, silencing the HO-1 gene in prostate carcinoma reduced cell invasion *in vitro*, and inhibited growth of primary and metastatic tumors *in vivo* [66]. Finally, clinical data demonstrate that high level of HO-1 expression is associated with lymph node metastasis in oral squamous cell carcinoma patients [227].

However, data on the role of HO-1 in tumor metastasis are inconsistent and may reflect the cell type-specific or development-stage specific effects. For example in MCF-7 breast cancer cell line, HO-1 inhibited the invasion induced by TPA, what resulted from HO-1-dependent decrease in ROS generation, and CO-mediated reduction in MMP-9

expression [228]. Inhibition of MMP-1, MMP-2, MMP-7, and MMP-9 by HO-1-derived CO has also been reported in other cell types [229-231], although MMPs regulation, especially the influence of oxidative stress, depends very much on cellular context [231]. Additionally, in some experiments performed in prostate cancer, the overexpression of HO-1 reduced cell migration and invasiveness *in vitro* [91], in contrast to the other published observations [66]. Finally, in colorectal [233, 234] and oral carcinomas [71, 235, 236], HO-1 was suggested as a marker of lower risk of metastasis. Thus, further studies are necessary to clarify its role. The important and hitherto underestimated aspect is also a distinction the effects of HO-1 expressed in neoplastic and stroma cells.

ROLE OF HO-1 IN TUMOR-RELATED INFLAMMATION

HO-1 is regarded as a potent anti-inflammatory enzyme. Many reports convinced that activation of HO-1 weakens inflammation and may result in immunosuppression [237]. The significance of HO-1 as an immunomodulator has been elegantly illustrated by studies performed in HO-1^{-/-} mice showing that such animals develop progressive inflammatory disease characterized by splenomegaly, lymphadenopathy, leukocytosis, as well as hepatic and renal inflammation [156]. Moreover, HO-1 deficiency lead to strongly enhanced generation of proinflammatory cytokines, including IL-1 β , interferon γ (IFN γ), TNF, and IL-6, so in general, Th1-weighted shift in immunity [238].

Many data suggest that overexpression of HO-1 may defend tissues from immune-mediated injury, either through protection against oxidative damage or via local immunomodulation [47, 61]. The exact mechanisms responsible for anti-inflammatory functions of HO-1 have not been fully elucidated. However, the signaling induced by CO combined with antioxidant properties of biliverdin/bilirubin and ferritin-mediated sequestration of iron could all contribute to this suppression.

CO has been described as an anti-inflammatory mediator in several models of inflammation and tissue injury. Thus, in lipopolysaccharide (LPS)-stimulated macrophages, it decreased the generation of inducible nitric oxide synthase (iNOS)-derived NO and production of TNF, IL-1 β , MIP-1 β , or IL-6 [61, 239], and these effects were exerted by interfering with AP-1 activity via a c-Jun N-terminal kinase (JNK) pathway [240]. CO reduced also the production of granulocyte macrophage colony-stimulating factor (GM-CSF), which is known to increase the secretion of proinflammatory mediators and promote the maturation of macrophages and neutrophils [241, 242]. Conversely, exposure to CO can increase the synthesis of anti-inflammatory IL-10 both *in vitro* and *in vivo* [61, 231].

Recent data suggest that HO-1/CO may reduce production of proinflammatory cytokines through augmenting the interaction between cav-1 and toll-like receptor-4 (TLR4). Cav-1 inhibits inflammation through direct binding of TLR4, which prevents TLR4 association with MyD88 and TRIF, and blocks downstream activation of NF κ B pathway. Cav-1 was also shown to activate p38 MAPK [243], which

was required for translocation of HO-1 to caveolae in LPS-treated macrophages [244] (Fig. 6).




Fig. (6). Schematic demonstration of the role of HO-1 in regulation of inflammatory-related pathways in tumors.

Similarly, inflammatory response can be diminished by biliverdin and bilirubin. For example, they blunt the inflammatory cascade in a model of rodent sepsis [245], and reduce the adhesion of leukocytes to vascular endothelium via inhibition of NFκB-dependent E-selectin, intracellular adhesion molecule-1 (ICAM-1), and vascular cell adhesion molecule-1 (VCAM-1) expressions [246]. The importance of biliverdin/bilirubin was also confirmed in HO-1-mediated reduction of monocyte chemotaxis [247]. Interestingly, anti-inflammatory effects of far infrared (FIR) therapy in human endothelium were reversed by pharmacological or genetic inhibition of HO-1, and could be restored by bilirubin [248]. Finally, unconjugated bilirubin can inhibit the initial step in the classical pathway of complement cascade activation, an important effector mechanism of immune system [249] (Fig. 6).

Tumors are infiltrated by many types of leukocytes, including macrophages, dendritic cells, neutrophils, mastocytes, eosinophils, natural killer (NK) cells, T and B lymphocytes, with macrophages being often the most abundant population [250]. Microenvironmental cytokines may not only selectively promote a Th1 or Th2 immune response, but also polarize macrophages toward the M1 or M2 phenotype [251]. The phenotype of tumor-associated macrophages (TAM) usually resembles that of M2 cells, with pro-tumor functions that include inhibition of Th1 immune response, and production of growth factors, such as epidermal growth factor (EGF), IL-6, IL-8, VEGF, platelet-derived growth factor (PDGF), or transforming growth factor-β (TGF-β). TAM can also induce the degradation and remodelling of extracellular matrix (ECM) via the expression of MMPs [252, 253]. In turn, lymphocytes may play a dual role in

tumor progression. Thus, tumor-specific cytotoxic T lymphocytes are able to eliminate tumor cells by direct or antibody-dependent cytotoxicity. However, the recruitment of regulatory T lymphocytes (Treg) attenuates the immune response and may neutralize the cytolytic potential of cytotoxic effectors [251], while tumor microenvironment supports the conversion of naive T lymphocytes into Foxp3⁺/CD25⁺ Tregs [254, 255]. It is supposed that activation of Treg cells is one of the main if not the main barrier precluding effective cancer immunotherapies [256, 257].

Inflammation is a key factor in tumor growth and metastasis. Therefore, one can expect that its regulation by HO-1 may affect the progression of tumors. Unfortunately, the experimental data illustrating such an influence are very limited. Interestingly, HO-1 can contribute to macrophage polarization toward M2 phenotype [258] that resembles that of protumorigenic TAMs. It also suppresses the T cell- and NK cell-mediated effector functions [259]. In accordance, elevated HO-1 activity may lead to clonal deletion of CD4⁺ T cells, resulting in a specific immunomodulation, as illustrated by prolongation of transplanted organ survival [260]. Moreover, it has been demonstrated that HO-1 expression is significantly higher in CD4⁺CD25⁺ Tregs than in CD4⁺CD25⁻ population [47], and induction of FoxP3 followed by stimulation of regulatory activity in CD4⁺CD25⁻ T lymphocytes correlates with HO-1 upregulation [261].

Recent evidence suggests that expression of HO-1 in dendritic cells may further strengthen the tolerogenic effects [262]. It blocks both their maturation and proinflammatory




Fig. (7). Schematic demonstration of the role of HO-1 in regulation of actions of inflammatory cells in tumors.

functions but preserves ability to the IL-10 synthesis [231, 263]. Thus, HO-1-expressing dendritic cells may activate suppressor T lymphocytes, thereby boosting suppression of local immune responses. Accordingly, adenoviral transduction of dendritic cells with HO-1 in rats receiving heart transplants led to the long-term allograft survival accompanied by an inhibition of cellular antigrraft immune responses [264]. Of interest, naturally occurring CD8⁺ T lymphocytes specific for HO-1 have been shown to suppress cellular immune responses. Such cells were detected *ex vivo* and *in situ* among T lymphocytes from malignant melanoma, renal cell carcinoma, and breast cancer patients. HO-1-specific T cells isolated from the peripheral blood blocked the cytokine release, proliferation, and cytotoxicity of other immune cells. Notably, the inhibitory effect of HO-1-specific T lymphocytes was far more prominent than that of conventional CD4⁺CD25⁺ Tregs [265]. Thus, HO-1 might be considered as a potential target for accessory strategies facilitating cancer immunotherapies. However, the molecular mechanism(s) responsible for the anti-inflammatory and immunomodulatory effects of HO-1 in tumors have not been elucidated and require further investigations (Fig. 7).

EFFECT OF HO-1 POLYMORPHISMS IN CANCER PATIENTS

Analysis of clinical data and comparison of HO-1 allele frequency distribution in healthy people and cancer patients can suggest the role of HO-1 in cancer development and anticancer therapy. Importantly, the efficacy of HO-1 pathway can vary in human population due to variable number of (GT)_n repeats, ranging from 11 to 40, in the proximal part of the HO-1 promoter [266]. Shorter fragments of repeats (S alleles) are related to higher transcription rates. It has been shown that lymphoblastoid cell lines derived from human subjects possessing S/S genotype showed much higher activity of HO-1 enzyme after H₂O₂ stimulation than did their L/L counterparts, and only S/S promoters provided protection of lymphoblastoid cells from H₂O₂-induced apoptosis [267]. We have demonstrated that primary human endothelial cells carrying the S allele display a higher expression and activity of HO-1 both in control conditions and in response to some oxidants or inflammatory mediators. Moreover, it is associated with the augmented resistance to oxidative stress, more favorable GSH:GSSG ratio, reduced synthesis of proinflammatory cytokines, and increased cell proliferation [268]. Finally, constitutive expression of HO-1 protein in urothelial cancers was associated with the presence of S alleles [88].

Interestingly, recent reports indicate that lack of short alleles and resulting lower activity of HO-1 can coincide with a higher risk of cancers. The frequency of L alleles was significantly higher in male smokers with lung adenocarcinoma than in control subjects [269]. Furthermore, the longer (GT)_n repeats in the HO-1 promoter were linked to the increased risk of areca-induced oral squamous cell carcinoma (OSCC) in males [270]. The frequencies of L-alleles were also significantly higher in esophageal squamous cell carcinoma patients (especially those heavy drinking) than in controls [271]. In turn, the higher risk of breast cancer was observed among postmenopausal women with high iron intake, carrying the HO-1 LL or LM genotypes [272].

Additionally, patients possessing both NFκB ins (resulting in increased activity of the factor) and HO-1 L alleles showed significant risks for various subsets of OSCC, including those of advanced stage with node metastasis [236]. Finally, the long (GT)_n sequence in the HO-1 promoter coincided with a higher frequency of gastric adenocarcinoma, whereas the M alleles were associated with a lower frequency of lymphovascular invasion in tumors [273].

However, the effect of HO-1 promoter polymorphism may depend on the type of tumor. For example, the calculated risk for acquiring primary malignant melanoma in L-allele carriers was twofold lower than in those having S/S genotype [274]. In turn, the more active variants of HO-1 and iNOS promoters were associated with increased risk of gastric cancer in women [275].

ROLE OF HO-1 IN CARCINOGENESIS

As mentioned above HO-1 can protect tumor cells against variety of harmful compounds (e.g. oxidants, chemotherapeutics), favoring tumor progression. On the other hand, it inhibits actions of dangerous stimuli, including carcinogens, in healthy tissues. Therefore, the fundamental question is whether HO-1 takes part in healthy - pre-malignant - malignant transformation or rather blocks this process. To our best knowledge, no studies address the problem directly. Although, it might be speculated that HO-1 could enhance the resistance of cells to carcinogenesis, especially that one induced by ROS or compounds producing ROS, like heme. Genotoxic actions of heme were shown in the human colon cell line and primary human colonocytes [276], whereas carcinogenic only in the colon [277]. On the other hand, antiapoptotic properties of HO-1 could be responsible for preservation of oncogenic mutations and acceleration of tumor transformation.

Importantly, the influence of HO-1 on the expression or activity of oncogenes or cancer-related genes was not elucidated yet. Although, it was shown that HO-1 or its inhibitor, tin mesoporphyrin-IX (SnMPiX) did not affect telomerase and telomerase reverse transcriptase (TERT) [278].

Nevertheless, there are some studies showing that treatment of animals with carcinogens can results in HO-1 induction. Enhanced levels of HO-1 were demonstrated in rats treated with alachlor, the inducer of olfactory mucosal tumors [279]. Correspondingly, the increase of HO-1 was observed in livers exposed to dietary p-dimethylaminoazobenzene (DAB) [65], whereas the downregulation of HO-1 expression was related with malignancy progression [280]. Unfortunately, it is difficult to distinguish whether alteration of HO-1 expression is a reason or outcome of transformation.

Additionally, different levels of HO-1 expression have been detected in the rat strains varying in the vulnerability to carcinogens. For example, carcinogen-resistant DRH rats displaying a significantly lower incidence of liver tumors than the carcinogen-sensitive Donryu strain in response to DAB, have shown much weaker inductions of HO-1 and HGF. Interestingly, these alterations were regarded as an indicator of tissue injury, whereas different levels of expres-

sion of cytochrome P-450 2E, glutathione S-transferase and γ -glutamyltranspeptidase were suggested to be the reason underlying the dissimilar sensitivity of DRH and Donryu rats to carcinogens [281]. Accordingly, the high levels of expression of HO-1 were observed in the Long-Evans with a cinnamon-like color (LEC) rats that spontaneously develops acute hepatitis and hepatoma. Surprisingly, HO-1 overexpression was observed not in cancer lesions but in surrounding healthy tissues, suggesting that HO-1 induction could be rather an adaptive response to oxidative stress [282].

Finally, HO-1 and proliferating cell nuclear antigen (PCNA) expression was correlated with the degree of epithelial dysplasia in human specimen. Interestingly, oral squamous cell carcinoma also showed enhanced expression of HO-1, but this level was not higher than in severe oral epithelial dysplasia (OED) or carcinoma *in situ*. Authors proposed that the augmentation of HO-1 in premalignant oral lesions is part of an early cytoprotection mechanism against carcinogenesis in the oral mucosa [72].

In conclusion, the participation of HO-1 in carcinogenesis needs further investigations, because data available can support both protective and damaging effects. Finally, the discrimination between the primary effect of HO-1 activity on induction of cancer and secondary influence of carcinogenesis on expression of HO-1 is crucial for understanding the problem.

ROLE OF HO-1 IN ANTI-CANCER THERAPY

Studies on the role of HO-1 are important not only for better understanding of tumor-development process but also for clinical practice. Although some reports describe a selective reducing of HO-1 expression in malignant cells, such as adenocarcinoma or tongue squamous carcinoma [235], the majority of analyses indicate that the levels of HO-1 are strongly increased in various tumors [64, 68, 69, 71, 73, 74, 283], especially in those subjected to chemotherapy or exposed to radiation or photodynamic therapy [76, 87, 164] (Table 2).

This upregulation may have important consequences. In the chronic myeloid leukemia-derived cell line K562,

induction of HO-1 has been shown to counteract imatinib-induced apoptosis [78]. In accordance, high HO-1 expression in pancreatic cancer cell lines was associated with increased chemoresistance to gemcitabine [142]. Similarly, upregulated expression of HO-1 in adenocarcinoma treated with photodynamic therapy led to much faster regrowth of tumors in mice [87]. These results support the idea that HO-1 may be a good candidate in antitumor therapies. Thus, pharmacological inhibition of HO-1 has been suggested as a new therapeutic option and potential sensitizer to chemotherapy, radiotherapy, or photodynamic therapy for chronic myeloid leukemia [78], colon carcinoma [164], adenocarcinoma [87], pancreatic cancer [76], melanoma [108, 167], neoplastic mast cells [109], lung cancer [109], hepatoma [124], and urothelial cancer [88]. The usefulness of such treatments has been confirmed in animal models, as administration of ZnPPiX considerably reduced the growth of hepatoma in rats [69], and sarcoma [284] lung cancer [114] or B-cell lymphoma in mice [124].

However, ZnPPiX is weakly soluble in water, which limits its practical use as a drug. Therefore, PEG conjugation was performed to modify ZnPPiX, resulting in a highly water-soluble complex PEG-ZnPPiX [285]. *In vivo* pharmacokinetic analysis revealed that PEG-ZnPPiX administered intravenously had a much longer plasma residence time, 40 times of nonconjugated free ZnPPiX. In addition, PEG-ZnPPiX preferentially accumulated in solid tumors taking the advantage of EPR (*Enhanced Permeability and Retention*) phenomenon [284]. As a result of EPR effect, macromolecules and lipids (bigger than 40 kDa) preferentially and spontaneously leak out of tumor vessels, and remain there at high concentrations for a long time [286]. Recently, another highly water-soluble micellar form of ZnPPiX was obtained, by the use of amphiphilic styrene-maleic acid copolymer (SMA), named SMA-ZnPPiX. It showed similar HO-1 inhibitory activity as native ZnPPiX and exhibited a potent antitumor potential without any apparent side effects [287].

PEG-ZnPPiX and SMA-ZnPPiX inhibited proliferation and induced apoptosis of canine [288] and human mast neoplastic cells *in vitro* [109], while PEG-ZnPPiX-treated

Table 2. Effects of HO-1 Activation in Tumors

Effect of HO-1 Activity	Type of Cancer
↑Proliferation	Prostate cancer [66], sarcoma [69], hepatoma [69], hepatocellular cancer [70], endothelioma [75], pancreatic cancer [76, 113], angioma [77], urothelial cancer cells [88], melanoma [108], neoplastic mast cells [109], lung cancer [114]
↓Proliferation	Prostate cancer [91], lung adenocarcinoma [110], breast cancer [111]
↑Apoptosis,	Breast cancer [198]
↓Apoptosis	Hormone-refractory prostate cancer [66], hepatoma [69, 70, 168], sarcoma [69, 168], pancreatic cancer [76], chronic myeloid leukemia [78, 165], urothelial carcinoma [88], melanoma [108, 167], thyroid carcinoma [122], gastric carcinoma [123], colon carcinoma [158, 164, 168], lung carcinoma [159, 160], acute myeloid leukemia [162]
↑Angiogenesis	Hepatocellular carcinoma [70], glioma [74], melanoma [74, 108], endothelioma [77], pancreatic cancer [113], lung carcinoma [114], fibrosarcoma [208], breast carcinoma [209]
↑Metastasis	Prostate cancer [66], melanoma [108], pancreatic cancer [113], oral squamous carcinoma [228]
↓Metastasis	Oral carcinoma [71, 236, 237], breast cancer [229], colorectal cancer [234, 235]

colon cancer cells exhibited higher rate of apoptosis [284] and became more vulnerable to hydrogen peroxide, t-butyl hydroperoxide, camptothecin and doxorubicin [164]. These promising results have been confirmed *in vivo*. Namely, treatment with PEG-ZnPPIX significantly reduced the growth of murine sarcoma tumors [284]. Moreover, PEG-ZnPPIX and SMA-ZnPPIX administered to mice allowed to overcome resistance of chronic myelocytic leukemia to imatinib [165]. Finally, it is worth to mention that ZnPPIX or its polymer derivatives PEG-ZnPPIX and SMA-ZnPPIX can also be applied in photodynamic therapies, as porphyrins are known photosensitizers [289]. One important advantage of PEG-ZnPPIX/SMA-ZnPPIX over conventional photosensitizer, is the fact that ZnPPIX sufficiently generates highly reactive singlet oxygen under illumination of visible light and xenon light, not only of laser [287].

It should be kept in mind that pharmacological HO-1 inhibitors, such as SnPPIX, ZnPPIX, zinc mesoporphyrin-IX (ZnMPPIX), or zinc deuteroporphyrin-IX (ZnDPPIX), as well as HO-1 activators, such as heme, CoPPIX, or CoCl_2 , are known to exert strong HO-1 independent activities [290, 291]. For example, both ZnPPIX (HO-1 inhibitor) and copper protoporphyrin-IX (CuPPIX, a compound that does not affect HO-1 activity) have been shown to equally diminish tumor blood flow in rats [292, 293], while even a relatively high dose of ZnPPIX (45 $\mu\text{mol/kg}$) injected intraperitoneally was unable to block the activity of HO-1 in tumors [293]. ZnPPIX induced also an accumulation of ROS in sarcoma [284], colon carcinoma [184], and lung carcinoma [160], causing cell apoptosis, and inhibited expression of cyclin-D1 in colon carcinoma, thereby decreasing the rate of proliferation [184]. Again, these effects seemed to be HO-1 independent. Finally in hepatoma and chronic myelogenous leukemia cells, the inhibition of proliferation by ZnPPIX could not be mimicked by SnPPIX, indicating the HO-1 independence [124].

Much more specific strategy is the employment of siRNA. Experiments performed *in vitro* and in animal models clearly confirm that HO-1 could be a promising target of anticancer therapy. For example, targeted knockdown of HO-1 achieved by transfection of siRNA induced apoptosis of cultured colon carcinoma [158, 284] or leukemic cells [162, 165], and resulted in diminished proliferation, reduced growth and decreased angiogenesis in orthotopic hepatocellular tumors [70]. The same approach in prostate cancer cells led to reduced cell proliferation, decreased cell survival, and attenuated cell invasion *in vitro*, what was reflected by inhibition of prostate tumor growth and lymph node and lung metastases *in vivo* [66]. Similarly, HO-1 siRNA increased apoptosis, augmented the cisplatin cytotoxicity toward lung cancer cells [85, 160], and sensitized pancreatic carcinoma to oxidative stress, gemcitabine, or γ -radiation [76].

The RNAi approach has been widely used for drug development and several clinical trials are ongoing [294]. However, there are still some concerns and challenges to be overcome, such as efficacy and specificity of oligonucleotide delivery, potential off-target effects, and activation of innate immune responses. Various siRNA delivery strategies have been established, including employing of nanoparticles, cationic lipids, antibodies, cholesterol, aptamers and viral

vectors for short hairpin RNAs (shRNAs). In some cases the oral or aerosol-mediated delivery might be more favorable. For example in mice, the aerosol delivery of anti-Akt1 siRNA was successfully carried out to treat the lung cancer [295].

Ectopically applied siRNAs can modify the expression of non-targeted transcripts [296, 297]. Thus, it has been shown that inhibition of neovascularization by siRNA in murine model of age-related macular degeneration was not due to a specific knockdown of VEGF and its receptor, but due to a dsRNA-dependent activation of TLR3, triggering the $\text{IFN}\gamma$ and IL-12 production [298]. Moreover, certain motifs in siRNAs can trigger the type I IFN production via activation of TLR7 and TLR8 [299, 300]. Fortunately, this off-targeting can be controlled by a 2'-OMe modification of siRNA [301]. It should be also kept in mind that the levels of siRNA have to be controlled, as too high expression in studies with AAV vectors, where Pol III was used to express shRNA, led to in lethality of mice due to acute liver failure [302]. Thus, the siRNA strategy could be a promising tool for specific and targeted HO-1 inhibition in cancers.

CONCLUSIONS

Protumoral activities of HO-1 seem to be related to its cytoprotective and antiapoptotic properties, which result in better tumor cell survival and higher resistance to different types of therapies. HO-1 affects also tumor cell proliferation and differentiation, which may lead to development of more aggressive phenotype. Moreover, HO-1 can facilitate angiogenesis, a key process in growth of primary and secondary tumors. Finally, HO-1 modulates behavior of inflammatory cells, favoring state of immunosuppression that assists tumor progression. Thus, HO-1 seems to accelerate tumor growth and metastasis. Therefore, we propose that inhibition of HO-1 activity might be considered as an independent therapeutic strategy or as a sensitizing approach in combination with chemotherapy, radiation, or photodynamic therapy to potentiate their anticancer effects (Fig. 8). On the other hand, HO-1 can protect healthy tissues against carcinogenesis, and can be beneficial if expressed in stroma cells. Consequently, it seems that such inhibition should be targeted specifically to the cancer cells.




Fig. (8). Schematic summary of HO-1 effects on tumor biology.

ACKNOWLEDGEMENTS

We thank our coworkers who contributed to the experiments described in this review. Supported by grants No. N401 297835, N301 314837, 311/N-COST/2008 and 347/N-INCA/2008 from the Ministry of Science and Higher Education. AJ is a recipient of the Wellcome Trust Senior Research Fellowship in Basic Biomedical Science. Department of Medical Biotechnology participates in the COST CM0602 Action (ANGIOKEM), and TD0901 (HypoxiaNet) both supported by European Commission. The Faculty of Biochemistry, Biophysics and Biotechnology of the Jagiellonian University is a beneficiary of the structural funds from the European Union (grant No: POIG. 01.01.02-00-109/09, 02.01.00-12-064/08 and 02.02.00-00-014/08).

REFERENCES

- [1] Femiano F, Scully C. DNA cytometry of oral leukoplakia and oral lichen planus. *Med Oral Patol Oral Cir Bucal* 2005; 10 (Suppl 1): E9-14.
- [2] Sen S. Aneuploidy and cancer. *Curr Opin Oncol* 2000; 12: 82-8.
- [3] Zhang L, Rosin MP. Loss of heterozygosity: a potential tool in management of oral premalignant lesions? *J Oral Pathol Med* 2001; 30: 513-20.
- [4] Fujita H, Nagata M, Hoshina H, *et al.* Clinical significance and usefulness of quantification of telomerase activity in oral malignant and nonmalignant lesions. *Int J Oral Maxillofac Surg* 2004; 33: 693-9.
- [5] Todd R, Wong DT. Oncogenes. *Anticancer Res* 1999; 19: 4729-46.
- [6] Levine AJ. p53, the cellular gatekeeper for growth and division. *Cell* 1997; 88: 323-31.
- [7] Haupt Y, Barak Y, Oren M. Cell type-specific inhibition of p53-mediated apoptosis by mdm2. *EMBO J* 1996; 15: 1596-606.
- [8] Potgens AJ, Westphal HR, de Waal RM, Ruiter DJ. The role of vascular permeability factor and basic fibroblast growth factor in tumor angiogenesis. *Biol Chem Hoppe Seyler* 1995; 376: 57-70.
- [9] Asahara T, Masuda H, Takahashi T, *et al.* Bone marrow origin of endothelial progenitor cells responsible for postnatal vasculogenesis in physiological and pathological neovascularization. *Circ Res* 1999; 85: 221-8.
- [10] Mantovani A, Allavena P, Sica A, Balkwill F. Cancer-related inflammation. *Nature* 2008; 454: 436-44.
- [11] Karin M. Nuclear factor- κ B in cancer development and progression. *Nature* 2006; 441: 431-6.
- [12] de Martel C, Franceschi S. Infections and cancer: established associations and new hypotheses. *Crit Rev Oncol Hematol* 2009; 70: 183-94.
- [13] Takahashi H, Ogata H, Nishigaki R, Broide DH, Karin M. Tobacco smoke promotes lung tumorigenesis by triggering IKK β - and JNK1-dependent inflammation. *Cancer Cell* 2010; 17: 89-97.
- [14] Park EJ, Lee JH, Yu GY, *et al.* Dietary and genetic obesity promote liver inflammation and tumorigenesis by enhancing IL-6 and TNF expression. *Cell* 2010; 140: 197-208.
- [15] Khasawneh J, Schulz MD, Walch A, *et al.* Inflammation and mitochondrial fatty acid β -oxidation link obesity to early tumor promotion. *Proc Natl Acad Sci USA* 2009; 106: 3354-9.
- [16] Ershler WB, Keller ET. Age-associated increased interleukin-6 gene expression, late-life diseases, and frailty. *Annu Rev Med* 2000; 51: 245-70.
- [17] Rodier F, Coppe JP, Patil CK, *et al.* Persistent DNA damage signalling triggers senescence-associated inflammatory cytokine secretion. *Nat Cell Biol* 2009; 11: 973-9.
- [18] Davies KJ. Oxidative stress: the paradox of aerobic life. *Biochem Soc Symp* 1995; 61: 1-31.
- [19] Aruoma OI, Grootveld M, Baharun T. Free radicals in biology and medicine: from inflammation to biotechnology. *Biofactors* 2006; 27: 1-3.
- [20] Demple B, Harrison L. Repair of oxidative damage to DNA: enzymology and biology. *Annu Rev Biochem* 1994; 63: 915-48.
- [21] Tenhunen R, Marver HS, Schmid R. The enzymatic conversion of heme to bilirubin by microsomal heme oxygenase. *Proc Natl Acad Sci USA* 1968; 61: 748-55.
- [22] Schacter BA. Heme catabolism by heme oxygenase: physiology, regulation, and mechanism of action. *Semin Hematol* 1988; 25: 349-69.
- [23] Graham BL, Mink JT, Cotton DJ. Dynamic measurements of CO diffusing capacity using discrete samples of alveolar gas. *J Appl Physiol* 1983; 54: 73-9.
- [24] Snyder SH, Baranano DE. Heme oxygenase: a font of multiple messengers. *Neuropsychopharmacology* 2001; 25: 294-8.
- [25] De Matteis F, Dawson SJ, Pons N, Pipino S. Bilirubin and uroporphyrinogen oxidation by induced cytochrome P4501A and cytochrome P4502B. Role of polyhalogenated biphenyls of different configuration. *Biochem Pharmacol* 2002; 63: 615-24.
- [26] Maines MD, Ibrahim NG, Kappas A. Solubilization and partial purification of heme oxygenase from rat liver. *J Biol Chem* 1977; 252: 5900-3.
- [27] Yoshida T, Kikuchi G. Purification and properties of heme oxygenase from pig spleen microsomes. *J Biol Chem* 1978; 253: 4224-9.
- [28] Yoshinaga T, Sassa S, Kappas A. Purification and properties of bovine spleen heme oxygenase. Amino acid composition and sites of action of inhibitors of heme oxidation. *J Biol Chem* 1982; 257: 7778-85.
- [29] Abraham NG, Lin JH, Dunn MW, Schwartzman ML. Presence of heme oxygenase and NADPH cytochrome P-450 (c) reductase in human corneal epithelium. *Invest Ophthalmol Vis Sci* 1987; 28: 1464-72.
- [30] Muller RM, Taguchi H, Shibahara S. Nucleotide sequence and organization of the rat heme oxygenase gene. *J Biol Chem* 1987; 262: 6795-802.
- [31] McCoubrey WK, Jr., Ewing JF, Maines MD. Human heme oxygenase-2: characterization and expression of a full-length cDNA and evidence suggesting that the two HO-2 transcripts may differ by choice of polyadenylation signal. *Arch Biochem Biophys* 1992; 295: 13-20.
- [32] Maines MD, Trakshel GM, Kutty RK. Characterization of two constitutive forms of rat liver microsomal heme oxygenase. Only one molecular species of the enzyme is inducible. *J Biol Chem* 1986; 261: 411-9.
- [33] Cruse I, Maines MD. Evidence suggesting that the two forms of heme oxygenase are products of different genes. *J Biol Chem* 1988; 263: 3348-53.
- [34] McCoubrey WK, Jr., Maines MD. The structure, organization and differential expression of the gene encoding rat heme oxygenase-2. *Gene* 1994; 139: 155-61.
- [35] Alam J, Shibahara S, Smith A. Transcriptional activation of the heme oxygenase gene by heme and cadmium in mouse hepatoma cells. *J Biol Chem* 1989; 264: 6371-5.
- [36] Lavrovsky Y, Schwartzman ML, Levere RD, Kappas A, Abraham NG. Identification of binding sites for transcription factors NF- κ B and AP-2 in the promoter region of the human heme oxygenase 1 gene. *Proc Natl Acad Sci USA* 1994; 91: 5987-91.
- [37] Foresti R, Clark JE, Green CJ, Motterlini R. Thiol compounds interact with nitric oxide in regulating heme oxygenase-1 induction in endothelial cells. Involvement of superoxide and peroxynitrite anions. *J Biol Chem* 1997; 272: 18411-7.
- [38] Terry CM, Clikeman JA, Hoidal JR, Callahan KS. Effect of tumor necrosis factor- α and interleukin-1 α on heme oxygenase-1 expression in human endothelial cells. *Am J Physiol* 1998; 274: H883-91.
- [39] Stuhlmeier KM. Activation and regulation of Hsp32 and Hsp70. *Eur J Biochem* 2000; 267: 1161-7.
- [40] Zhang WL, Tsuneishi S, Nakamura H. Induction of heat shock proteins and its effects on glial differentiation in rat C6 glioblastoma cells. *Kobe J Med Sci* 2001; 47: 77-95.
- [41] Jozkowicz A, Huk I, Nigisch A, *et al.* Effect of prostaglandin-J₂ on VEGF synthesis depends on the induction of heme oxygenase-1. *Antioxid Redox Signal* 2002; 4: 577-85.
- [42] Motterlini R, Foresti R, Bassi R, *et al.* Endothelial heme oxygenase-1 induction by hypoxia. Modulation by inducible nitric-oxide synthase and S-nitrosothiols. *J Biol Chem* 2000; 275: 13613-20.
- [43] Shibahara S, Nakayama M, Kitamuro T, Udono-Fujimori R, Takahashi K. Repression of heme oxygenase-1 expression as a defense strategy in humans. *Exp Biol Med (Maywood)* 2003; 228: 472-3.

- [44] Gonzalez-Michaca L, Farrugia G, Croatt AJ, Alam J, Nath KA. Heme: a determinant of life and death in renal tubular epithelial cells. *Am J Physiol Renal Physiol* 2004; 286: F370-7.
- [45] Bauer I, Rensing H, Florax A, *et al.* Expression pattern and regulation of heme oxygenase-1/heat shock protein 32 in human liver cells. *Shock* 2003; 20: 116-22.
- [46] Nishie A, Ono M, Shono T, *et al.* Macrophage infiltration and heme oxygenase-1 expression correlate with angiogenesis in human gliomas. *Clin Cancer Res* 1999; 5: 1107-13.
- [47] Pae HO, Oh GS, Choi BM, Chae SC, Chung HT. Differential expressions of heme oxygenase-1 gene in CD25- and CD25+ subsets of human CD4+ T cells. *Biochem Biophys Res Commun* 2003; 306: 701-5.
- [48] Balla J, Vercellotti GM, Jeney V, *et al.* Heme, heme oxygenase, and ferritin: how the vascular endothelium survives (and dies) in an iron-rich environment. *Antioxid Redox Signal* 2007; 9: 2119-37.
- [49] Balla J, Jacob HS, Balla G, *et al.* Endothelial-cell heme uptake from heme proteins: induction of sensitization and desensitization to oxidant damage. *Proc Natl Acad Sci USA* 1993; 90: 9285-9.
- [50] Dallegri F, Ballestrero A, Frumento G, Patrone F. Augmentation of neutrophil-mediated erythrocyte lysis by cells derived *in vitro* from human monocytes. *Blood* 1987; 70: 1743-9.
- [51] Weiss SJ. Neutrophil-mediated methemoglobin formation in the erythrocyte. The role of superoxide and hydrogen peroxide. *J Biol Chem* 1982; 257: 2947-53.
- [52] Foresti R, Bains S, Sulc F, *et al.* The interaction of nitric oxide with distinct hemoglobins differentially amplifies endothelial heme uptake and heme oxygenase-1 expression. *J Pharmacol Exp Ther* 2006; 317: 1125-1133.
- [53] Graca-Souza AV, Arruda MA, de Freitas MS, Barja-Fidalgo C, Oliveira PL. Neutrophil activation by heme: implications for inflammatory processes. *Blood* 2002; 99: 4160-5.
- [54] Wagener FA, Eggert A, Boerman OC, *et al.* Heme is a potent inducer of inflammation in mice and is counteracted by heme oxygenase. *Blood* 2001; 98: 1802-11.
- [55] Wagener FA, Feldman E, de Witte T, Abraham NG. Heme induces the expression of adhesion molecules ICAM-1, VCAM-1, and E selectin in vascular endothelial cells. *Proc Soc Exp Biol Med* 1997; 216: 456-63.
- [56] Balla G, Vercellotti GM, Eaton JW, Jacob HS. Iron loading of endothelial cells augments oxidant damage. *J Lab Clin Med* 1990; 116: 546-54.
- [57] Eskew JD, Vanacore RM, Sung L, Morales PJ, Smith A. Cellular protection mechanisms against extracellular heme. heme-hemopexin, but not free heme, activates the N-terminal c-jun kinase. *J Biol Chem* 1999; 274: 638-48.
- [58] Bunn HF, Jandl JH. Exchange of heme among hemoglobins and between hemoglobin and albumin. *J Biol Chem* 1968; 243: 465-75.
- [59] Wang R, Wu L. Interaction of selective amino acid residues of K(ca) channels with carbon monoxide. *Exp Biol Med (Maywood)* 2003; 228: 474-80.
- [60] L'Abbate A, Neglia D, Vecoli C, *et al.* Beneficial effect of heme oxygenase-1 expression on myocardial ischemia-reperfusion involves an increase in adiponectin in mildly diabetic rats. *Am J Physiol Heart Circ Physiol* 2007; 293: H3532-41.
- [61] Otterbein LE, Bach FH, Alam J, *et al.* Carbon monoxide has anti-inflammatory effects involving the mitogen-activated protein kinase pathway. *Nat Med* 2000; 6: 422-8.
- [62] Paller MS, Jacob HS. Cytochrome P-450 mediates tissue-damaging hydroxyl radical formation during reoxygenation of the kidney. *Proc Natl Acad Sci USA* 1994; 91: 7002-6.
- [63] Schacter BA, Kurz P. Alterations in hepatic and splenic microsomal electron transport system components, drug metabolism, heme oxygenase activity, and cytochrome P-450 turnover in Murphy-Sturm lymphosarcoma-bearing rats. *Cancer Res* 1982; 42: 3557-64.
- [64] Maines MD, Abrahamsson PA. Expression of heme oxygenase-1 (HSP32) in human prostate: normal, hyperplastic, and tumor tissue distribution. *Urology* 1996; 47: 727-33.
- [65] Sacca P, Meiss R, Casas G, *et al.* Nuclear translocation of haeme oxygenase-1 is associated to prostate cancer. *Br J Cancer* 2007; 97: 1683-9.
- [66] Alaoui-Jamali MA, Bismar TA, Gupta A, *et al.* A novel experimental heme oxygenase-1-targeted therapy for hormone-refractory prostate cancer. *Cancer Res* 2009; 69: 8017-24.
- [67] Hara E, Takahashi K, Tominaga T, *et al.* Expression of heme oxygenase and inducible nitric oxide synthase mRNA in human brain tumors. *Biochem Biophys Res Commun* 1996; 224: 153-8.
- [68] Goodman AI, Choudhury M, da Silva JL, Schwartzman ML, Abraham NG. Overexpression of the heme oxygenase gene in renal cell carcinoma. *Proc Soc Exp Biol Med* 1997; 214: 54-61.
- [69] Doi K, Akaike T, Fujii S, *et al.* Induction of haem oxygenase-1 nitric oxide and ischaemia in experimental solid tumours and implications for tumour growth. *Br J Cancer* 1999; 80: 1945-54.
- [70] Sass G, Leukel P, Schmitz V, *et al.* Inhibition of heme oxygenase 1 expression by small interfering RNA decreases orthotopic tumor growth in livers of mice. *Int J Cancer* 2008; 123: 1269-77.
- [71] Tsuji MH, Yanagawa T, Iwasa S, *et al.* Heme oxygenase-1 expression in oral squamous cell carcinoma as involved in lymph node metastasis. *Cancer Lett* 1999; 138: 53-9.
- [72] Lee J, Lee SK, Lee BU, *et al.* Upregulation of heme oxygenase-1 in oral epithelial dysplasias. *Int J Oral Maxillofac Surg* 2008; 37: 287-92.
- [73] Deininger MH, Meyermann R, Trautmann K, *et al.* Heme oxygenase (HO)-1 expressing macrophages/microglial cells accumulate during oligodendroglioma progression. *Brain Res* 2000; 882: 1-8.
- [74] Torisu-Itakura H, Furue M, Kuwano M, Ono M. Co-expression of thymidine phosphorylase and heme oxygenase-1 in macrophages in human malignant vertical growth melanomas. *Jpn J Cancer Res* 2000; 91: 906-10.
- [75] McAllister SC, Hansen SG, Ruhl RA, *et al.* Kaposi sarcoma-associated herpesvirus (KSHV) induces heme oxygenase-1 expression and activity in KSHV-infected endothelial cells. *Blood* 2004; 103: 3465-73.
- [76] Berberat PO, Dambrauskas Z, Gulbinas A, *et al.* Inhibition of heme oxygenase-1 increases responsiveness of pancreatic cancer cells to anticancer treatment. *Clin Cancer Res* 2005; 11: 3790-8.
- [77] Marinissen MJ, Tanos T, Bolos M, *et al.* Inhibition of heme oxygenase-1 interferes with the transforming activity of the Kaposi sarcoma herpesvirus-encoded G protein-coupled receptor. *J Biol Chem* 2006; 281: 11332-46.
- [78] Mayerhofer M, Florian S, Krauth MT, *et al.* Identification of heme oxygenase-1 as a novel BCR/ABL-dependent survival factor in chronic myeloid leukemia. *Cancer Res* 2004; 64: 3148-54.
- [79] Liu J, Benbrahim-Tallaa L, Qian X, *et al.* Further studies on aberrant gene expression associated with arsenic-induced malignant transformation in rat liver TRL1215 cells. *Toxicol Appl Pharmacol* 2006; 216: 407-15.
- [80] Calvisi DF, Ladu S, Hironaka K, Factor VM, Thorgeirsson SS. Vitamin E down-modulates iNOS and NADPH oxidase in c-Myc/TGF α transgenic mouse model of liver cancer. *J Hepatol* 2004; 41: 815-22.
- [81] Wen WN. Methemoglobin contributes to the growth of human tumor cells. *Life Sci* 2002; 70: 907-16.
- [82] Morita K, Lee MS, Her S. Possible relation of heme-induced HO-1 expression to the upregulation of VEGF and BDNF mRNA levels in rat C6 glioma cells. *J Mol Neurosci* 2009; 38: 31-40.
- [83] Inoue K, Takahashi T, Uehara K, *et al.* Protective role of heme oxygenase 1 in the intestinal tissue injury in hemorrhagic shock in rats. *Shock* 2008; 29: 252-61.
- [84] Hsu JT, Kan WH, Hsieh CH, *et al.* Mechanism of estrogen-mediated attenuation of hepatic injury following trauma-hemorrhage: Akt-dependent HO-1 up-regulation. *J Leukoc Biol* 2007; 82: 1019-26.
- [85] Kuroda H, Takeno M, Murakami S, *et al.* Inhibition of heme oxygenase-1 with an epidermal growth factor receptor inhibitor and cisplatin decreases proliferation of lung cancer A549 cells. *Lung Cancer* 2010; 67: 31-6.
- [86] Kocanova S, Buytaert E, Matroule JY, *et al.* Induction of heme-oxygenase 1 requires the p38MAPK and PI3K pathways and suppresses apoptotic cell death following hypericin-mediated photodynamic therapy. *Apoptosis* 2007; 12: 731-41.
- [87] Nowis D, Legat M, Grzela T, *et al.* Heme oxygenase-1 protects tumor cells against photodynamic therapy-mediated cytotoxicity. *Oncogene* 2006; 25: 3365-74.
- [88] Miyake M, Ishii M, Kawashima K, *et al.* siRNA-mediated knockdown of the heme synthesis and degradation pathways: modulation of treatment effect of 5-aminolevulinic acid-based photodynamic therapy in urothelial cancer cell lines. *Photochem Photobiol* 2009; 85: 1020-7.

- [89] Rapozzi V, Miculan M, Xodo LE. Evidence that photoactivated pheophorbide a causes in human cancer cells a photodynamic effect involving lipid peroxidation. *Cancer Biol Ther* 2009; 8: 1318-27.
- [90] Boschetto P, Zeni E, Mazzetti L, *et al.* Decreased heme-oxygenase (HO)-1 in the macrophages of non-small cell lung cancer. *Lung Cancer* 2008; 59: 192-7.
- [91] Gueron G, De Siervi A, Ferrando M, *et al.* Critical role of endogenous heme oxygenase 1 as a tuner of the invasive potential of prostate cancer cells. *Mol Cancer Res* 2009; 7: 1745-55.
- [92] Lin Q, Weis S, Yang G, *et al.* Heme oxygenase-1 protein localizes to the nucleus and activates transcription factors important in oxidative stress. *J Biol Chem* 2007; 282: 20621-33.
- [93] Li Volti G, Ientile R, Abraham NG, *et al.* Immunocytochemical localization and expression of heme oxygenase-1 in primary astroglial cell cultures during differentiation: effect of glutamate. *Biochem Biophys Res Commun* 2004; 315: 517-24.
- [94] Suttner DM, Sridhar K, Lee CS, *et al.* Protective effects of transient HO-1 overexpression on susceptibility to oxygen toxicity in lung cells. *Am J Physiol* 1999; 276: L443-51.
- [95] Giordano A, Nisoli E, Tonello C, *et al.* Expression and distribution of heme oxygenase-1 and -2 in rat brown adipose tissue: the modulatory role of the noradrenergic system. *FEBS Lett* 2000; 487: 171-5.
- [96] Hanselmann C, Mauch C, Werner S. Haem oxygenase-1: a novel player in cutaneous wound repair and psoriasis? *Biochem J* 2001; 353: 459-66.
- [97] Clark JE, Green CJ, Motterlini R. Involvement of the heme oxygenase-carbon monoxide pathway in keratinocyte proliferation. *Biochem Biophys Res Commun* 1997; 241: 215-20.
- [98] Jozkowicz A, Huk I, Nigisch A, *et al.* Heme oxygenase and angiogenic activity of endothelial cells: stimulation by carbon monoxide and inhibition by tin protoporphyrin-IX. *Antioxid Redox Signal* 2003; 5: 155-62.
- [99] Lee SS, Gao W, Mazzola S, *et al.* Heme oxygenase-1, carbon monoxide, and bilirubin induce tolerance in recipients toward islet allografts by modulating T regulatory cells. *FASEB J* 2007; 21: 3450-7.
- [100] Brusko TM, Wasserfall CH, Agarwal A, Kapturczak MH, Atkinson MA. An integral role for heme oxygenase-1 and carbon monoxide in maintaining peripheral tolerance by CD4+CD25+ regulatory T cells. *J Immunol* 2005; 174: 5181-6.
- [101] Petrache I, Otterbein LE, Alam J, Wiegand GW, Choi AM. Heme oxygenase-1 inhibits TNF α -induced apoptosis in cultured fibroblasts. *Am J Physiol Lung Cell Mol Physiol* 2000; 278: L312-9.
- [102] Duckers HJ, Boehm M, True AL, *et al.* Heme oxygenase-1 protects against vascular constriction and proliferation. *Nat Med* 2001; 7: 693-8.
- [103] Ollinger R, Bilban M, Erat A, *et al.* Bilirubin: a natural inhibitor of vascular smooth muscle cell proliferation. *Circulation* 2005; 112: 1030-9.
- [104] Kim JY, Cho HJ, Sir JJ, *et al.* Sulfasalazine induces haem oxygenase-1 via ROS-dependent Nrf2 signalling, leading to control of neointimal hyperplasia. *Cardiovasc Res* 2009; 82: 550-60.
- [105] Aizawa T, Ishizaka N, Kurokawa K, *et al.* Different effects of angiotensin II and catecholamine on renal cell apoptosis and proliferation in rats. *Kidney Int* 2001; 59: 645-53.
- [106] Song R, Mahidhara RS, Zhou Z, *et al.* Carbon monoxide inhibits T lymphocyte proliferation via caspase-dependent pathway. *J Immunol* 2004; 172: 1220-6.
- [107] Song W, Su H, Song S, Paudel HK, Schipper HM. Over-expression of heme oxygenase-1 promotes oxidative mitochondrial damage in rat astroglia. *J Cell Physiol* 2006; 206: 655-63.
- [108] Was H, Cichon T, Smolarczyk R, *et al.* Overexpression of heme oxygenase-1 in murine melanoma: increased proliferation and viability of tumor cells, decreased survival of mice. *Am J Pathol* 2006; 169: 2181-98.
- [109] Kondo R, Gleixner KV, Mayerhofer M, *et al.* Identification of heat shock protein 32 (Hsp32) as a novel survival factor and therapeutic target in neoplastic mast cells. *Blood* 2007; 110: 661-9.
- [110] Lee PJ, Alam J, Wiegand GW, Choi AM. Overexpression of heme oxygenase-1 in human pulmonary epithelial cells results in cell growth arrest and increased resistance to hyperoxia. *Proc Natl Acad Sci USA* 1996; 93: 10393-8.
- [111] Hill M, Pereira V, Chauveau C, *et al.* Heme oxygenase-1 inhibits rat and human breast cancer cell proliferation: mutual cross inhibition with indoleamine 2,3-dioxygenase. *FASEB J* 2005; 19: 1957-68.
- [112] La P, Fernando AP, Wang Z, *et al.* Zinc protoporphyrin regulates cyclin-D1 expression independent of heme oxygenase inhibition. *J Biol Chem* 2009; 284: 36302-11.
- [113] Sunamura M, Duda DG, Ghattas MH, *et al.* Heme oxygenase-1 accelerates tumor angiogenesis of human pancreatic cancer. *Angiogenesis* 2003; 6: 15-24.
- [114] Hirai K, Sasahira T, Ohmori H, Fujii K, Kuniyasu H. Inhibition of heme oxygenase-1 by zinc protoporphyrin IX reduces tumor growth of LL/2 lung cancer in C57BL mice. *Int J Cancer* 2007; 120: 500-5.
- [115] Meiller A, Alvarez S, Drane P, *et al.* p53-dependent stimulation of redox-related genes in the lymphoid organs of γ -irradiated-mice identification of haeme-oxygenase 1 as a direct p53 target gene. *Nucleic Acids Res* 2007; 35: 6924-34.
- [116] DeYulia GJ, Jr., Carcamo JM, Borquez-Ojeda O, Shelton CC, Golde DW. Hydrogen peroxide generated extracellularly by receptor-ligand interaction facilitates cell signaling. *Proc Natl Acad Sci USA* 2005; 102: 5044-9.
- [117] Ruiz-Ramos R, Lopez-Carrillo L, Rios-Perez AD, De Vizcaya-Ruiz A, Cebrian ME. Sodium arsenite induces ROS generation, DNA oxidative damage, HO-1 and c-Myc proteins, NF- κ B activation and cell proliferation in human breast cancer MCF-7 cells. *Mutat Res* 2009; 674: 109-15.
- [118] Ball KL. p21: structure and functions associated with cyclin-CDK binding. *Prog Cell Cycle Res* 1997; 3: 125-34.
- [119] Karjalainen JM, Eskelinen MJ, Kellokoski JK, *et al.* p21(WAF1/CIP1) expression in stage I cutaneous malignant melanoma: its relationship with p53, cell proliferation and survival. *Br J Cancer* 1999; 79: 895-902.
- [120] Li MY, Hsin MK, Yip J, *et al.* PPAR γ activation extinguishes smoking carcinogen by inhibiting NNN-mediated proliferation. *Am J Respir Cell Mol Biol* 2010; 42: 113-22.
- [121] Ohata M, Nakamura S, Fujita H, Isemura M. Prognostic implications of p21 (Waf1/Cip1) immunolocalization in multiple myeloma. *Biomed Res* 2005; 26: 91-8.
- [122] Chen GG, Liu ZM, Vlantis AC, *et al.* Heme oxygenase-1 protects against apoptosis induced by tumor necrosis factor- α and cycloheximide in papillary thyroid carcinoma cells. *J Cell Biochem* 2004; 92: 1246-56.
- [123] Liu ZM, Chen GG, Ng EK, *et al.* Upregulation of heme oxygenase-1 and p21 confers resistance to apoptosis in human gastric cancer cells. *Oncogene* 2004; 23: 503-13.
- [124] La P, Fernando AP, Wang Z, *et al.* Zinc protoporphyrin regulates cyclin-D1 expression independent of heme oxygenase inhibition. *J Biol Chem* 2009; 284: 36302-11.
- [125] Li Volti G, Sacerdoti D, Sangras B, *et al.* Carbon monoxide signaling in promoting angiogenesis in human microvessel endothelial cells. *Antioxid Redox Signal* 2005; 7: 704-10.
- [126] Jeon EM, Choi HC, Lee KY, Chang KC, Kang YJ. Hemin inhibits hypertensive rat vascular smooth muscle cell proliferation through regulation of cyclin D and p21. *Arch Pharm Res* 2009; 32: 375-82.
- [127] Schwer CI, Guerrero AM, Humar M, *et al.* Heme oxygenase-1 inhibits the proliferation of pancreatic stellate cells by repression of the extracellular signal-regulated kinase1/2 pathway. *J Pharmacol Exp Ther* 2008; 327: 863-71.
- [128] Woo SW, Lee SH, Ko G, Kim YC, Sohn DH. Isoliquiritigenin inhibits cell proliferation by a heme oxygenase-dependent pathway in rat hepatic stellate cells. *Planta Med* 2008; 74: 834-9.
- [129] Iwasa H, Han J, Ishikawa F. Mitogen-activated protein kinase p38 defines the common senescence-signalling pathway. *Genes Cells* 2003; 8: 131-44.
- [130] Kim HP, Wang X, Nakao A, *et al.* Caveolin-1 expression by means of p38 β mitogen-activated protein kinase mediates the antiproliferative effect of carbon monoxide. *Proc Natl Acad Sci USA* 2005; 102: 11319-24.
- [131] Cohen AW, Hnasko R, Schubert W, Lisanti MP. Role of caveolae and caveolins in health and disease. *Physiol Rev* 2004; 84: 1341-79.
- [132] Volonte D, Zhang K, Lisanti MP, Galbati F. Expression of caveolin-1 induces premature cellular senescence in primary cultures of murine fibroblasts. *Mol Biol Cell* 2002; 13: 2502-17.

- [133] Engelman JA, Wykoff CC, Yasuhara S, *et al.* Recombinant expression of caveolin-1 in oncogenically transformed cells abrogates anchorage-independent growth. *J Biol Chem* 1997; 272: 16374-81.
- [134] Kim HP, Wang X, Galbiati F, Ryter SW, Choi AM. Caveolae compartmentalization of heme oxygenase-1 in endothelial cells. *FASEB J* 2004; 18: 1080-9.
- [135] Taille C, Almolkhi A, Benhamed M, *et al.* Heme oxygenase inhibits human airway smooth muscle proliferation via a bilirubin-dependent modulation of ERK1/2 phosphorylation. *J Biol Chem* 2003; 278: 27160-8.
- [136] Wagner EF, Nebreda AR. Signal integration by JNK and p38 MAPK pathways in cancer development. *Nat Rev Cancer* 2009; 9: 537-49.
- [137] Fujita H, Yamamoto M, Yamagami T, *et al.* Sequential activation of genes for heme pathway enzymes during erythroid differentiation of mouse Friend virus-transformed erythroleukemia cells. *Biochim Biophys Acta* 1991; 1090: 311-6.
- [138] Richardson DR. Molecular mechanisms of iron uptake by cells and the use of iron chelators for the treatment of cancer. *Curr Med Chem* 2005; 12: 2711-29.
- [139] Darnell G, Richardson DR. The potential of iron chelators of the pyridoxal isonicotinoyl hydrazone class as effective antiproliferative agents III: the effect of the ligands on molecular targets involved in proliferation. *Blood* 1999; 94: 781-92.
- [140] Brodie C, Siriwardana G, Lucas J, *et al.* Neuroblastoma sensitivity to growth inhibition by deferrioxamine: evidence for a block in G1 phase of the cell cycle. *Cancer Res* 1993; 53: 3968-75.
- [141] Donfrancesco A, Deb G, Dominici C, *et al.* Effects of a single course of deferrioxamine in neuroblastoma patients. *Cancer Res* 1990; 50: 4929-30.
- [142] Nuhn P, Kunzli BM, Hennig R, *et al.* Heme oxygenase-1 and its metabolites affect pancreatic tumor growth *in vivo*. *Mol Cancer* 2009; 8: 37-43.
- [143] Wegiel B, Chin BY, Otterbein LE. Inhale to survive, cycle or die? Carbon monoxide and cellular proliferation. *Cell Cycle* 2008; 7: 1379-84.
- [144] Zwerina J, Tzima S, Hayer S, *et al.* Heme oxygenase 1 (HO-1) regulates osteoclastogenesis and bone resorption. *FASEB J* 2005; 19: 2011-3.
- [145] Lin TH, Tang CH, Hung SY, *et al.* Upregulation of heme oxygenase-1 inhibits the maturation and mineralization of osteoblasts. *J Cell Physiol* 2010; 222: 757-68.
- [146] Zarjou A, Jeney V, Arosio P, *et al.* Ferritin prevents calcification and osteoblastic differentiation of vascular smooth muscle cells. *J Am Soc Nephrol* 2009; 20: 1254-63.
- [147] Listopad J, Asadullah K, Sievers C, *et al.* Heme oxygenase-1 inhibits T cell-dependent skin inflammation and differentiation and function of antigen-presenting cells. *Exp Dermatol* 2007; 16: 661-70.
- [148] Vanella L, Kim DH, Asprinio D, *et al.* HO-1 expression increases mesenchymal stem cell-derived osteoblasts but decreases adipocyte lineage. *Bone* 2010; 46: 236-43.
- [149] Koiso Y, Fujimoto Y, Matsumura D, Nakajima O, Hashimoto Y. Modification by heme oxygenase inhibitor, tin protoporphyrin, of cellular differentiation of human myeloid leukemia K562 cell line. *Biol Pharm Bull* 1999; 22: 439-40.
- [150] Owusu-Ansah E, Banerjee U. Reactive oxygen species prime *Drosophila* haematopoietic progenitors for differentiation. *Nature* 2009; 461: 537-41.
- [151] Wang JY, Lee YT, Chang PF, Chau LY. Hemin promotes proliferation and differentiation of endothelial progenitor cells via activation of AKT and ERK. *J Cell Physiol* 2009; 219: 617-25.
- [152] Savickiene J, Borutinskaite VV, Treigyte G, Magnusson KE, Navakasienė R. The novel histone deacetylase inhibitor BML-210 exerts growth inhibitory, proapoptotic and differentiation stimulating effects on the human leukemia cell lines. *Eur J Pharmacol* 2006; 549: 9-18.
- [153] Stocker R, Yamamoto Y, McDonagh AF, Glazer AN, Ames BN. Bilirubin is an antioxidant of possible physiological importance. *Science* 1987; 235: 1043-6.
- [154] Hunt RC, Handy I, Smith A. Heme-mediated reactive oxygen species toxicity to retinal pigment epithelial cells is reduced by hemopexin. *J Cell Physiol* 1996; 168: 81-6.
- [155] Eisenstein RS, Garcia-Mayol D, Pettingell W, Munro HN. Regulation of ferritin and heme oxygenase synthesis in rat fibroblasts by different forms of iron. *Proc Natl Acad Sci USA* 1991; 88: 688-92.
- [156] Poss KD, Tonegawa S. Reduced stress defense in heme oxygenase 1-deficient cells. *Proc Natl Acad Sci USA* 1997; 94: 10925-30.
- [157] Fang J, Akaike T, Maeda H. Antiapoptotic role of heme oxygenase (HO) and the potential of HO as a target in anticancer treatment. *Apoptosis* 2004; 9: 27-35.
- [158] Busserolles J, Megias J, Terencio MC, Alcaraz MJ. Heme oxygenase-1 inhibits apoptosis in Caco-2 cells via activation of Akt pathway. *Int J Biochem Cell Biol* 2006; 38: 1510-7.
- [159] Park JH, Kim EJ, Jang HY, *et al.* Combination treatment with arsenic trioxide and sulindac enhances apoptotic cell death in lung cancer cells via activation of oxidative stress and mitogen-activated protein kinases. *Oncol Rep* 2008; 20: 379-84.
- [160] Kim HR, Kim S, Kim EJ, *et al.* Suppression of Nrf2-driven heme oxygenase-1 enhances the chemosensitivity of lung cancer A549 cells toward cisplatin. *Lung Cancer* 2008; 60: 47-56.
- [161] Park MK, Kang YJ, Ha YM, *et al.* EP2 receptor activation by prostaglandin E2 leads to induction of HO-1 via PKA and PI3K pathways in C6 cells. *Biochem Biophys Res Commun* 2009; 379: 1043-7.
- [162] Rushworth SA, MacEwan DJ. HO-1 underlies resistance of AML cells to TNF-induced apoptosis. *Blood* 2008; 111: 3793-801.
- [163] Nagai T, Kikuchi S, Ohmine K, *et al.* Hemin reduces cellular sensitivity to imatinib and anthracyclins via Nrf2. *J Cell Biochem* 2008; 104: 680-91.
- [164] Fang J, Sawa T, Akaike T, Greish K, Maeda H. Enhancement of chemotherapeutic response of tumor cells by a heme oxygenase inhibitor, pegylated zinc protoporphyrin. *Int J Cancer* 2004; 109: 1-8.
- [165] Mayerhofer M, Gleixner KV, Mayerhofer J, *et al.* Targeting of heat shock protein 32 (Hsp32)/heme oxygenase-1 (HO-1) in leukemic cells in chronic myeloid leukemia: a novel approach to overcome resistance against imatinib. *Blood* 2008; 111: 2200-10.
- [166] Reghehy M, Greish K, Rancan F, *et al.* Water-soluble polymer conjugates of ZnPP for photodynamic tumor therapy. *Bioconjug Chem* 2007; 18: 494-9.
- [167] Frank J, Lornejad-Schafer MR, Schoffl H, *et al.* Inhibition of heme oxygenase-1 increases responsiveness of melanoma cells to ALA-based photodynamic therapy. *Int J Oncol* 2007; 31: 1539-45.
- [168] Tanaka S, Akaike T, Fang J, *et al.* Antiapoptotic effect of haem oxygenase-1 induced by nitric oxide in experimental solid tumour. *Br J Cancer* 2003; 88: 902-9.
- [169] Shi L, Fang J. Implication of heme oxygenase-1 in the sensitivity of nasopharyngeal carcinomas to radiotherapy. *J Exp Clin Cancer Res* 2008; 27: 13-9.
- [170] Sassa S, Nagai T. The role of heme in gene expression. *Int J Hematol* 1996; 63: 167-78.
- [171] Moi P, Chan K, Asunis I, Cao A, Kan YW. Isolation of NF-E2-related factor 2 (Nrf2), a NF-E2-like basic leucine zipper transcriptional activator that binds to the tandem NF-E2/AP1 repeat of the beta-globin locus control region. *Proc Natl Acad Sci USA* 1994; 91: 9926-30.
- [172] Oyake T, Itoh K, Motohashi H, *et al.* Bach proteins belong to a novel family of BTB-basic leucine zipper transcription factors that interact with MafK and regulate transcription through the NF-E2 site. *Mol Cell Biol* 1996; 16: 6083-95.
- [173] Dhakshinamoorthy S, Jain AK, Bloom DA, Jaiswal AK. Bach1 competes with Nrf2 leading to negative regulation of the antioxidant response element (ARE)-mediated NAD(P)H:quinone oxidoreductase 1 gene expression and induction in response to antioxidants. *J Biol Chem* 2005; 280: 16891-900.
- [174] Li CQ, Kim MY, Godoy LC, *et al.* Nitric oxide activation of Keap1/Nrf2 signaling in human colon carcinoma cells. *Proc Natl Acad Sci USA* 2009; 106: 14547-51.
- [175] Bonnovilas ID, Tsiftoglou AS. Hemin counteracts the repression of Bcl-2 and Nrf2 genes and the cell killing induced by imatinib in human Bcr-Abl(+) CML cells. *Oncol Res* 2009; 17: 535-47.
- [176] Ogborne RM, Rushworth SA, O'Connell MA. Epigallocatechin activates haem oxygenase-1 expression via protein kinase Cdelta and Nrf2. *Biochem Biophys Res Commun* 2008; 373: 584-8.
- [177] Kweon MH, Adhami VM, Lee JS, Mukhtar H. Constitutive overexpression of Nrf2-dependent heme oxygenase-1 in A549 cells contributes to resistance to apoptosis induced by epigallocatechin 3-gallate. *J Biol Chem* 2006; 281: 33761-72.

- [178] Zhang J, Hung AC, Ng PY, *et al.* PKCdelta mediates Nrf2-dependent protection of neuronal cells from NO-induced apoptosis. *Biochem Biophys Res Commun* 2009; 386: 750-6.
- [179] Li MH, Jang JH, Na HK, Cha YN, Surh YJ. Carbon monoxide produced by heme oxygenase-1 in response to nitrosative stress induces expression of glutamate-cysteine ligase in PC12 cells via activation of phosphatidylinositol 3-kinase and Nrf2 signaling. *J Biol Chem* 2007; 282: 28577-86.
- [180] Granato A, Gores G, Vilei MT, *et al.* Bilirubin inhibits bile acid induced apoptosis in rat hepatocytes. *Gut* 2003; 52: 1774-8.
- [181] Fondevila C, Shen XD, Tsuchiyashi S, *et al.* Biliverdin therapy protects rat livers from ischemia and reperfusion injury. *Hepatology* 2004; 40: 1333-41.
- [182] Lee HJ, Lee J, Min SK, *et al.* Differential induction of heme oxygenase-1 against nicotine-induced cytotoxicity via the PI3K, MAPK, and NF- κ B pathways in immortalized and malignant human oral keratinocytes. *J Oral Pathol Med* 2008; 37: 278-86.
- [183] Kim DS, Chae SW, Kim HR, Chae HJ. CO and bilirubin inhibit doxorubicin-induced cardiac cell death. *Immunopharmacol Immunotoxicol* 2009; 31: 64-70.
- [184] Nowis D, Bugajski M, Winiarska M, *et al.* Zinc protoporphyrin IX, a heme oxygenase-1 inhibitor, demonstrates potent antitumor effects but is unable to potentiate antitumor effects of chemotherapeutics in mice. *BMC Cancer* 2008; 8: 197.
- [185] Cesaratto L, Calligaris SD, Vascotto C, *et al.* Bilirubin-induced cell toxicity involves PTEN activation through an APE1/Ref-1-dependent pathway. *J Mol Med* 2007; 85: 1099-112.
- [186] Ollinger R, Kogler P, Troppmair J, *et al.* Bilirubin inhibits tumor cell growth via activation of ERK. *Cell Cycle* 2007; 6: 3078-85.
- [187] Cherrington JM, Strawn LM, Shawver LK. New paradigms for the treatment of cancer: the role of anti-angiogenesis agents. *Adv Cancer Res* 2000; 79: 1-38.
- [188] Brouard S, Otterbein LE, Anrather J, *et al.* Carbon monoxide generated by heme oxygenase 1 suppresses endothelial cell apoptosis. *J Exp Med* 2000; 192: 1015-26.
- [189] Tsui TY, Siu YT, Schlitt HJ, Fan ST. Heme oxygenase-1-derived carbon monoxide stimulates adenosine triphosphate generation in human hepatocyte. *Biochem Biophys Res Commun* 2005; 336: 898-902.
- [190] Gunther L, Berberat PO, Haga M, *et al.* Carbon monoxide protects pancreatic β -cells from apoptosis and improves islet function/survival after transplantation. *Diabetes* 2002; 51: 994-9.
- [191] Suliman HB, Carraway MS, Ali AS, *et al.* The CO/HO system reverses inhibition of mitochondrial biogenesis and prevents murine doxorubicin cardiomyopathy. *J Clin Invest* 2007; 117: 3730-41.
- [192] Lee SK, Park DY, Lee HJ, *et al.* Functional interaction between nitric oxide-induced iron homeostasis and heme oxygenase-1 in immortalized and malignant oral keratinocytes. *Cancer Lett* 2007; 249: 283-93.
- [193] Cermak J, Balla J, Jacob HS, *et al.* Tumor cell heme uptake induces ferritin synthesis resulting in altered oxidant sensitivity: possible role in chemotherapy efficacy. *Cancer Res* 1993; 53: 5308-13.
- [194] Baldi A, Lombardi D, Russo P, *et al.* Ferritin contributes to melanoma progression by modulating cell growth and sensitivity to oxidative stress. *Clin Cancer Res* 2005; 11: 3175-83.
- [195] Kiessling MK, Klemke CD, Kaminski MM, *et al.* Inhibition of constitutively activated nuclear factor- κ B induces reactive oxygen species- and iron-dependent cell death in cutaneous T-cell lymphoma. *Cancer Res* 2009; 69: 2365-74.
- [196] Lin F, Girotti AW. Hyperresistance of leukemia cells to photodynamic inactivation after long-term exposure to hemin. *Cancer Res* 1996; 56: 4636-43.
- [197] Torti FM, Torti SV. Regulation of ferritin genes and protein. *Blood* 2002; 99: 3505-16.
- [198] Andreadi CK, Howells LM, Atherfold PA, Manson MM. Involvement of Nrf2, p38, B-Raf, and nuclear factor- κ B, but not phosphatidylinositol 3-kinase, in induction of hemoxygenase-1 by dietary polyphenols. *Mol Pharmacol* 2006; 69: 1033-40.
- [199] Liu XM, Chapman GB, Wang H, Durante W. Adenovirus-mediated heme oxygenase-1 gene expression stimulates apoptosis in vascular smooth muscle cells. *Circulation* 2002; 105: 79-84.
- [200] David H. Rudolf Virchow and modern aspects of tumor pathology. *Pathol Res Pract* 1988; 183: 356-64.
- [201] Rijhsinghani K, Greenblatt M, Shubik P. Vascular abnormalities induced by benzo[a]pyrene: an *in vivo* study in the hamster cheek pouch. *J Natl Cancer Inst* 1968; 41: 205-16.
- [202] Ehrmann RL, Knott M. Choriocarcinoma. Transfilter stimulation of vasoproliferation in the hamster cheek pouch. Studied by light and electron microscopy. *J Natl Cancer Inst* 1968; 41: 1329-41.
- [203] Folkman J. Tumor angiogenesis: therapeutic implications. *N Engl J Med* 1971; 285: 1182-6.
- [204] Cisowski J, Loboda A, Jozkowicz A, *et al.* Role of heme oxygenase-1 in hydrogen peroxide-induced VEGF synthesis: effect of HO-1 knockout. *Biochem Biophys Res Commun* 2005; 326: 670-6.
- [205] Jazwa A, Loboda A, Golda S, *et al.* Effect of heme and heme oxygenase-1 on vascular endothelial growth factor synthesis and angiogenic potency of human keratinocytes. *Free Radic Biol Med* 2006; 40: 1250-63.
- [206] Dulak J, Jozkowicz A, Foresti R, *et al.* Heme oxygenase activity modulates vascular endothelial growth factor synthesis in vascular smooth muscle cells. *Antioxid Redox Signal* 2002; 4: 229-40.
- [207] Suzuki M, Iso-o N, Takeshita S, *et al.* Facilitated angiogenesis induced by heme oxygenase-1 gene transfer in a rat model of hindlimb ischemia. *Biochem Biophys Res Commun* 2003; 302: 138-43.
- [208] Martin MJ, Tanos T, Garcia AB, *et al.* The Galpha12/13 family of heterotrimeric G proteins and the small GTPase RhoA link the Kaposi sarcoma-associated herpes virus G protein-coupled receptor to heme oxygenase-1 expression and tumorigenesis. *J Biol Chem* 2007; 282: 34510-24.
- [209] Kim DH, Kim JH, Kim EH, *et al.* 15-Deoxy- $\Delta^{12,14}$ -prostaglandin- J_2 upregulates the expression of heme oxygenase-1 and subsequently matrix metalloproteinase-1 in human breast cancer cells: possible roles of iron and ROS. *Carcinogenesis* 2009; 30: 645-54.
- [210] Moghaddam A, Zhang HT, Fan TP, *et al.* Thymidine phosphorylase is angiogenic and promotes tumor growth. *Proc Natl Acad Sci USA* 1995; 92: 998-1002.
- [211] Sengupta S, Sellers LA, Matheson HB, Fan TP. Thymidine phosphorylase induces angiogenesis *in vivo* and *in vitro*: an evaluation of possible mechanisms. *Br J Pharmacol* 2003; 139: 219-31.
- [212] Malinda KM, Goldstein AL, Kleinman HK. Thymosin- β_4 stimulates directional migration of human umbilical vein endothelial cells. *FASEB J* 1997; 11: 474-81.
- [213] Brown NS, Jones A, Fujiyama C, Harris AL, Bicknell R. Thymidine phosphorylase induces carcinoma cell oxidative stress and promotes secretion of angiogenic factors. *Cancer Res* 2000; 60: 6298-302.
- [214] Li W, Tanaka K, Morioka K, *et al.* Thymidine phosphorylase gene transfer inhibits vascular smooth muscle cell proliferation by upregulating heme oxygenase-1 and p27KIP1. *Arterioscler Thromb Vasc Biol* 2005; 25: 1370-5.
- [215] Maggiora P, Gambarotta G, Olivero M, *et al.* Control of invasive growth by the HGF receptor family. *J Cell Physiol* 1997; 173: 183-6.
- [216] Horiguchi N, Takayama H, Toyoda M, *et al.* Hepatocyte growth factor promotes hepatocarcinogenesis through c-Met autocrine activation and enhanced angiogenesis in transgenic mice treated with diethylnitrosamine. *Oncogene* 2002; 21: 1791-9.
- [217] Tacchini L, Dansi P, Matteucci E, Desiderio MA. Hepatocyte growth factor signalling stimulates hypoxia inducible factor-1 (HIF-1) activity in HepG2 hepatoma cells. *Carcinogenesis* 2001; 22: 1363-71.
- [218] Hillen F, Griffioen AW. Tumour vascularization: sprouting angiogenesis and beyond. *Cancer Metastasis Rev* 2007; 26: 489-502.
- [219] Duda DG, Cohen KS, Kozin SV, *et al.* Evidence for incorporation of bone marrow-derived endothelial cells into perfused blood vessels in tumors. *Blood* 2006; 107: 2774-6.
- [220] Lyden D, Hattori K, Dias S, *et al.* Impaired recruitment of bone-marrow-derived endothelial and hematopoietic precursor cells blocks tumor angiogenesis and growth. *Nat Med* 2001; 7: 1194-201.
- [221] Aghi M, Cohen KS, Klein RJ, Scadden DT, Chiocca EA. Tumor stromal-derived factor-1 recruits vascular progenitors to mitotic neovasculature, where microenvironment influences their differentiated phenotypes. *Cancer Res* 2006; 66: 9054-64.

- [222] Reddy K, Zhou Z, Jia SF, *et al.* Stromal cell-derived factor-1 stimulates vasculogenesis and enhances Ewing's sarcoma tumor growth in the absence of vascular endothelial growth factor. *Int J Cancer* 2008; 123: 831-7.
- [223] Deshane J, Chen S, Caballero S, *et al.* Stromal cell-derived factor 1 promotes angiogenesis via a heme oxygenase 1-dependent mechanism. *J Exp Med* 2007; 204: 605-18.
- [224] Lin HH, Chen YH, Chang PF, *et al.* Heme oxygenase-1 promotes neovascularization in ischemic heart by coinduction of VEGF and SDF-1. *J Mol Cell Cardiol* 2008; 45: 44-55.
- [225] Lin HH, Chen YH, Yet SF, Chau LY. After vascular injury, heme oxygenase-1/carbon monoxide enhances re-endothelialization via promoting mobilization of circulating endothelial progenitor cells. *J Thromb Haemost* 2009; 7: 1401-8.
- [226] Price JT, Thompson EW. Mechanisms of tumour invasion and metastasis: emerging targets for therapy. *Expert Opin Ther Targets* 2002; 6: 217-33.
- [227] Lee SS, Yang SF, Tsai CH, *et al.* Upregulation of heme oxygenase-1 expression in areca-quid-chewing-associated oral squamous cell carcinoma. *J Formos Med Assoc* 2008; 107: 355-63.
- [228] Lin CW, Shen SC, Hou WC, Yang LY, Chen YC. Heme oxygenase-1 inhibits breast cancer invasion via suppressing the expression of matrix metalloproteinase-9. *Mol Cancer Ther* 2008; 7: 1195-206.
- [229] Desmard M, Amara N, Lanone S, Motterlini R, Boczkowski J. Carbon monoxide reduces the expression and activity of matrix metalloproteinases 1 and 2 in alveolar epithelial cells. *Cell Mol Biol (Noisy-le-grand)* 2005; 51: 403-8.
- [230] Megias J, Busserolles J, Alcaraz MJ. The carbon monoxide-releasing molecule CORM-2 inhibits the inflammatory response induced by cytokines in Caco-2 cells. *Br J Pharmacol* 2007; 150: 977-86.
- [231] Lee TS, Chau LY. Heme oxygenase-1 mediates the anti-inflammatory effect of interleukin-10 in mice. *Nat Med* 2002; 8: 240-6.
- [232] Vincenti MP, Brinckerhoff CE. Signal transduction and cell-type specific regulation of matrix metalloproteinase gene expression: can MMPs be good for you? *J Cell Physiol* 2007; 213: 355-64.
- [233] Becker JC, Fukui H, Imai Y, *et al.* Colonic expression of heme oxygenase-1 is associated with a better long-term survival in patients with colorectal cancer. *Scand J Gastroenterol* 2007; 42: 852-8.
- [234] Ishikawa T, Yoshida N, Higashihara H, *et al.* Different effects of constitutive nitric oxide synthase and heme oxygenase on pulmonary or liver metastasis of colon cancer in mice. *Clin Exp Metastasis* 2003; 20: 445-550.
- [235] Yanagawa T, Omura K, Harada H, *et al.* Heme oxygenase-1 expression predicts cervical lymph node metastasis of tongue squamous cell carcinomas. *Oral Oncol* 2004; 40: 21-7.
- [236] Lin SC, Liu CJ, Yeh WI, *et al.* Functional polymorphism in NFKB1 promoter is related to the risks of oral squamous cell carcinoma occurring on older male areca (betel) chewers. *Cancer Lett* 2006; 243: 47-54.
- [237] Willis D, Moore AR, Frederick R, Willoughby DA. Heme oxygenase: a novel target for the modulation of the inflammatory response. *Nat Med* 1996; 2: 87-90.
- [238] Kapturczak MH, Wasserfall C, Brusko T, *et al.* Heme oxygenase-1 modulates early inflammatory responses: evidence from the heme oxygenase-1-deficient mouse. *Am J Pathol* 2004; 165: 1045-53.
- [239] Sawle P, Foresti R, Mann BE, *et al.* Carbon monoxide-releasing molecules (CO-RMs) attenuate the inflammatory response elicited by lipopolysaccharide in RAW264.7 murine macrophages. *Br J Pharmacol* 2005; 145: 800-10.
- [240] Morse D, Pischke SE, Zhou Z, *et al.* Suppression of inflammatory cytokine production by carbon monoxide involves the JNK pathway and AP-1. *J Biol Chem* 2003; 278: 36993-8.
- [241] Ryter SW, Alam J, Choi AM. Heme oxygenase-1/carbon monoxide: from basic science to therapeutic applications. *Physiol Rev* 2006; 86: 583-650.
- [242] Song R, Ning W, Liu F, *et al.* Regulation of IL-1 β -induced GM-CSF production in human airway smooth muscle cells by carbon monoxide. *Am J Physiol Lung Cell Mol Physiol* 2003; 284: L50-6.
- [243] Wang XM, Kim HP, Song R, Choi AM. Caveolin-1 confers antiinflammatory effects in murine macrophages via the MKK3/p38 MAPK pathway. *Am J Respir Cell Mol Biol* 2006; 34: 434-42.
- [244] Wang XM, Kim HP, Nakahira K, Ryter SW, Choi AM. The heme oxygenase-1/carbon monoxide pathway suppresses TLR4 signaling by regulating the interaction of TLR4 with caveolin-1. *J Immunol* 2009; 182: 3809-18.
- [245] Overhaus M, Moore BA, Barbato JE, *et al.* Biliverdin protects against polymicrobial sepsis by modulating inflammatory mediators. *Am J Physiol Gastrointest Liver Physiol* 2006; 290: G695-703.
- [246] Hayashi S, Takamiya R, Yamaguchi T, *et al.* Induction of heme oxygenase-1 suppresses venular leukocyte adhesion elicited by oxidative stress: role of bilirubin generated by the enzyme. *Circ Res* 1999; 85: 663-771.
- [247] Ishikawa K, Navab M, Leitinger N, Fogelman AM, Lusis AJ. Induction of heme oxygenase-1 inhibits the monocyte transmigration induced by mildly oxidized LDL. *J Clin Invest* 1997; 100: 1209-16.
- [248] Lin CC, Liu XM, Peyton K, *et al.* Far infrared therapy inhibits vascular endothelial inflammation via the induction of heme oxygenase-1. *Arterioscler Thromb Vasc Biol* 2008; 28: 739-45.
- [249] Basiglio CL, Arriaga SM, Pelusa F, *et al.* Complement activation and disease: protective effects of hyperbilirubinaemia. *Clin Sci (Lond)* 2009; 118: 99-113.
- [250] Lin EY, Pollard JW. Role of infiltrated leucocytes in tumour growth and spread. *Br J Cancer* 2004; 90: 2053-8.
- [251] Mantovani A, Romero P, Palucka AK, Marincola FM. Tumour immunity: effector response to tumour and role of the microenvironment. *Lancet* 2008; 371: 771-83.
- [252] Mantovani A, Sozzani S, Locati M, Allavena P, Sica A. Macrophage polarization: tumor-associated macrophages as a paradigm for polarized M2 mononuclear phagocytes. *Trends Immunol* 2002; 23: 549-55.
- [253] Allavena P, Sica A, Solinas G, Porta C, Mantovani A. The inflammatory micro-environment in tumor progression: the role of tumor-associated macrophages. *Crit Rev Oncol Hematol* 2008; 66: 1-9.
- [254] Liu VC, Wong LY, Jang T, *et al.* Tumor evasion of the immune system by converting CD4+CD25- T cells into CD4+CD25+ T regulatory cells: role of tumor-derived TGF β . *J Immunol* 2007; 178: 2883-92.
- [255] Nishikawa H, Kato T, Tanida K, *et al.* CD4+ CD25+ T cells responding to serologically defined autoantigens suppress antitumor immune responses. *Proc Natl Acad Sci USA* 2003; 100: 10902-6.
- [256] Curiel TJ. Regulatory T cells and treatment of cancer. *Curr Opin Immunol* 2008; 20: 241-6.
- [257] Zou W. Regulatory T cells, tumour immunity and immunotherapy. *Nat Rev Immunol* 2006; 6: 295-307.
- [258] Weis N, Weigert A, von Knethen A, Brune B. Heme oxygenase-1 contributes to an alternative macrophage activation profile induced by apoptotic cell supernatants. *Mol Biol Cell* 2009; 20: 1280-8.
- [259] Yamashita K, Ollinger R, McDaid J, *et al.* Heme oxygenase-1 is essential for and promotes tolerance to transplanted organs. *Faseb J* 2006; 20: 776-8.
- [260] McDaid J, Yamashita K, Chora A, *et al.* Heme oxygenase-1 modulates the allo-immune response by promoting activation-induced cell death of T cells. *FASEB J* 2005; 19: 458-60.
- [261] Walker MR, Kasproicz DJ, Gersuk VH, *et al.* Induction of FoxP3 and acquisition of T regulatory activity by stimulated human CD4+CD25- T cells. *J Clin Invest* 2003; 112: 1437-43.
- [262] Chauveau C, Remy S, Royer PJ, *et al.* Heme oxygenase-1 expression inhibits dendritic cell maturation and proinflammatory function but conserves IL-10 expression. *Blood* 2005; 106: 1694-702.
- [263] Ricchetti GA, Williams LM, Foxwell BM. Heme oxygenase 1 expression induced by IL-10 requires STAT-3 and phosphoinositol-3 kinase and is inhibited by lipopolysaccharide. *J Leukoc Biol* 2004; 76: 719-26.
- [264] Braudeau C, Bouchet D, Tesson L, *et al.* Induction of long-term cardiac allograft survival by heme oxygenase-1 gene transfer. *Gene Ther* 2004; 11: 701-10.
- [265] Andersen MH, Sorensen RB, Brimnes MK, *et al.* Identification of heme oxygenase-1-specific regulatory CD8+ T cells in cancer patients. *J Clin Invest* 2009; 119: 2245-56.
- [266] Kimpara T, Takeda A, Watanabe K, *et al.* Microsatellite polymorphism in the human heme oxygenase-1 gene promoter and

- its application in association studies with Alzheimer and Parkinson disease. *Hum Genet* 1997; 100: 145-7.
- [267] Hirai H, Kubo H, Yamaya M, *et al.* Microsatellite polymorphism in heme oxygenase-1 gene promoter is associated with susceptibility to oxidant-induced apoptosis in lymphoblastoid cell lines. *Blood* 2003; 102: 1619-21.
- [268] Taha H, Skrzypek K, Guevara I, *et al.* Role of heme oxygenase-1 in human endothelial cells – lesson from the promoter allelic variants. *Arterioscler Thromb Vasc Biol* 2010; 30: 1634-41.
- [269] Kikuchi A, Yamaya M, Suzuki S, *et al.* Association of susceptibility to the development of lung adenocarcinoma with the heme oxygenase-1 gene promoter polymorphism. *Hum Genet* 2005; 116: 354-60.
- [270] Chang KW, Lee TC, Yeh WI, *et al.* Polymorphism in heme oxygenase-1 (HO-1) promoter is related to the risk of oral squamous cell carcinoma occurring on male areca chewers. *Br J Cancer* 2004; 91: 1551-5.
- [271] Hu JL, Li ZY, Liu W, *et al.* Polymorphism in heme oxygenase-1 (HO-1) promoter and alcohol are related to the risk of esophageal squamous cell carcinoma on Chinese males. *Neoplasma* 2010; 57: 86-92.
- [272] Hong CC, Ambrosone CB, Ahn J, *et al.* Genetic variability in iron-related oxidative stress pathways (Nrf2, NQO1, NOS3, and HO-1), iron intake, and risk of postmenopausal breast cancer. *Cancer Epidemiol Biomarkers Prev* 2007; 16: 1784-94.
- [273] Lo SS, Lin SC, Wu CW, *et al.* Heme oxygenase-1 gene promoter polymorphism is associated with risk of gastric adenocarcinoma and lymphovascular tumor invasion. *Ann Surg Oncol* 2007; 14: 2250-6.
- [274] Okamoto I, Krogler J, Endler G, *et al.* A microsatellite polymorphism in the heme oxygenase-1 gene promoter is associated with risk for melanoma. *Int J Cancer* 2006; 119: 1312-5.
- [275] Sawa T, Mounawar M, Tatemichi M, *et al.* Increased risk of gastric cancer in Japanese subjects is associated with microsatellite polymorphisms in the heme oxygenase-1 and the inducible nitric oxide synthase gene promoters. *Cancer Lett* 2008; 269: 78-84.
- [276] Glei M, Klenow S, Sauer J, *et al.* Hemoglobin and heme induce DNA damage in human colon tumor cells HT29 clone 19A and in primary human colonocytes. *Mutat Res* 2006; 594: 162-71.
- [277] Pierre F, Tache S, Petit CR, Van der Meer R, Corpet DE. Meat and cancer: haemoglobin and heme in a low-calcium diet promote colorectal carcinogenesis at the aberrant crypt stage in rats. *Carcinogenesis* 2003; 24: 1683-90.
- [278] Ghattas MH, Chuang LT, Kappas A, Abraham NG. Protective effect of HO-1 against oxidative stress in human hepatoma cell line (HepG2) is independent of telomerase enzyme activity. *Int J Biochem Cell Biol* 2002; 34: 1619-28.
- [279] Genter MB, Burman DM, Vijayakumar S, Ebert CL, Aronow BJ. Genomic analysis of alachlor-induced oncogenesis in rat olfactory mucosa. *Physiol Genomics* 2002; 12: 35-45.
- [280] Caballero F, Meiss R, Gimenez A, Batlle A, Vazquez E. Immunohistochemical analysis of heme oxygenase-1 in preneoplastic and neoplastic lesions during chemical hepatocarcinogenesis. *Int J Exp Pathol* 2004; 85: 213-22.
- [281] Yan Y, Higashi K, Yamamura K, *et al.* Different responses other than the formation of DNA-adducts between the livers of carcinogen-resistant rats (DRH) and carcinogen-sensitive rats (Donryu) to 3'-methyl-4-dimethylaminoazobenzene administration. *Jpn J Cancer Res* 1998; 89: 806-13.
- [282] Matsumoto A, Hanayama R, Nakamura M, *et al.* A high expression of heme oxygenase-1 in the liver of LEC rats at the stage of hepatoma: the possible implication of induction in uninvolved tissue. *Free Radic Res* 1998; 28: 383-91.
- [283] Henry F, Bretaudeau L, Barbieux I, Meflah K, Gregoire M. Induction of antigen presentation by macrophages after phagocytosis of tumour apoptotic cells. *Res Immunol* 1998; 149: 673-9.
- [284] Fang J, Sawa T, Akaike T, *et al.* *In vivo* antitumor activity of pegylated zinc protoporphyrin: targeted inhibition of heme oxygenase in solid tumor. *Cancer Res* 2003; 63: 3567-74.
- [285] Sahoo SK, Sawa T, Fang J, *et al.* Pegylated zinc protoporphyrin: a water-soluble heme oxygenase inhibitor with tumor-targeting capacity. *Bioconjug Chem* 2002; 13: 1031-8.
- [286] Matsumura Y, Maeda H. A new concept for macromolecular therapeutics in cancer chemotherapy: mechanism of tumorotropic accumulation of proteins and the antitumor agent smancs. *Cancer Res* 1986; 46: 6387-92.
- [287] Iyer AK, Greish K, Seki T, *et al.* Polymeric micelles of zinc protoporphyrin for tumor targeted delivery based on EPR effect and singlet oxygen generation. *J Drug Target* 2007; 15: 496-506.
- [288] Hadzijušević E, Rebutti L, Gleixner KV, *et al.* Targeting of heat-shock protein 32/heme oxygenase-1 in canine mastocytoma cells is associated with reduced growth and induction of apoptosis. *Exp Hematol* 2008; 36: 1461-70.
- [289] Schiwon K, Brauer HD, Gerlach B, Müller CM, Montforts FP. Potential photosensitizers for photodynamic therapy. IV. Photophysical and photochemical properties of azaporphyrin and azachlorin derivatives. *J Photochem Photobiol B* 1994; 23: 239-43.
- [290] Jozkowicz A, Dulak J. Effects of protoporphyrins on production of nitric oxide and expression of vascular endothelial growth factor in vascular smooth muscle cells and macrophages. *Acta Biochim Pol* 2003; 50: 69-79.
- [291] Loboda A, Jazwa A, Wegiel B, Jozkowicz A, Dulak J. Heme oxygenase-1-dependent and -independent regulation of angiogenic genes expression: effect of cobalt protoporphyrin and cobalt chloride on VEGF and IL-8 synthesis in human microvascular endothelial cells. *Cell Mol Biol (Noisy-le-grand)* 2005; 51: 347-55.
- [292] Khelifi AF, Prise VE, Tozer GM. Effects of tin-protoporphyrin IX on blood flow in a rat tumor model. *Exp Biol Med (Maywood)* 2003; 228: 481-5.
- [293] Tozer GM, Prise VE, Motterlini R, *et al.* The comparative effects of the NOS inhibitor, Nomega-nitro-L-arginine, and the haemoxygenase inhibitor, zinc protoporphyrin IX, on tumour blood flow. *Int J Radiat Oncol Biol Phys* 1998; 42: 849-53.
- [294] Tiemann K, Rossi JJ. RNAi-based therapeutics-current status, challenges and prospects. *EMBO Mol Med* 2009; 1: 142-51.
- [295] Xu CX, Jere D, Jin H, *et al.* Poly(ester amine)-mediated, aerosol-delivered Akt1 small interfering RNA suppresses lung tumorigenesis. *Am J Respir Crit Care Med* 2008; 178: 60-73.
- [296] Jackson AL, Bartz SR, Schelter J, *et al.* Expression profiling reveals off-target gene regulation by RNAi. *Nat Biotechnol* 2003; 21: 635-7.
- [297] Jackson AL, Burchard J, Schelter J, *et al.* Widespread siRNA "off-target" transcript silencing mediated by seed region sequence complementarity. *RNA* 2006; 12: 1179-87.
- [298] Kleinman ME, Yamada K, Takeda A, *et al.* Sequence- and target-independent angiogenesis suppression by siRNA via TLR3. *Nature* 2008; 452: 591-7.
- [299] Hornung V, Guenther-Biller M, Bourquin C, *et al.* Sequence-specific potent induction of IFN α by short interfering RNA in plasmacytoid dendritic cells through TLR7. *Nat Med* 2005; 11: 263-70.
- [300] Judge A, MacLachlan I. Overcoming the innate immune response to small interfering RNA. *Hum Gene Ther* 2008; 19: 111-24.
- [301] Jackson AL, Burchard J, Leake D, *et al.* Position-specific chemical modification of siRNAs reduces "off-target" transcript silencing. *Rna* 2006; 12: 1197-205.
- [302] Grimm D, Streetz KL, Jopling CL, *et al.* Fatality in mice due to oversaturation of cellular microRNA/short hairpin RNA pathways. *Nature* 2006; 441: 537-41.