

Essential fatty acids as possible mediators of the actions of statins

U. N. Das

EFA Sciences LLC, Norwood, MA, USA

Summary Statins and polyunsaturated fatty acids have similar actions: both enhance endothelial nitric oxide synthesis, inhibit the production of pro-inflammatory cytokines, lower cholesterol levels, prevent atherosclerosis and are of benefit in coronary heart disease, stroke and osteoporosis. Statins enhance the conversion of linoleic acid and eicosapentaenoic acid to their long chain derivatives. Animals with essential fatty acid deficiency show an increase in HMG-CoA reductase activity, which reverts to normalcy following topical application of linoleic acid. Similarly to statins, polyunsaturated fatty acids also inhibit HMG-CoA reductase activity. In view of the similarity in their actions and as statins influence essential fatty acid metabolism, it is suggested that essential fatty acids and their metabolites may serve as second messengers of the actions of statins. © 2001 Harcourt Publishers Ltd

INTRODUCTION

HMG-CoA reductase inhibitors or statins inhibit the rate-limiting step in cholesterol biosynthesis and hence are used to lower cholesterol and prevent the development of atherosclerosis.¹ Recently, it has been recognized that statins are of benefit to patients with myocardial infarction even when they have no effect on cholesterol levels or formation of atherosclerosis,^{4,5} prevent stroke,² and increase bone-mineral density in post-menopausal women.³ These multiple actions of statins have been ascribed to their ability to enhance endothelial nitric oxide synthesis,⁶ inhibit the production of pro-inflammatory cytokines, tumor necrosis factor α (TNF α)⁷ and interleukin-6 (IL-6),⁸ and C-reactive protein,⁹ plasminogen activator inhibitor type-1 (PAI-1),¹⁰ and monocyte chemotactic protein-1 (MCP-1).¹¹ Statins can increase the expression of bone morphogenetic protein 2 (BMP-2) and thus increase bone formation, even in postmenopausal women.¹² In addition, statins inhibit cellular proliferation, induce apoptosis of tumor cells in vitro, though they are of little use as anti-tumor agents in vivo in

view of their selective hepatic uptake and low systemic availability.¹³

EFAS HAVE ACTIONS SIMILAR TO STATINS

It is interesting to note that essential fatty acids (EFAs) and their metabolites such as gamma-linolenic acid (GLA, 18:3n-6), dihomog-LA (20:3n-6), arachidonic acid (AA, 20:4n-6), eicosapentaenoic acid (EPA, 20:5n-3) and docosahexaenoic acid (DHA, 22:6n-3) also have actions similar to statins. EFAs, linoleic acid (LA, 18:2n-6) and alpha-linolenic acid (ALA, 18:3n-3), are widely distributed in our diet.¹⁴ Both LA and ALA are converted to their respective long-chain polyunsaturated fatty acids (PUFAs) by the enzymes: δ -6- and δ -5-desaturases. DGLA, AA and EPA form precursors to 1, 2 and 3 series of eicosanoids respectively. Similarly to statins, PUFAs can lower cholesterol, triglyceride and low density lipoprotein levels,¹⁵ block HMG-CoA reductase activity^{16,17} and enhance NO synthesis from the endothelial cells.^{18,19} Marine PUFAs protect against ischemic heart disease,^{20,21} inhibit platelet aggregation²² and smooth muscle cell proliferation and thus, prevent atherosclerosis.²³ Similarly to statins, EFAs/PUFAs are useful in the prevention and treatment of osteoporosis,²⁴ and suppress the production of pro-inflammatory cytokines.^{25,26} In view of the similarity in their actions, is it possible that statins and EFAs interact with each other?

Received 16 January 2001

Accepted 7 February 2001

Correspondence to: Dr U. N. Das, MD, FAMS, EFA Sciences LLC, 1420 Providence Highway, Suite #266, Norwood, MA 02062, USA. Phone: 781-278-9919; Fax: 781-278-9959; E-mail: undurti@hotmail.com

EFAS AND STATINS

It has been shown that statins can enhance the conversion of LA and EPA to their long chain PUFA derivatives.²⁷ Animals with EFA deficiency showed an increase in HMG-CoA reductase activity, which reverted to normalcy following topical application of LA.²⁸ Further, PUFAs can inhibit HMG CoA reductase activity.^{29–31} Hence, I suggest that statins bring about some, if not all, of their actions by interacting with EFAs and their metabolites and by augmenting EFA metabolism. It is possible that EFAs and their metabolites may serve as endogenous second messengers of statins.

In view of this, PUFAs and statins may form a potent combination in the treatment of a variety of disorders including, but not limited to, hyperlipidemias, atherosclerosis, stroke, ischaemic heart disease, osteoporosis, and possibly cancer. Such a combination may suppress inflammation and autoimmune disorders such as rheumatoid arthritis and systemic lupus erythematosus in view of their action on pro-inflammatory cytokines. A combination of PUFAs and statins may particularly be useful in the treatment of cancer. This is so since mevalonate is not only the precursor of cholesterol but also of a variety of isoprenoid containing compounds, which are necessary for the posttranslational lipid modification (prenylation) and hence, the function of *ras* and other small GTPases.¹³ Chapkin et al.³² showed that n-3 PUFAs are capable of suppressing carcinogen induced *ras* activation in the colon prior to overt neoplasia. Furthermore, various PUFAs can selectively kill tumor cells by a free radical dependent process,^{33–36} and by suppressing *Bcl-2* expression,³⁷ an anti-apoptotic gene.

Recently, it was suggested that statins can substantially lower the risk of developing dementia including Alzheimer's disease.³⁸ If this is true, even EFAs and their metabolites should also be useful in Alzheimer's disease. Yehuda and Carasso³⁹ showed that preparations with w3-to-w6 ratios ranging from 1:3.5 to 1:5 produced significant favourable effects on learning, pain thresholds and thermoregulation. On the basis of these results, it was opined that the beneficial effects of the PUFAs can be related to optimal incorporation of w3 fatty acids into the brain membranes. The presence of appropriate amounts of PUFAs in the neuronal membranes is important for their structural integrity, membrane fluidity, and membrane-related functions such as receptor, enzyme, and ion channel kinetics, as well as eicosanoid functions (reviewed in 39). It is also known that arachidonic acid and linolenic acid stimulate glucose uptake in cerebral cortical astrocytes.⁴⁰ Furthermore, hippocampal acetylcholine release during memory testing in rats can be potentiated by glucose.⁴¹ This suggests that AA and ALA

and, possibly, other PUFAs can improve memory by enhancing glucose uptake by neuronal cells, which in turn augments acetylcholine release, a molecule that is known to be involved in memory formation. This coupled with the observation that PUFAs are potent neuroprotectors,⁴² DHA prevents neuronal apoptosis,⁴³ and low serum DHA is a significant risk factor for Alzheimer's disease,⁴⁴ indicates that provision of adequate amounts of various PUFAs (in the right proportion) can also prevent Alzheimer's disease and other dementias similarly to statins. Hence, it is likely that the beneficial effect of statins in the prevention of Alzheimer's disease and other dementias may also be due to their ability to enhance EFA metabolism. Furthermore, both statins and PUFAs have anti-inflammatory actions. Since inflammation is believed to play a major role in Alzheimer's disease, the anti-inflammatory actions of statins and PUFAs may contribute to their beneficial action(s) in Alzheimer's disease and other dementias. It is also known that small GTPases, which are prenylated products of the mevalonate pathway, may have a negative control on the expression of BMP-2 and other BMPs. Thus, inhibition of the mevalonate pathway by statins and PUFAs may prevent the function of small GTPases and enhance the expression of BMPs. It is now known that BMPs have multiple functions in the developing nervous system. Primary cells derived from the septal area of E14 mice when were treated with human recombinant BMP-2, BMP-4, BMP-6, BMP-7, BMP-9 and BMP-12 a several-fold increase in cellular acetylcholine levels was observed.⁴⁵ Further, BMP-9 directly induced the expression of the cholinergic gene locus encoding choline acetyltransferase and the vesicular acetylcholine transporter and up-regulated acetylcholine synthesis.⁴⁵ It is known that DHA increases cerebral acetylcholine levels and improves learning ability in rats.⁴⁶ It is possible that PUFAs are able to enhance brain acetylcholine levels by increasing the formation of BMPs as a result of their action on HMG-CoA reductase activity. Thus, it is suggested that both statins and PUFAs are able to improve memory and prevent Alzheimer's disease by augmenting the concentrations of BMPs in the brain.

In view of the similarity in their actions, it is suggested that combined administration of statins and PUFAs in the right proportion may prove to be highly beneficial in a variety of clinical conditions. Preliminary evidence for such potentiation of the action of statins by PUFAs in the treatment hyperlipidemias is already available.^{47,48}

REFERENCES

1. Goldstein J. L., Brown M. S. Regulation of the mevalonate pathway. *Nature* 1990; **343**: 425–430.
2. Sacks F. M., et al. The effect of pravastatin on coronary events after myocardial infarction in patients with average cholesterol

- levels. Cholesterol and Recurrent Events Trial Investigators. *N Engl J Med* 1996; **335**: 1001–1009.
3. Lefer A. M., et al. Simvastatin preserves the ischemia-reperfused myocardium in normocholesterolemic rat hearts. *Circulation* 1999; **100**: 178–184.
 4. Rosendorff C. Statins for prevention of stroke. *Lancet* 1998; **351**: 1002–1003.
 5. Edwards C. J., Hart D. J., Spector T. D. Oral statins and increased bone mineral density in postmenopausal women. *Lancet* 2000; **355**: 2218–2219.
 6. Kureishi Y., et al. The HMG-CoA reductase inhibitor simvastatin activates the protein kinase Akt and promotes angiogenesis in normocholesterolemic animals. *Nature Med* 2000; **6**: 1004–1010.
 7. Rosenson R. S., Tangney C. C., Casey L. C. Inhibition of proinflammatory cytokine production by pravastatin. *Lancet* 1999; **353**: 983–984.
 8. Ikeda U., Shimada K. Statins and monocytes. *Lancet* 1999; **353**: 2070.
 9. Strandberg T. E., Vanhanen H., Tikkanen M. J. Effect of statins on C-reactive protein in patients with coronary artery disease. *Lancet* 1999; **353**: 118–119.
 10. Lopez S., Peiretti F., Bonardo B., Juhan-Vague I., Nalbone G. Effect of atorvastatin and fluvastatin on the expression of plasminogen activator inhibitor type-1 in cultured endothelial cells. *Atherosclerosis* 2000; **152**: 359–366.
 11. Romano M., Diomedea L., Sironi M., et al. Inhibition of monocyte chemotactic protein-1 synthesis by statins. *Lab Invest* 2000; **80**: 1095–1100.
 12. Mundy G., et al. Stimulation of bone formation in vitro and in rodents by statins. *Science* 1999; **286**: 1946–1949.
 13. Rogers M. J. Statins: lower lipids and better bones? *Nature Med* 2000; **6**: 21–23.
 14. Das U. N., et al. Clinical significance of essential fatty acids. *Nutrition* 1988; **4**: 337–341.
 15. Das U. N. Beneficial effect(s) of n-3 fatty acids in cardiovascular diseases: but, why and how? *Prostaglandins Leukotrienes Essen Fatty Acids* 2000; **63**: 351–362.
 16. Watanabe I. M., Umekawa H., Takahashi T., Furuichi Y. Effects of dietary alpha- or gamma-linolenic acid on levels and fatty acid compositions of serum and hepatic lipids, and activity and mRNA abundance of 3-hydroxy-3-methylglutaryl CoA reductase in rats. *Comp Biochem Physiol A Mol Integr Physiol* 1999; **122**: 213–220.
 17. Proksch E., Feingold K. R., Elias F. M. Epidermal HMG CoA reductase activity in essential fatty acid deficiency: barrier requirements rather than eicosanoid generation regulate cholesterol synthesis. *J Invest Dermatol* 1992; **99**: 216–220.
 18. McVeigh G. E., Brennan G. M., Johnson G. D., et al. Dietary fish oil augments nitric oxide production or release in patients with type 2 (non-insulin dependent) diabetes mellitus. *Diabetologia* 1993; **36**: 33–38.
 19. Lawson D. L., Mehta J. L., Saldeen K., Mehta P., Saldeen T. G. Omega-3 polyunsaturated fatty acids augment endothelium dependent vasorelaxation by enhanced release of EDRF and vasodilator prostaglandins. *Eicosanoids* 1991; **4**: 217–223.
 20. Prichard B. N. C., et al. Fish oils and cardiovascular disease. *Br Med J* 1995; **310**: 819–820.
 21. GISSI Prevenzione Investigators. Dietary supplementation with n-3 polyunsaturated fatty acids and vitamin E after myocardial infarction: results of the GISSI-Prevenzione trial. *Lancet* 1999; **354**: 447–455.
 22. Haines A. P., Sanders T. A. B., Imeson J. D., et al. Effects of a fish oil supplement on platelet function, haemostatic variables and albuminuria in insulin-dependent diabetics. *Thromb Res* 1986; **43**: 643–655.
 23. Cornwell D. G., Panganamala R. V. Atherosclerosis: an intracellular deficiency in essential fatty acids. *Prog Lipid Res* 1981; **20**: 365–376.
 24. Das U. N. Essential fatty acids and osteoporosis. *Nutrition* 2000; **16**: 386–390.
 25. Kumar G. S., Das U. N. Effect of prostaglandins and their precursors on the proliferation of human lymphocytes and their secretion of tumor necrosis factor and various interleukins. *Prostaglandins Leukotrienes Essen Fatty Acids* 1994; **50**: 331–334.
 26. Das U. N. Beneficial effect of eicosapentaenoic acid and docosahexaenoic acid in the management of systemic lupus erythematosus and its relationship to the cytokine network. *Prostaglandins Leukotrienes Essen Fatty Acids* 1994; **51**: 207–213.
 27. Risc P., Colombo C., Galli C. Effects of simvastatin on the metabolism of polyunsaturated fatty acids and on glycerolipid, cholesterol, and de novo lipid synthesis in THP-1 cells. *J Lipid Res* 1997; **38**: 1299–1307.
 28. Proksch E., Feingold K. R., Elias F. M. Epidermal HMG CoA reductase activity in essential fatty acid deficiency: barrier requirements rather than eicosanoid generation regulate cholesterol synthesis. *J Invest Dermatol* 1992; **99**: 216–220.
 29. Field F. J., Albright E. J., Mathur S. N. Effect of dietary n-3 fatty acids on HMG-CoA reductase and ACAT activities in liver and intestine of the rabbit. *J Lipid Res* 1987; **28**: 50–58.
 30. Hromadova M., Sebkova E., Klimes I. HMG-CoA reductase activity in the liver of rats with hereditary hypertriglyceridemia: effect of dietary fish oil. *Endocr Regul* 1994; **28**: 211–214.
 31. El-Sohemy A., Archer M. C. Regulation of mevalonate synthesis in rat mammary glands by dietary n-3 and n-6 polyunsaturated fatty acids. *Cancer Res* 1997; **57**: 3685–3687.
 32. Chapkin R. S., Jinag Y. H., Davidson L. A., Lupton J. R. Modulation of intracellular second messengers by dietary fat during colonic tumor development. *Adv Exp Med Biol* 1997; **422**: 85–96.
 33. Das U. N. Tumoricidal action of cis-unsaturated fatty acids and their relationship to free radicals and lipid peroxidation. *Cancer Lett* 1991; **56**: 235–243.
 34. Das U. N., Begin M. E., Ells G., Horrobin D. F. Polyunsaturated fatty acids augment free radical generation in tumor cells in vitro. *Biochem Biophys Res Commun* 1987; **145**: 15–24.
 35. Das U. N., Huang Y. S., Begin M. E., Ells G., Horrobin D. F. Uptake and distribution of cis-unsaturated fatty acids and their effect on free radical generation in normal and tumor cells in vitro. *Free Radical Biol Med* 1987; **3**: 9–14.
 36. Das U. N., Prasad V. V. S. K., Reddy D. R. Local application of gamma-linolenic acid in the treatment of human gliomas. *Cancer Lett* 1995; **94**: 147–155.
 37. Das U. N. Essential fatty acids, lipid peroxidation and apoptosis. *Prostaglandins Leukotrienes Essen Fatty Acids* 1999; **61**: 157–163.
 38. Jick H., Zornberg G. L., Jick S. S., Seshadri S., Drachman D. A. Statins and the risk of dementia. *Lancet* 2000; **356**: 1627–1631.
 39. Yehuda S., Carasso R. L. Modulation of learning, pain thresholds, and thermoregulation in the rat by preparations of free purified α -linolenic and linoleic acids: Determination of the optimal w3-to-w6 ratio. *Proc Natl Acad Sci USA* 1993; **90**: 10345–10349.
 40. Yu N., Martin J.-L., Stella N., Magistretti P. J. Arachidonic acid stimulates glucose uptake in cerebral astrocytes. *Proc Natl Acad Sci USA* 1993; **90**: 4042–4046.
 41. Ragozzino M. E., Unick K. E., Gold P. E. Hippocampal acetylcholine release during memory testing in rats: Potentiation by glucose. *Proc Natl Acad Sci USA* 1996; **93**: 4693–4698.
 42. Lauritzen I., Blondeau N., Heurteaux C., Widmann C., Romey G., Lazdunski M. Polyunsaturated fatty acids are potent neuroprotectors. *EMBO J* 2000; **19**: 1784–1793.

43. Kim H-Y, Akbar M., Lau A., Edsall L. Inhibition of neuronal apoptosis by docosahexaenoic acid (22:6n-3). *J Biol Chem* 2000; **275**: 35 215–35 223.
44. Kyle D. J., Schaefer E., Patton G., Beiser A. Low serum docosahexaenoic acid is a significant risk factor for Alzheimer's dementia. *Lipids* 1999; **34** (Suppl): S245.
45. Coviella I. L., Berse B., Krauss R., Thies R. S., Blusztajn J. K. Induction and maintenance of the neuronal cholinergic phenotype in the central nervous system by BMP-9. *Science* 2000; **289**: 313–316.
46. Minami M., Kimura S., Endo T., Hamaue N., Hirafuji M., Togashi H., Matsumoto M., Yoshioka M., Saito H., Watanabe S., Kobayashi T., Okuyama H. Dietary docosahexaenoic acid increases cerebral acetylcholine levels and improves passive avoidance performance in stroke-prone spontaneously hypertensive rats. *Pharmacol Biochem Behav* 1997; **58**: 1123–1129.
47. Nordoy A., Bonna K. H., Nilsen H., et al. Effects of simvastatin and omega-3 fatty acids on plasma lipoproteins and lipid peroxidation in patients with combined hyperlipidaemia. *J Intern Med* 1998; **243**: 163–170.
48. Nordoy A., Bonna K. H., Sandset P. M., et al. Effect of omega-3 fatty acids and simvastatin on homeostatic risk factors and postprandial hyperlipidemia in patients with combined hyperlipidemia. *Arterioscler Thromb Vasc Biol* 2000; **20**: 259–265.