



# Cataracts: Role of the unfolded protein response

Toshimichi Shinohara \*, Kengo Ikesugi, Michael L. Mulhern

*Department of Ophthalmology and Visual Sciences, University of Nebraska Medical Center,  
985840 Nebraska Medical Center, Omaha, NE 68198-5840, USA*

Received 28 July 2005; accepted 1 August 2005

**Summary** Many cataracts are caused by unfolded protein aggregates in highly oxidized lenses, but the underlying mechanisms of their formation are poorly understood. A literature search has shown that many cataractogenic stressors are also endoplasmic reticulum (ER) stressors, which induce the unfolded protein response (UPR) in a wide range of cell types. Since the lumen of the ER is highly oxidized, ER stressors might generate unfolded protein aggregates, which activate the UPR leading to the production of reactive oxygen species (ROS) in lens epithelial cells (LECs). ROS decrease the amount of free glutathione from whole lenses and elicit a more oxidized environment, where unfolded protein aggregates are formed and grown to large protein aggregate particles to scatter light. Recently, we have shown that ER stressors, homocysteine, tunicamycin,  $\text{Ca}^{2+}$  ionophore (A23187), and glucose deprivation induce the UPR in LECs. Here we hypothesize the cataractogenic stressors induce ER stress, initiate the UPR and ROS production in LECs with or without apoptosis and eventually resulted in cataracts.  
© 2005 Elsevier Ltd. All rights reserved.

## Introduction

The pivotal role of the endoplasmic reticulum (ER) in the life of cells has been known for several decades, but the part it plays in the progression of human diseases has only recently been appreciated. Although many causes of cataracts have been identified, the underlying mechanisms of initiation of cataract formation are poorly understood. It is increasingly evident that a few common pathways are shared in the responses to multiple, seemingly divergent stresses. A number of recent reports support idea that multiple intracellular stress pathways converge into a single event: the unfolded

protein response (UPR). The UPR is induced by unfolded protein aggregates in the ER (ER stress) [1–3].

The ER is a principal site for protein synthesis and is the location where the vast majority of secreted, glycosylated, modified, and lipid proteins are folded into their tertiary and quaternary structure. The lumen of the ER is a highly oxidized compartment and the concentration ratio of reduced glutathione (GSH):oxidized GSH (GSSG) is 3:1, whereas in the cytosol the ratio is 100:1 [4]. Only those proteins that are properly folded are formed into oligomers and are transferred to the Golgi complex for further modification. Otherwise, Bip/GRP78, a central sensing protein in the ER, binds to improperly folded proteins, which are subsequently degraded by ubiquitin [2,4,5]. If this response is not capable of removing unfolded

\* Corresponding author. Tel.: +1 402 559 4205; fax: +1 402 559 3869.

E-mail address: [tshinohara@unmc.edu](mailto:tshinohara@unmc.edu) (T. Shinohara).

protein aggregates, the cells are able to activate death pathways that include the production of ROS and an activation of multiple caspases. The basic mechanisms of the UPR induced by ER stress and enzymes involved in the UPR were recently elsewhere reviewed in depth [2,3]. We shall here describe multiple cataractogenic stressors inducing the UPR in the lenses.

Decades of study on cataract formation have shown that a wide variety of stressors induce cataracts [6–10]. Interestingly, all types of known cataracts are associated with a significant loss of free glutathione (GSH) [9,11,12] and the generation of a highly oxidized environment, where unfolded crystallin aggregate particles are formed and grow to larger than  $5 \times 10^7$  Da to scatter light [13].

Similar chemical interactions must be provoked by cataractogenic stressors in the highly oxidized lumen of the ER, generating unfolded protein aggregates. Recent reports show that the ER produces a significant amount of ROS, where electrons are generated from the formation of disulfide bonds, and oxygen molecules serve as the terminal electron acceptor [14–16]. This electron transfer then leads to the production of ROS and ultimately to the oxidation of GSH. In addition, deprivation of GSH induces dysfunctional mitochondria, which further generate ROS [17]. Thus the UPR may generate significant levels of ROS from the ER and mitochondria.

### The UPR induced by ER stress may be a common cellular basis for cataract development

Careful examination of the different stressors involved in cataractogenesis overwhelmingly shows that these stressors also induce ER stress and activate the UPR in various cell types.

**Oxidative imbalance:** Decades of studies have shown that oxidative insults induce cataract formation in the lens [11,18]. Rat lenses cultured in 0.5–25  $\mu\text{M}$  of  $\text{H}_2\text{O}_2$  induce neither cataract nor apoptosis [19]; by contrast, LECs exposed to 1000  $\mu\text{M}$  of  $\text{H}_2\text{O}_2$  underwent necrosis [20]. Our preliminary study showed that Bip/GRP78 was elevated and cell death was increased in human LECs treated with between 30 and 50  $\mu\text{M}$  of  $\text{H}_2\text{O}_2$  for 24 h, suggesting that a certain concentration of oxidative stress may induce the UPR. In contrast, deprivation of oxygen induces cataracts [21] and UPR dependent apoptosis [22,23]. Thus, appropriate oxygen tension is essential for cell survival and lack of oxygen induces the UPR in LECs which may be associated with cataract formation.

**Glucose imbalance:** Diabetic complications induce anterior and posterior sub-capsular cataracts [24,25]. Several papers reported that hypoglycemia induces the UPR in various cell types [26,27] presumably by inhibiting N-linked protein glycosylation in the ER. In the lens, osmotic and oxidative stresses are reported to contribute to cataract development at higher glucose levels [10,28]. Deprivation of glucose induces reversible cataracts [29,30], and an acute cataract has been observed in diabetics when the blood glucose has been brought down very rapidly [31,32]. In addition, ischemia (deprivation of glucose, oxygen, and various factors) induces the UPR [22,23,33] and sub-capsular and nuclear cataracts [34].

**$\text{Ca}^{2+}$  imbalance:** Although  $\text{Ca}^{2+}$  plays numerous functional roles, intralumenal  $\text{Ca}^{2+}$  storage is important for the generation of  $\text{Ca}^{2+}$  signals as well as for the correct folding and post-translational processing of proteins entering the ER after synthesis. Influx of  $\text{Ca}^{2+}$  into LECs treated with  $\text{Ca}^{2+}$  ionophore (A23187) induces the UPR [1,5,35] and cataract formation [36], while a hypo- or hyper- $\text{Ca}^{2+}$  environment causes LEC death and cataract [37].

**DNA damage (radiation insults)** has been produced by irradiation of the eye with most regions of the electromagnetic spectrum from  $\gamma$ -rays to microwaves [6]. Posterior subcapsular cataracts (PSC) have been produced by irradiation of the eye with  $\gamma$ -rays or X-rays [6,38–40]. Death of peripheral LECs appears to be associated with development of lens opacity. Dividing cells are more susceptible to X-rays, which may also explain why very young animals are more susceptible to X-ray cataracts than older animals. DNA damage beyond that which is repaired is known to induce cataract and cell death in animals [41,42] and the UPR [43–45].

**Metal ions:** Lead, copper, and mercury induce cataracts [46], the UPR and cell death. Methylmercury is known to bind thiol groups and be incorporated into proteins, which may generate unfolded protein aggregates [47]. Copper and lead induce depolarization of the mitochondria membrane [48] and lead induces the UPR by direct binding to Bip/GRP78 [47,49].

**Selenium** is an essential trace element, but at high concentrations it induces nuclear cataract in various animals, which display ionic imbalances,  $\text{Ca}^{2+}$  accumulation, calpain induced apoptosis, lens opacification, and LEC death [50]. In selenite injected rat lenses, apoptosis occurs in 7–8% of epithelial cells in the germinative zone [51] and the UPE is induced [52].

*Homocystinuria* is also highly associated with cataract formation [53,54]. Homocysteine interacts with proteins through a disulfide bond, and resulting modified protein aggregates are recognized in the ER and induce the UPR. A prolonged exposure to higher levels of homocysteine induces UPR dependent apoptosis [55,56].

*Tunicamycin* blocks the glycosylation of many glycoproteins, which generates unfolded proteins that are recognized by the ER and induce the UPR [5], LEC death, and sub-capsular opacity within 48–72 h in culture [1].

*Naphthalene cataracts*: Cortical cataracts have been produced in animals by feeding them large doses of naphthalene. The auto-oxidized toxic agent appears to be 1,2-naphthoquinone, which in turn can react with the thiol groups of glutathione and lens crystallins [57]. Recently, naphthalene/quinone has been reported to induce ER stress and apoptosis [58,59].

*Deficiency cataracts*: Amino acid and vitamin deficiencies lead to cataract formation [60]. A lack of tryptophane has been most often implicated in cataract [60,61] and also induces the UPR [45]. Riboflavin deficiency is the only vitamin deficiency which has produced cataract consistently in animals for several decades [62,63], and its deficiency has also been reported to induce the UPR and apoptosis in Jurkat cells [64,65]. A novel conserved flavin adenine dinucleotide (FAD) dependent enzyme (Ero1p) directly interacts with FAD to oxidize protein disulfide isomerase (PDI), which then subsequently oxidizes the folding protein directly, therefore riboflavin deficiency results in a striking defect in oxidative protein folding [66].

*Malnutrition*: Chronic diarrhea [67], which is common in some tropical countries, and anorexia nervosa [68,69] are associated with cataracts and probably generate deprivation of essential amino acids (such as tryptophane) and riboflavin, and induce dehydration, acidosis, and increased plasma urea concentration, which can induce carbamylation.

*Viral infection*: Herpes, retrovirus, and rubella induce cataracts [70–73]. Viral proteins in the ER activate the UPR [74]. Chronic hepatitis B virus infection induces oxidative stress and DNA damage and results in UPR induced apoptosis [75–77]. Retrovirus infection also induces Bip/GRP78 and the UPR [78]. Alpha virus induces the UPR and cell death [79]. Expression of viral protein such as viral E7 gene [80] or RB gene, a tumor repressor gene [81] in transgenic mice induce cataract, but we do not know whether the UPR is induced in these animals. Furthermore, prion protein is also known to induce the UPR [82].

*Nitric Oxide (NO)*: NO is speculated to induce cataract [83] and reported to induce the UPR [84] in macrophages.

*U18666A*: A cholesterol biosynthesis inhibitor induces subcapsular cataracts in animals and increases the risk of cataract in humans [85–88]. Several reports indicated that U18666A induces  $\text{Ca}^{2+}$  activated proteolysis, protein modification, lipid peroxidation, and loss of junction. It is apparent that apoptosis is induced in LECs shortly after being treated with U18666A [88], and two recent papers showed that the suppression of cholesterol apparently induces the UPR [89,90].

*Toxic cataracts*: Over-dose of ethanol induces cataracts [6]; acetaldehyde, a major metabolite of the alcohol, is able to bind to proteins and induce the UPR [91]. Methionine sulfoximine induces anterior and peripheral cataracts [92] and ER stress [93]. Dinitrophenol induces cataract in various animals and man [94] and also induces the UPR [95]. 3-Aminotriazole induces posterior cataracts [96] and the UPR [15].

To date, no information is available whether the following cataractogenic stressors induce the UPR; allopurinol, diquat, hyperglycemia, hyperbaric tank, uremia (carbamylation), mutant proteins in the ER, phenothiazines, or steroids.

In conclusion, it is not surprising that many cataractogenic stressors are also ER stressors, which induce similar chemical interactions in the highly oxidized lumens of the ER and induce unfolded protein aggregates in the lens. The unfolded protein aggregates induce the UPR, which induces ROS production with or without induction of apoptosis [1]. The UPR may play a central role in the initiation and progression of many types of cataract formation. The prevention of cataracts would entail the removal of any ER stressor in daily life.

## Acknowledgments

The authors are grateful to Drs. J. Harding, J. Horowitz, D. Giblin, and P. Zelenka, for critical reading of the manuscript and discussion prior to publication. This work was supported in part by the RPB and fund from the Department of Ophthalmology and Visual Sciences, UNMC.

## References

- [1] Ikesugi K, Yamamoto R, Mulhern ML, Shinohara T. Role of the unfolded protein response (UPR) in cataract formation; 2005 [still in Revision].

- [2] Rutkowski DT, Kaufman RJ. A trip to the ER: coping with stress. *Trends Cell Biol* 2004;14:20–8.
- [3] Schroder M, Kaufman RJ. The mammalian unfolded protein response. *Annu Rev Biochem* 2005;74:739–89.
- [4] Hwang C, Sinskey AJ, Lodish HF. Oxidized redox state of glutathione in the endoplasmic reticulum. *Science* 1992;257:1496–502.
- [5] Szegezdi E, Fitzgerald U, Samali A. Caspase-12 and ER-stress-mediated apoptosis: the story so far. *Ann N Y Acad Sci* 2003;1010:186–94.
- [6] Harding JJ, Crabbe JC. The lens: development, proteins, metabolism and cataract. In: Davson H, editor. *The Eye*. 3rd ed. New York: Academic Press; 1984. p. 207–492.
- [7] Bloemendal H. Molecular biology of the eye lens. New York: Wiley; 1981. p. 1–47.
- [8] Spector A. Oxidative stress-induced cataract: mechanism of action. *FASEB J* 1995;9:1173–82.
- [9] Lou MF. Redox regulation in the lens. *Prog Retin Eye Res* 2003;22:657–82.
- [10] Harding JJ. Post-translational modification of lens proteins in cataract. *Lens Eye Toxic Res* 1991;8:245–50.
- [11] Spector A. Review: oxidative stress and disease. *J Ocul Pharmacol Ther* 2000;16:193–201.
- [12] Andley UP, Liang JJ, Lou M. Biochemical mechanisms of age-related cataract. In: Albert DM, Jackobiec FA, editors. *Principals and practice in ophthalmology*. Philadelphia: Saunders; 2000. p. 1428–49.
- [13] Benedek GB. Cataract as a protein condensation disease: the Proctor Lecture. *Invest Ophthalmol Vis Sci* 1997;38:1911–21.
- [14] Harding HP, Zhang Y, Zeng H, et al. An integrated stress response regulates amino acid metabolism and resistance to oxidative stress. *Mol Cell* 2003;11:619–33.
- [15] Tu BP, Weissman JS. Oxidative protein folding in eukaryotes: mechanisms and consequences. *J Cell Biol* 2004;164:341–6.
- [16] Haynes CM, Titus EA, Cooper AA. Degradation of misfolded proteins prevents ER-derived oxidative stress and cell death. *Mol Cell* 2004;15:767–76.
- [17] Merad-Boudia M, Nicole A, Santiard-Baron D, Saille C, Ceballos-Picot I. Mitochondrial impairment as an early event in the process of apoptosis induced by glutathione depletion in neuronal cells: relevance to Parkinson's disease. *Biochem Pharmacol* 1998;56:645–55.
- [18] Giblin FJ. Glutathione: a vital lens antioxidant. *J Ocul Pharmacol Ther* 2000;16:121–35.
- [19] Spector A. The lens and oxidative stress. In: Sies H, editor. *Oxidative stress, oxidants and antioxidants*. London: Academic press; 1991. p. 529–58.
- [20] Ueda N, Shah SV. Endonuclease-induced DNA damage and cell death in oxidant injury to renal tubular epithelial cells. *J Clin Invest* 1992;90:2593–7.
- [21] Ziegler AL. Cataracts and anoxia in the premature infants. *Ann Paediatr* 1960;194:76–89.
- [22] Marjon PL, Bobrovnikova-Marjon EV, Abcouwer SF. Expression of the pro-angiogenic factors vascular endothelial growth factor and interleukin-8/CXCL8 by human breast carcinomas is responsive to nutrient deprivation and endoplasmic reticulum stress. *Mol Cancer* 2004;3:4–16.
- [23] Benavides A, Pastor D, Santos P, Tranque P, Calvo S. CHOP plays a pivotal role in the astrocyte death induced by oxygen and glucose deprivation. *Glia* 2005; July 6.
- [24] Kinoshita JH, Fukushi S, Kador P, Merola LO. Aldose reductase in diabetic complications of the eye. *Metabolism* 1979;28:462–9.
- [25] Bron AJ, Sparrow J, Brown NA, Harding JJ, Blakytyn R. The lens in diabetes. *Eye* 1993;7:260–75.
- [26] Doerrler WT, Lehrman MA. Regulation of the dolichol pathway in human fibroblasts by the endoplasmic reticulum unfolded protein response. *Proc Natl Acad Sci USA* 1999;96:13050–5.
- [27] Fernandez J, Bode B, Koromilas A, Diehl JA, Krukavets I, Snider MD, et al. Translation mediated by the internal ribosome entry site of the cat-1 mRNA is regulated by glucose availability in a PERK kinase-dependent manner. *J Biol Chem* 2002;277:11780–7.
- [28] Kinoshita JH. A thirty year journey in the polyol pathway. *Exp Eye Res* 1990;50:567–73.
- [29] Chylack Jr LT, Schaefer FL. Mechanism of "hypoglycemic" cataract formation in the rat lens. II. Further studies on the role of hexokinase instability. *Invest Ophthalmol Vis Sci* 1976;15:519–28.
- [30] Fagerholm P, Chylack Jr LT. Regional sensitivity to hypoglycemia within the rabbit lens epithelium. *Acta Ophthalmol (Copenh)* 1989;67:13–20.
- [31] Jackson RC. Temporary cataracts in diabetes mellitus. *Br J Ophthalmol* 1955;39:629–31.
- [32] Corral RJ. Coincidental changes in conscious level and lens translucency during treatment of diabetic ketoacidosis. *Br J Ophthalmol* 1975;59:233–5.
- [33] Hayashi T, Saito A, Okuno S, Ferrand-Drake M, Dodd RL, Chan PH. Damage to the endoplasmic reticulum and activation of apoptotic machinery by oxidative stress in ischemic neurons. *J Cereb Blood Flow Metab* 2005;25:41–53.
- [34] Pfister RR. The intraocular changes of anterior segment necrosis. *Eye* 1991;5:214–21.
- [35] Kaufman RJ. Orchestrating the unfolded protein response in health and disease. *J Clin Invest* 2002;110:1389–98.
- [36] Duncan G, Wormstone IM. Calcium cell signalling and cataract: role of the endoplasmic reticulum. *Eye* 1999;13:480–3.
- [37] Matsushima H, Mukai K, Yoshida S, Obara Y. Effects of calcium on human lens epithelial cells in vitro. *Jpn J Ophthalmol* 2004;48:97–100.
- [38] Shinohara T, Singh DP, Chylack Jr LT. Review: age-related cataract: immunity and lens epithelium-derived growth factor (LEDGF). *J Ocul Pharmacol Ther* 2000;16:181–91.
- [39] Pirie A. Biochemical changes in radiation cataract. *Trans Ophthalm Soc UK* 1956;76:461–75.
- [40] Zigman S. Ultraviolet A and cataracts: basic research and practical applications. *Int Ophthalmol Clin* 2005;45:29–40.
- [41] Treton JA, Courtois Y. Evolution of the distribution, proliferation and ultraviolet repair capacity of rat lens epithelial cells as a function of maturation and aging. *Mech Ageing Dev* 1981;5:251–67.
- [42] Treton JA, Courtois Y, Jeanny JC, Gontcharoff M. Analysis of the rat lens epithelial cell nucleus during ageing by scanning cytophotometry. *Mech Ageing Dev* 1983;22:265–77.
- [43] Wu S, Hu Y, Wang JL, Chatterjee M, Shi Y, Kaufman RJ. Ultraviolet light inhibits translation through activation of the unfolded protein response kinase PERK in the lumen of the endoplasmic reticulum. *J Biol Chem* 2002;277:18077–83.
- [44] van Laar T, Schouten T, Hoogervorst E, van Eck M, van der Eb AJ, Terleth C. The novel MMS-inducible gene Mif1/KIAA0025 is a target of the unfolded protein response pathway. *FEBS Lett* 2000;469:123–31.
- [45] Kaufman RJ, Scheuner D, Schroder M, et al. The unfolded protein response in nutrient sensing and differentiation. *Nat Rev Mol Cell Biol* 2002;3:411–21.
- [46] Garland D. Role of site-specific, metal-catalyzed oxidation in lens aging and cataract: a hypothesis. *Exp Eye Res* 1990;50:677–82.

- [47] Qian Y, Falahatpisheh MH, Zheng Y, Ramos KS, Tiffany-Castiglioni E. Induction of 78 kDa glucose-regulated protein (GRP78) expression and redox-regulated transcription factor activity by lead and mercury in C6 rat glioma cells. *Neurotox Res* 2001;3:581–9.
- [48] Qian Y, Zheng Y, Abraham L, Ramos KS, Tiffany-Castiglioni E. Differential profiles of copper-induced ROS generation in human neuroblastoma and astrocytoma cells. *Brain Res Mol Brain Res* 2005;134:323–32.
- [49] Qian Y, Zheng Y, Ramos KS, Tiffany-Castiglioni E. GRP78 compartmentalized redistribution in Pb-treated glia: role of GRP78 in lead-induced oxidative stress. *Neurotoxicology* 2005;26:267–75.
- [50] Shearer TR, Ma H, Fukiage C, Azuma M. Selenite nuclear cataract: review of the model. *Mol Vis* 1997;3:8–16.
- [51] Tamada Y, Fukiage C, Nakamura Y, Azuma M, Kim YH, Shearer TR. Evidence for apoptosis in the selenite rat model of cataract. *Biochem Biophys Res Commun* 2000;275:300–6.
- [52] Zu K, Ip C. Synergy between selenium and vitamin E in apoptosis induction is associated with activation of distinctive initiator caspases in human prostate cancer cells. *Cancer Res* 2003;63:6988–95.
- [53] Roybal CN, Yang S, Sun CW, Hurtado D, Vander Jagt DL, Townes TM, et al. Homocysteine increases the expression of vascular endothelial growth factor by a mechanism involving endoplasmic reticulum stress and transcription factor ATF4. *J Biol Chem* 2004;279:14844–52.
- [54] Sulochana KN, Amirthalakshmi S, Vasanthi SB, Tamilselvi R, Ramakrishnan S. Homocystinuria with congenital/developmental cataract. *Indian J Pediatr* 2000;67:725–8.
- [55] Zhang C, Cai Y, Adachi MT, Oshiro S, Aso T, Kaufman RJ, et al. Homocysteine induces programmed cell death in human vascular endothelial cells through activation of the unfolded protein response. *J Biol Chem* 2001;276:35867–74.
- [56] Austin RC, Lentz SR, Werstuck GH. Role of hyperhomocysteinemia in endothelial dysfunction and atherothrombotic disease. *Cell Death Differ* 2004;11:556–64.
- [57] Hoke H, Zellerhoff R. Metabolism and toxicity of diisopropyl naphthalene as compared to naphthalene and monoalkyl naphthalenes: a minireview. *Toxicology* 1998;126:1–7.
- [58] Feng W, Liu G, Xia R, Abramson JJ, Pessah IN. Site-selective modification of hyperreactive cysteines of ryanodine receptor complex by quinones. *Mol Pharmacol* 1999;55:821–31.
- [59] Gilmore WJ, Kirby GM. Endoplasmic reticulum stress due to altered cellular redox status positively regulates murine hepatic CYP2A5 expression. *J Pharmacol Exp Ther* 2004;308:600–8.
- [60] Dische Z, Elliott J, Pearson E, Merriam GR. Changes in proteins and protein synthesis; in tryptophane deficiency and radiation cataracts of rats. *Am J Ophthalmol* 1959;47:368–79.
- [61] Bunce GE, Hess JL, Fillnow GM. Investigation of low tryptophan induced cataract in weanling rats. *Exp Eye Res* 1978;26:399–405.
- [62] Pirie A. Vitamin deficiency and vision. *Br Med Bull* 1956;12:32–4.
- [63] Wynn M, Wynn A. Can improved diet contribute to the prevention of cataract? *Nutr Health* 1996;11:87–104.
- [64] Camporeale G, Zemleni J. Oxidative folding of interleukin-2 is impaired in flavin-deficient jurkat cells, causing intracellular accumulation of interleukin-2 and increased expression of stress response genes. *J Nutr* 2003;133:668–72.
- [65] Manthey KC, Chew YC, Zemleni J. Riboflavin deficiency impairs oxidative folding and secretion of apolipoprotein B-100 in HepG2 cells, triggering stress response systems. *J Nutr* 2005;135:978–82.
- [66] Tu BP, Ho-Schleyer SC, Travers KJ, Weissman JS. Biochemical basis of oxidative protein folding in the endoplasmic reticulum. *Science* 2000;290:1571–4.
- [67] West SK, Valmadrid CT. Epidemiology of risk factors for age-related cataract. *Surv Ophthalmol* 1995;39:323–34.
- [68] Miller D. A case of anorexia nervosa in a young woman with development of subcapsular cataracts. *Trans Ophthal Soc UK* 1958;78:217–22.
- [69] Archer AG. Cataract formation in anorexia nervosa. *Br Med J* 1981;282:274.
- [70] Matoba A. Ocular viral infections. *Pediatr Infect Dis* 1984;3:358–68.
- [71] Ficker LA, Kirkness CM, Rice NS, Steele AD. The changing management and improved prognosis for corneal grafting in herpes simplex keratitis. *Ophthalmology* 1989;96:1587–96.
- [72] Boniuk V. Rubella. *Int Ophthalmol Clin* 1975;15:229–41.
- [73] Raghu H, Subhan S, Jose RJ, Gangopadhyay N, Bhende J, Sharma S. Herpes simplex virus-1-associated congenital cataract. *Am J Ophthalmol* 2004;138:313–4.
- [74] Su HL, Liao CL, Lin YL. Japanese encephalitis virus infection initiates endoplasmic reticulum stress and an unfolded protein response. *J Virol* 2002;76:4162–71.
- [75] Waris G, Tardif KD, Siddiqui A. Endoplasmic reticulum (ER) stress: hepatitis C virus induces an ER-nucleus signal transduction pathway and activates NF-kappaB and STAT-3. *Biochem Pharmacol* 2002;64:1425–30.
- [76] Wang HC, Wu HC, Chen CF, Fausto N, Lei HY, Su IJ. Different types of ground glass hepatocytes in chronic hepatitis B virus infection contain specific pre-S mutants that may induce endoplasmic reticulum stress. *Am J Pathol* 2003;163:2441–9.
- [77] Hsieh YH, Su IJ, Wang HC, Chang WW, Lei HY, Lai MD, et al. Pre-S mutant surface antigens in chronic hepatitis B virus infection induce oxidative stress and DNA damage. *Carcinogenesis* 2004;25:2023–32.
- [78] Dimcheff DE, Faasse MA, McAtee FJ, Portis JL. Endoplasmic reticulum (ER) stress induced by a neurovirulent mouse retrovirus is associated with prolonged BiP binding and retention of a viral protein in the ER. *J Biol Chem* 2004;279:33782–90.
- [79] Li ML, Stollar V. Alphaviruses and apoptosis. *Int Rev Immunol* 2004;23:7–24.
- [80] Pan H, Griep AE. Altered cell cycle regulation in the lens of HPV-16 E6 or E7 transgenic mice: implications for tumor suppressor gene function in development. *Genes Dev* 1994;8:1285–99.
- [81] Morgenbesser SD, Williams BO, Jacks T, DePinho RA. p53-dependent apoptosis produced by Rb-deficiency in the developing mouse lens. *Nature* 1994;371:72–4.
- [82] Gotoh T, Oyadomari S, Mori K, Mori M. Nitric oxide-induced apoptosis in RAW 264.7 macrophages is mediated by endoplasmic reticulum stress pathway involving ATF6 and CHOP. *J Biol Chem* 2002;277:12343–50.
- [83] Hetz C, Russelakis-Carneiro M, Maundrell K, Castilla J, Soto C. Caspase-12 and endoplasmic reticulum stress mediate neurotoxicity of pathological prion protein. *EMBO J* 2003;22:5435–45.
- [84] Ornek K, Karel F, Buyukbingol Z. May nitric oxide molecule have a role in the pathogenesis of human cataract? *Exp Eye Res* 2003;76:23–7.
- [85] Kuszak JR, Khan AR, Cenedella RJ. An ultrastructural analysis of plasma membrane in the U18666A cataract. *Invest Ophthalmol Vis Sci* 1988;29:261–7.

- [86] Cenedella RJ, Fleschner CR. Selective association of crystallins with lens 'native' membrane during dynamic cataractogenesis. *Curr Eye Res* 1992;11:801–15.
- [87] Chandrasekhar G, Cenedella RJ. Calcium activated proteolysis and protein modification in the U18666A cataract. *Exp Eye Res* 1993;57:737–45.
- [88] Cenedella RJ, Jacob R, Borchman D, Tang D, Neely AR, Samadi A, et al. Direct perturbation of lens membrane structure may contribute to cataracts caused by U18666A, an oxidosqualene cyclase inhibitor. *J Lipid Res* 2004;45:1232–41.
- [89] Pedruzzi E, Guichard C, Ollivier V, et al. NAD (P)H oxidase Nox-4 mediates 7-ketocholesterol-induced endoplasmic reticulum stress and apoptosis in human aortic smooth muscle cells. *Mol Cell Biol* 2004;24:10703–17.
- [90] Lee JN, Ye J. Proteolytic activation of sterol regulatory element-binding protein induced by cellular stress through depletion of Insig-1. *J Biol Chem* 2004;279:45257–65.
- [91] Ji C, Kaplowitz N. Betaine decreases hyperhomocysteinemia, endoplasmic reticulum stress, and liver injury in alcohol-fed mice. *Gastroenterology* 2003;124:1488–99.
- [92] Bagchi K. The effects of methionine sulfoximine induced methionine deficiency on the crystalline lens of albino rats. *Indian J Med Res* 1959;47:437–47.
- [93] Mobley JA, Brueggemeier RW. Increasing the DNA damage threshold in breast cancer cells. *Toxicol Appl Pharmacol* 2002;180:219–26.
- [94] Ogino S, Yasukura K. Biochemical studies on cataract. VI. Production of cataracts in guinea pigs with dinitrophenol. *Am J Ophthalmol* 1957;43:936–46.
- [95] Kondoh M, Tsukada M, Kuronaga M, et al. Induction of hepatic metallothionein synthesis by endoplasmic reticulum stress in mice. *Toxicol Lett* 2004;148:133–9.
- [96] Bhuyan KC, Bhuyan DK. Regulation of hydrogen peroxide in eye humors. Effect of 3-amino-1H-1,2,4-triazole on catalase and glutathione peroxidase of rabbit eye. *Biochim Biophys Acta* 1977;497:641–51.

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

