

Influence of Nonesterified Fatty Acids and Lysolecithins on Thyroxine Binding to Thyroxine-Binding Globulin and Transthyretin

CHEN-FEE LIM,¹ S.L.A. MUNRO,² K.N. WYNNE,¹ D.J. TOPLISS,¹ and J.R. STOCKIGT¹

ABSTRACT

The hydrolysis of lecithin by phospholipase produces equimolar amounts of nonesterified fatty acids (NEFAs) and lysolecithin. In this study, we have evaluated the effect of lysolecithins and NEFAs on thyroid hormone binding by examining their interactions with thyroxine-binding globulin (TBG)(serum 1:10,000 dilution) and purified transthyretin (TTR). Unsaturated NEFAs (palmitoleic, oleic, linoleic, linolenic, arachidonic, eicosapentaenoic, and docosahexaenoic acid) inhibited [¹²⁵I]T₄ binding to TBG. Their affinities, relative to unlabeled T₄, ranged from 0.005 to 0.0016%, except for oleic acid with relative affinity of <0.0005%. Saturated NEFAs, lauric, myristic, palmitic, and stearic acid were inactive. After purification by high-performance liquid chromatography, 1-oleoyl and 2-oleoyl lysolecithin displaced [¹²⁵I]T₄ from TBG with an affinity of 0.0006 and 0.0005%, respectively. On a molar basis, this affinity was approximately 10-fold lower than arachidonic acid, the most potent NEFA in inhibiting T₄ binding to TBG in this assay system. Of all the NEFAs tested, only arachidonic acid inhibited [¹²⁵I]T₄ binding to TTR, with an affinity relative to unlabeled T₄ of 0.49%. 1-Oleoyl, 1-palmitoyl, and 1-stearoyl lysolecithin were without effect on TTR binding. The T₄-displacing effects of NEFAs are markedly attenuated by their extensive binding to albumin. Using purified [¹⁴C]NEFA preparations and heptane partitioning, the mean unbound percentages of linoleic, eicosapentaenoic, and docosahexaenoic acid in undiluted normal human serum were 0.00099, 0.0050, and 0.0042%, respectively (*n* = 3). In view of the very high degree of albumin binding of NEFAs, studies in diluted serum will grossly overestimate their competitor potency. The affinities of lysolecithins for the T₄ binding sites of TBG and TTR are lower than those of NEFAs and depend on the fatty acid component. Lysolecithins are unlikely to influence plasma protein binding of T₄ during critical illness.

INTRODUCTION

SEVERAL NONESTERIFIED FATTY ACIDS (NEFAs) have been shown to inhibit plasma protein binding of T₄ in healthy volunteers after prolonged exercise (1) and to inhibit hepatocyte T₄ uptake (2,3), which may impair deiodination of T₄ in patients with critical illness (2) and euthyroid obese subjects during calorie restriction (3). These studies suggest that NEFAs may be modulators of thyroid hormone delivery *in vivo*.

Lysolecithin and NEFA are produced in equimolar amounts from the hydrolysis of lecithin by phospholipase (4). Marked elevation of phospholipase A₂ has been reported in the sera of patients with a variety of critical illnesses, including acute pancreatitis, infections, multiple injuries, and rheumatoid arthritis

(5–7). It is not yet known whether lysolecithins may also directly modulate the binding of T₄ to plasma proteins.

While the effects of palmitic, stearic, oleic, linoleic, linolenic, and arachidonic acid on T₄ binding to human TBG have been reported (8,9), information on other long-chain NEFAs is lacking. In addition, no data are available on the interaction of NEFAs with T₄ binding to human transthyretin (TTR), but oleic acid, arachidonic acid, and docosahexaenoic acid have been shown to interact with rat and mouse TTR (10,11). The main effect of competitor interactions with the binding of T₄ to TBG will result in the alteration of serum levels of T₃ and T₄. Interaction with TTR binding in humans will be more subtle because the proportion of circulating T₄ bound to TTR is relatively small (about 20%), and any displaced T₄ would rapidly

¹ Ewen Downie Metabolic Unit and Monash University Department of Medicine, Alfred Hospital, Commercial Road, Melbourne, Victoria, Australia, 3181.

² School of Pharmaceutical Chemistry, Victorian College of Pharmacy (Monash University), Parkville, Victoria, Australia, 3052.

equilibrate with the unoccupied sites on TBG and albumin. However, interaction of a competitor with TTR may be important in individuals with complete TBG deficiency, in non-primate species in which TTR is the predominant T_4 -binding protein (12), and in tissues such as the choroid plexus where TTR has a postulated T_4 -transport function (13).

Heparin-induced alterations of [125 I] T_4 and drug binding have previously been reported (14–16). In these studies, the authors were unable to reproduce the alteration of binding by *in vitro* addition of concentrations of NEFAs equivalent to those generated by intravenous heparin administration (15,16). Such observations suggest that some unidentified factors in the serum could be responsible for altered binding of these ligands. Because lysolecithins are candidates for these effects, we have studied the interaction of lysolecithins with T_4 in comparison to that of NEFAs.

We have previously emphasized the importance of studying the interaction of competitors with T_4 binding in undiluted serum, in order to avoid dilution-related over- or underestimates of competitor potency (17). However, such studies preclude analysis of competitor interaction at specific T_4 binding sites on TBG and TTR. In this study, we have compared the competitive potency of numerous saturated and unsaturated NEFAs and lysolecithins for the binding of [125 I] T_4 to TBG and TTR. Using a TBG-specific system of normal human serum at very high dilution (1:10,000) and an isolated preparation of human TTR, we have identified marked differences in the hierarchy of competition by NEFAs and lysolecithins for T_4 binding to TBG and TTR.

MATERIALS AND METHODS

Materials

NEFAs and lysolecithins were purchased from Sigma (St. Louis, MO). The serum pool used for the binding study was from hospital outpatients who were not taking any drug known to influence the serum concentration or the binding of thyroid hormones or NEFAs. Concentrations of albumin, total NEFA, and total T_4 in this serum pool were 41 g/L (0.62 mM), 0.7 mM, and 95 nM, respectively.

NEFAs (highest concentration 0.1 mM) were dissolved in 5% ethanol. Lysolecithins (highest concentration 10 mM) were dissolved in 50% ethanol. After dilution with buffer, the final concentration of ethanol in the assay mixture was $\leq 0.5\%$, a concentration that did not interfere in the binding reaction.

TTR was isolated from pooled plasma from normal human subjects by a modification of the method of Dwulet and Benson (18) as previously described (19).

Purification of 1-acyl lysolecithins

Our initial findings suggested that commercial 1-acyl lysolecithin preparations were highly potent in displacing T_4 (20), but further evaluation showed a high level of phospholipase contamination. Hence, these phospholipids were purified using high-performance liquid chromatography (HPLC) according to the method of Gross and Sobel (21), using a SCX Radial-Pak cartridge (10 \times 0.8 cm; Waters Associates), a mobile phase consisting of acetonitrile:methanol:water (300:100:70, v/v/v), and a flow rate of 2 mL/min. The absorbance

of 1-acyl lysolecithin was monitored at 203 nm. One milligram of 1-acyl lysolecithin, dissolved in 20 μ L of chloroform:methanol (2:1, v/v), was injected with each run. Preliminary experiments established that the retention time of 1-oleoyl, 1-palmitoyl, and 1-stearoyl lysolecithin was approximately 12.0 min. Therefore, during routine purification, effluent between 10.0 and 14.5 min was collected. The effluent was evaporated in a rotary evaporator and the residue transferred to a screw-capped vial. The percentage yield for the purification was 70% for 1-palmitoyl lysolecithin and 100% for 1-oleoyl and 1-stearoyl lysolecithin.

Preparation of 2-oleoyl lysolecithin

2-Oleoyl lysolecithin was prepared according to the method of Subbaiah et al. (22) by incubating 10 mg of di-oleoyl lysolecithin with 10^6 units of lipase suspension (Sigma L4384, Type XI from *Rhizopus arrhizus*). The mixture was incubated at 37°C for 75 min. At the end of incubation, the mixture was extracted by the method of Nishijima et al. (23) using 2 mL methanol and 8 mL petroleum ether:ethyl ether (1:1, v/v). The phospholipid containing the 2-oleoyl lysolecithin was purified by a modified HPLC procedure described by Creer and Gross (24) using a C18 μ -Bondapak cartridge (Waters Associates) with methanol:water (85:15, v/v) as mobile phase and a flow rate of 1.0 mL/min. In a typical preparation, the yield of 2-oleoyl lysolecithins was approximately 60%.

[125 I] T_4 binding to TBG (1:10,000 serum dilution)

The interaction of various NEFAs and lysolecithins with [125 I] T_4 binding to TBG was assessed in normal human serum at 1:10,000 dilution. The concentration of added [125 I] T_4 (1500 μ Ci/ μ g; Amersham, Aylesbury, Buckinghamshire, U.K.) was approximately 2 pM, giving a total concentration of T_4 (endogenous plus tracer) of less than 20 pM. The displacement assay was performed by the addition of 0.1–100 μ M NEFAs or lysolecithins to the reaction mixture. After overnight (16–18 h) incubation at 4°C, non-protein-bound [125 I] T_4 was separated by the addition of 50% volume of ice-cold dextran-charcoal with immediate centrifugation (4°C, 20 min at 2000 g). This method has been extensively validated as a specific assessment of T_4 binding to TBG: (i) lack of detectable T_4 binding ($<1\%$) to other sites at 1:10,000 dilution of serum from males with hereditary total TBG deficiency; (ii) Scatchard analysis of T_4 binding by 1:10,000 diluted serum shows mean affinity and capacity corresponding to purified TBG; (iii) mixing of high TBG and zero TBG serum samples in various proportions shows uniform affinity with capacity proportional to the concentration of TBG; and (iv) an order of potency of iodothyronine inhibition of [125 I] T_4 binding that corresponds to known specificity of TBG (19,25–28). In the absence of binding proteins, oleic acid and arachidonic acid (0.1 mM) did not influence adsorption of [125 I] T_4 to dextran-charcoal (data not shown). Minimum detectable inhibition corresponded to an affinity 0.0005% that of unlabeled T_4 .

[125 I] T_4 binding to transthyretin

Binding of [125 I] T_4 to isolated TTR was studied using [125 I] T_4 (0.1–0.3 nM T_4 ; 40–60 μ Ci/ μ g; Amersham, Aylesbury,

Buckinghamshire, U.K.) with unlabeled T₄, NEFA, or lysolecithin and diluted TTR solution (3.6–7.2 µg/mL) in Tris-Cl buffer, pH 7.4. After overnight incubation at 4°C, free and protein-bound [¹²⁵I]T₄ were separated by adsorption of free [¹²⁵I]T₄ to methyl-cellulose charcoal suspension at 0°C, as previously validated (19). These modifications minimize "stripping" of [¹²⁵I]T₄ bound to TTR by charcoal. Minimum detectable inhibition in this system corresponded to an affinity 0.19% that of unlabeled T₄.

[¹⁴C]NEFAs binding in serum

Serum binding of [¹⁴C]-labeled linoleic acid, eicosapentaenoic acid, and docosahexaenoic acid was measured using heptane partition (29,30), modified for use with serum by adjustment of serum pH to 7.4 and by preliminary purification of [¹⁴C]-NEFAs (30). [¹⁴C]NEFA (1 to 2 × 10⁵ cpm/mL) was added to the heptane phase and incubated with diluted serum (1:16) with intermittent rotation at 37°C for 90 min. Provided that [¹⁴C]NEFAs were purified before use, this method shows a linear relationship with the unbound fraction over a 4096-fold range of serum dilution (30). Hence, the unbound fraction was corrected for dilution and the results expressed for undiluted serum.

RESULTS

[¹²⁵I]T₄ binding to TBG

Effect of NEFA. The inhibition of [¹²⁵I]T₄ binding to TBG by seven unsaturated (palmitoleic, oleic, linoleic, linolenic, arachidonic, eicosapentaenoic, and docosahexaenoic acid) and four saturated (lauric, myristic, palmitic, and stearic acid) long-chain NEFAs was examined using serum diluted 1:10,000. The unsaturated NEFAs displaced [¹²⁵I]T₄ binding from TBG with an affinity at least four orders of magnitude lower than that of T₄ itself (Fig. 1). The four saturated NEFA were inactive at final concentration up to 10 µM (data not shown).

The affinities of unsaturated NEFA, relative to T₄ are shown in Table 1. The two most potent unsaturated NEFAs in displacing [¹²⁵I]T₄ from TBG are arachidonic acid and docosahexaenoic acid, both with a relative affinity of 0.005%. There were, however, only minor differences in the relative affinities of unsaturated fatty acids for the T₄ binding site on TBG which ranged from 0.005 to 0.0016%, except for oleic acid, which has a relative affinity of <0.0005%.

In view of the very high degree of albumin binding of NEFAs (30), it was important to establish whether the estimates of relative affinity were influenced by albumin binding of added NEFA in the 1:10,000 diluted serum. Accordingly, T₄ displacement activity of linoleic acid and arachidonic acid was compared using normal serum and serum from a patient with total hereditary analbuminemia (a generous gift of Professor E. Kallee, Tuebingen, Germany). In view of the high TBG concentration (60 mg/L) found in this analbuminemic serum, the dilution was adjusted (1:30,000) to achieved comparable initial binding of T₄ by normal and analbuminemic serum. Results indicated that the relative affinities of these NEFAs for T₄ binding site of TBG were virtually identical using normal and analbuminemic sera, suggesting that the results in normal serum

(1:10,000 dilution) are little influenced by residual albumin binding of NEFAs.

Effect of lysolecithins. Figure 2 shows the displacement of [¹²⁵I]T₄ binding from TBG by unlabeled T₄, 1-oleoyl, 2-oleoyl, 1-palmitoyl, and 1-stearoyl lysolecithins in comparison to the effect of arachidonic acid, the most potent NEFA in inhibiting T₄ binding to TBG in this assay system.

Both 1-oleoyl and 2-oleoyl lysolecithin displaced [¹²⁵I]T₄ from TBG, with similar competitive potency. Relative to unlabeled T₄, the affinities of 1-oleoyl and 2-oleoyl lysolecithin were 0.0006 and 0.0005%, respectively. On a molar basis, this affinity was about one order of magnitude less potent than arachidonic acid. Both 1-stearoyl and 1-palmitoyl lysolecithin were inactive (Table 1).

[¹²⁵I]T₄ binding to transthyretin

Effect of NEFAs. We investigated the ability of four unsaturated (oleic, linoleic, linolenic, and arachidonic acid) and three saturated (myristic, palmitic, and stearic acid) long-chain NEFAs to displace [¹²⁵I]T₄ from TTR. Of the NEFAs studied, only arachidonic acid showed demonstrable interaction with [¹²⁵I]T₄ binding to isolated TTR, with an affinity relative to unlabeled T₄ of 0.49%.

Effect of 1-acyl lysolecithins. Neither 1-oleoyl, 1-palmitoyl, nor 1-stearoyl lysolecithin (0.1–10 mM) showed any detectable displacement of [¹²⁵I]T₄ bound to isolated TTR.

[¹⁴C]NEFAs binding in serum

The unbound percentages of linoleic acid, eicosapentaenoic acid, and docosahexaenoic acid in normal human serum were 0.00099 ± 0.000035, 0.0050 ± 0.00011, and 0.0042 ± 0.00023, respectively (mean ± SD, *n* = 3), unbound fractions




FIG. 1. Molar comparison of the effect of unlabeled T₄ and five unsaturated NEFA (palmitoleic acid 16:1; oleic acid 18:1; linoleic acid 18:2; linolenic acid 18:3; arachidonic acid 20:4) in inhibiting [¹²⁵I]T₄ binding to TBG (serum 1:10,000), Tris-Cl buffer, pH 7.4. Bound and free [¹²⁵I]T₄ were separated by charcoal adsorption. Each data point is the mean of 3–6 determinations in duplicate. The inhibition of [¹²⁵I]T₄ binding by eicosapentaenoic (20:5) and docosahexaenoic acid (22:6) is not shown because the curves overlapped with linoleic and arachidonic acid, respectively.

TABLE 1. RELATIVE AFFINITIES OF NEFA AND PHOSPHOLIPIDS FOR [125 I]T₄ BINDING TO TBG

Compound	Concentration at 50% B/F ^a (M)	Relative affinity ^b (%)
(T ₄)	5.0×10^{-11}	100
NEFA		
Palmitoleic acid 16:1	3.2×10^{-6}	0.0016
Oleic acid 18:1	$>10^{-5}$	<0.0005
Linoleic acid 18:2	1.6×10^{-6}	0.0032
Linolenic acid 18:3	2.0×10^{-6}	0.0025
Arachidonic acid 20:4	1.0×10^{-6}	0.0050
Eicosapentaenoic acid 20:5	1.6×10^{-6}	0.0032
Docosahexaenoic acid 20:6	1.0×10^{-6}	0.0050
Phospholipids		
1-Oleoyl lysolecithin	8.3×10^{-6}	0.0006
2-Oleoyl lysolecithin	5.0×10^{-5}	0.0005

^aConcentration obtained from Figure 1 and Figure 2.^bExpressed relative to T₄, taken as 100%; mean of 3–6 determinations in duplicate.

similar to those previously reported for myristic acid (0.0072%), palmitic acid (0.00067%), oleic acid (0.00090%), linolenic acid (0.0019%), and arachidonic acid (0.0020%) (30).

Estimates of the unbound concentration of the various NEFAs in the pooled serum (total NEFA concentration 700 μ M) using the proportions of individual NEFAs previously published (31), ranged from 1.6 nM for oleic acid to 0.12 nM for linolenic acid.

DISCUSSION

This study demonstrates large differences between various NEFAs and phospholipids in their hierarchy of T₄ displacement from TBG and TTR. While all the unsaturated NEFAs tested

interacted weakly with the binding of T₄ to TBG, there is a general lack of effect of the unsaturated NEFAs on the binding of T₄ to TTR (except arachidonic acid). With regard to T₄-TBG binding, it is important to note that even for the most potent NEFAs, arachidonic acid and docosahexaenoic acid, the potency relative to that of T₄ is only about 0.005%. This affinity is approximately 20-fold lower than furosemide, the most potent drug competitor for T₄-TBG binding so far described (19).

Of the four lysolecithins tested, only 1-oleoyl and 2-oleoyl lysolecithins inhibited the binding of [125 I]T₄ to TBG, with an affinity relative to unlabeled T₄ of about 0.0005%. On a molar basis, this affinity was about one order of magnitude less than arachidonic acid, the most potent unsaturated NEFA for T₄-TBG binding. Serum concentration of lysolecithins in normal human serum is between 0.2 and 0.3 mM (22), but this may increase during critical illness as a result of the marked elevation of the phospholipase A₂ levels (5–7). Given that the threshold concentration of unsaturated NEFAs for the displacement of [125 I]T₄ from plasma proteins in whole serum is between 1.5 and 2.0 mM (1,2), and taking into consideration the relative affinity of the lysolecithins for T₄-TBG binding, these findings suggest that lysolecithins are unlikely to influence the binding of T₄ to plasma proteins during critical illness.

The minute unbound fraction of linoleic acid, eicosapentaenoic acid, and docosahexaenoic acid in the present study, together with values for five other NEFAs previously reported (30), indicates almost complete binding of these lipids to serum proteins, principally to albumin (32). Such extensive binding limits the unbound concentrations, the physiologically important moiety that determines ligand interactions. However, despite the key role of albumin in determining the unbound concentration of NEFAs, numerous workers have attributed biological activities to various NEFAs, using systems either devoid of albumin or in diluted serum (33–38). The significance of these studies needs to be reassessed, taking heed of the unbound concentration of NEFAs at physiological albumin concentrations.

In conclusion, our study demonstrates widely varying affinities of various NEFAs and lysolecithins for TBG and TTR.




FIG. 2. Inhibition of [125 I]T₄ binding to TBG at serum dilution 1:10,000 by unlabeled T₄ and HPLC-purified 1-oleoyl (1-OLPC), 1-palmitoyl (1-PLPC), 1-stearoyl (1-SPLC), and 2-oleoyl (2-OLPC) lysolecithins, showing their activity relative to arachidonic acid (20:4), which is the most potent NEFA in inhibiting [125 I]T₄ binding to TBG. Bound and free [125 I]T₄ were separated by dextran-charcoal adsorption. Each point is the mean of two determinations in duplicate.

While unsaturated NEFA and 1-oleoyl and 2-oleoyl lysolecithins all showed interactions with TBG, only arachidonic acid interacted with T₄ binding to TTR. At normal concentrations, NEFAs are unlikely to influence the plasma binding of T₄ at physiological albumin concentrations, but marked increases in NEFAs concentrations might still influence T₄ binding *in vivo*, as, for example, during prolonged exercise (1) or when elevated NEFAs are associated with marked decreases in albumin concentrations (2,39,40). The weak interactions of lysolecithins with the binding of T₄ to TBG suggest that these compounds are unlikely to influence T₄ binding *in vivo*.

ACKNOWLEDGMENTS

This work was supported in part by Project Grants from the National Health and Medical Research Council of Australia, and by Alfred Hospital Medical Research Scholarships. We thank Ms. Hilary Kay for secretarial assistance.

REFERENCES

- Liewendahl K, Helenius T, Naveri H, Tikkanen H 1992 Fatty acid-induced increase in serum dialyzable free thyroxine after physical exercise: Implication for nonthyroidal illness. *J Clin Endocrinol Metab* 74:1361–1365.
- Lim C-F, Docter R, Visser TJ, et al. 1993 Inhibition of thyroxine transport into cultured rat hepatocytes by serum of nonuremic critically ill patients: Effects of bilirubin and nonesterified fatty acids. *J Clin Endocrinol Metab* 76:1165–1172.
- Lim C-F, Docter R, Krenning EP, et al. 1994 Transport of thyroxine into cultured hepatocytes: Effects of mild non-thyroidal illness and calorie restriction in obese subjects. *Clin Endocrinol* 40:79–85.
- White A, Handler P, Smith EL (eds) 1968 The lipids. In: *Principles of Biochemistry*, 4th ed. McGraw-Hill, New York; pp 57–86.
- Vadas P 1984 Elevated plasma phospholipase A₂ levels: Correlation with the hemodynamic and pulmonary changes in gram-negative septic shock. *J Lab Clin Med* 104:873–881.
- Pruzanski W, Vadas P, Stefanski E, Urowitz MB 1985 Phospholipase A₂ activity in sera and synovial fluids in rheumatoid arthritis and osteoarthritis. Its possible role as a proinflammatory enzyme. *J Rheumatol* 12:211–216.
- Grönroos JM, Kuttilla K, Nevalainen TJ 1994 Group II phospholipase A₂ in serum in critically ill surgical patients. *Crit Care Med* 22:956–959.
- Tabachnick M, Korcek L 1986 Effect of long-chain fatty acids on the binding of thyroxine and triiodothyronine to human thyroxine-binding globulin. *Biochim Biophys Acta* 881:292–296.
- Benavenga S, Calzi LL, Robbins J 1987 Effect of free fatty acids and nonlipid inhibitors of thyroid hormone binding in the immunoradiometric assay of thyroxine-binding globulin. *Clin Chem* 33:1752–1755.
- Vranckx R, Savu L, Maya M, Nunez EA 1990 Characterization of a major development-regulated serum thyroxine-binding globulin in the euthyroid mouse. *Biochem J* 271:373–379.
- Savu L, Vranckx R, Maya M, Nunez EA 1989 Binding activities of thyroxine binding globulin versus thyroxine binding prealbumin in rat sera: Differential modulation by thyroid hormone ligands, oleic acid and pharmacological drugs. *Biochem Biophys Res Commun* 159:919–926.
- Larsson M, Pettersson T, Carlstrom A 1985 Thyroid hormone binding in serum of 15 vertebrate species: Isolation of thyroxine-binding globulin and prealbumin analogs. *Gen Comp Endocrinol* 58:360–375.
- Dickson PW, Aldred AR, Menting JGT, Marley PD, Sawyer WH, Schreiber G 1987 Thyroxine transport in choroid plexus. *J Biol Chem* 262:13907–13915.
- Mendel CM, Frost PH, Kunitake ST, Cavalieri RR 1987 Mechanism of the heparin-induced increase in the concentration of free thyroxine in plasma. *J Clin Endocrinol Metab* 65:1259–1264.
- Hollander CS, Scott RL, Burgess JA, Rabinowitz D, Merimee TJ, Oppenheimer JH 1967 Free fatty acids: A possible regulator of free thyroid hormone levels in man. *J Clin Endocrinol Metab* 27:1219–1223.
- Nilsen OG, Storstein L, Jacobsen S 1977 Effect of heparin and fatty acids on the binding of quinidine and warfarin in plasma. *Biochem Pharmacol* 26:229–235.
- Lim C-F, Bai Y, Topliss DJ, Barlow JW, Stockigt JR 1988 Drug and fatty acid effects on serum thyroid hormone binding. *J Clin Endocrinol Metab* 67:682–688.
- Dwulet FE, Benson MD 1983 Polymorphism of human plasma thyroxine binding prealbumin. *Biochem Biophys Res Commun* 114:657–662.
- Munro SL, Lim C-F, Hall JG, et al. 1989 Drug competition for thyroxine binding to transthyretin (prealbumin): Comparison with effects on thyroxine-binding globulin. *J Clin Endocrinol Metab* 68:1141–1147.
- Lim C-F, Wynne KN, Stockigt JR 1987 Influence of lysolecithins (lysophosphatidylcholines, LPC) on T₄ binding in serum: An effect mediated by changes in free fatty acids (FFA) binding. *European Thyroid Association 16th Meeting, Lausanne, Switzerland, Annales d'endocrinologie* 48:104 (Abstract #55).
- Gross RW, Sobel BE 1980 Isocratic high-performance liquid chromatography separation of phosphoglycerides and lysophosphoglycerides. *J Chromatogr* 197:79–85.
- Subbaiah PV, Chen C-H, Bagdade JD, Albers JJ 1985 Substrate specificity of plasma lysolecithin acyl-transferase and the molecular species of lecithin formed by the reaction. *J Biol Chem* 260:5308–5314.
- Nishijima M, Akamatsu Y, Nojima S 1974 Purification and properties of a membrane-bound phospholipase A₁ from mycobacterium phlei. *J Biol Chem* 249:5658–5667.
- Creer MH, Gross RW 1985 Separation of isomeric lysophospholipids by reverse phase HPLC. *Lipids* 20:922–928.
- Ingbar SH 1963 Observations concerning the binding of thyroid hormones by human serum prealbumin. *J Clin Invest* 42:143–160.
- Robbins J, Cheng S-Y, Gershengorn MC, Glinioer D, Cahnmann HJ, Edelhoch H 1978 Thyroxine transport proteins of plasma, molecular properties and biosynthesis. *Recent Prog Horm Res* 34:477–519.
- Snyder SM, Cavalieri RR, Goldfine ID, Ingbar SH, Jorgensen EC 1976 Binding of thyroid hormones and their analogues to thyroxine-binding globulin in human serum. *J Biol Chem* 251:6489–6494.
- Mohr VS, Barlow JW, Topliss DJ, O'Dea K, Stockigt JR 1987 Evaluation of T₄ and T₃ binding kinetics in the thyroxine binding globulin abnormality of Australian aborigines. *Clin Endocrinol* 26:531–540.
- Spector AA 1975 Fatty acid binding to plasma protein. *J Lipid Res* 16:165–179.
- Lim C-F, Wynne KN, Stockigt JR 1989 Measurement of the unbound fraction of long chain nonesterified fatty acids (NEFA) in serum: Methodological considerations. *Clin Chim Acta* 186:31–38.
- Diem K, Lentner C (eds) 1971 Scientific Tables, 7th ed. Ciba-Geigy, Basel, p 604.
- Brown JR, Shockley P 1982 Serum albumin: Structure and characterization of its ligand binding sites. In: Jost PC, Griffith PH (eds) *Lipid-Protein Interactions*, Vols 1–2. John Wiley, New York, pp 26–68.
- Martin ME, Vranckx R, Benassayag C, Nunez EA 1986 Modifications of the properties of human sex steroid-binding protein by nonesterified fatty acids. *J Biol Chem* 261:2954–2959.
- Martin ME, Benassayag C, Nunez EA 1988 Selective changes in binding and immunological properties of human corticosteroid binding globulin by free fatty acids. *Endocrinology* 123:1178–1186.

35. Chopra IJ, Huang T-S, Hurd RE, Beredo A, Solomon DH 1984 A competitive ligand binding assay for measurement of thyroid hormone-binding inhibitor in serum and tissues. *J Clin Endocrinol Metab* **58**: 619–628.
36. Rudman D, Bixler TJ, Del Rio AE 1971 Effect of free fatty acids on binding of drugs by bovine serum albumin, by human serum albumin and by rabbit serum. *J Pharm Exp Ther* **176**:261–272.
37. Tamura M, Kuwano H, Kinoshita T, Inagami T 1985 Identification of linoleic and oleic acids as endogenous Na^+ , K^+ -ATPase inhibitors from acute volume-expanded hog plasma. *J Biol Chem* **260**: 9672–9677.
38. Kelly RA, O'Hara DS, Mitch WE, Smith TW 1986 Identification of Na^+ , K^+ -ATPase inhibitors in human plasma as nonesterified fatty acids and lysophospholipids. *J Biol Chem* **261**:11704–11711.
39. Mendel CM, Frost PH, Cavalieri RR 1986 Effect of free fatty acids on the concentration of free thyroxine in human serum: The role of albumin. *J Clin Endocrinol Metab* **63**:1394–1399.
40. Chopra IJ, Huang T-S, Solomon DH, Chaudhuri G, Chua Teco GN 1986 The role of thyroxine (T_4)-binding serum proteins in oleic acid-induced increase in free T_4 in nonthyroidal illnesses. *J Clin Endocrinol Metab* **63**:776–779.

Address reprint requests to:

Chen-Fee Lim, Ph.D.

Ewen Downie Metabolic Unit

Alfred Hospital

Commercial Road

Melbourne, Victoria, Australia, 3181