

On the Density of Intracellular Water

JAMES S. CLEGG

Laboratory for Quantitative Biology, Department of Biology, University of Miami, Coral Gables, Florida 33124

and

W. DROST-HANSEN

Laboratory for Water Research, University of Miami, Coral Gables, Florida 33124

ABSTRACT: The density of individual *Artemia* cysts has been determined by sedimentation velocity measurements at unit gravity. Dried cyst (< 0.02 g H₂O/g dry weight) densities, ρ_s , were obtained by successive sedimentation in two nonpenetrating organic solvents. This removes geometric terms from the equation relating density to sedimentation velocity. Hydrated cysts (≈ 1.68 g H₂O/g dry weight) were sedimented in 0.0750 M NaCl to obtain their density (ρ_c). Values of ρ_s , ρ_c , and their ratios were found to be independent of cyst volume; therefore, the weight fraction of water in hydrated cysts is very nearly the same in cysts of greatly different size. It can be concluded that measurement of the water content of large populations of these cysts accurately reflects the water content of individual cysts, a point which has been assumed in previous work on this system. If ρ_s does not change appreciably when dried cysts are fully hydrated then the density of their water, ρ_w , can be calculated to be 1.022 g/cm³ ($\pm 0.0011 \sigma$). That value is significantly higher than the density of pure water and is very close to estimates of ρ_w in skeletal muscle and amphibian oocytes obtained by others. However, the assumption that ρ_s is independent of hydration is open to serious criticism, for all these studies. Consequently, conclusions and interpretations derived from such measurements must be considered to be tentative and uncertain.

INTRODUCTION

ALTHOUGH MOST RESEARCH on the physical properties of intracellular water has involved the use of nuclear magnetic resonance (NMR) spectroscopy (for recent reviews see Hazlewood, 1979; Finch, 1979; Mathur-DeVré, 1979; Ratkovic', 1981; and papers by Burnell, et al., 1981; Seitz, et al., 1981), a significant amount of information has also been obtained from microwave dielectrics (see Schwan and Foster, 1977; Grant, et al., 1978; Pethig, 1979; Foster, et al., 1980), differential scanning calorimetry (see Aubin, et al., 1978; Brown and Sturtevant, 1980) and solute distribution methods (see Paine and Horowitz, 1980; Hempling, 1981; Ling and Negendank, 1980; Ling, 1981; Garlid, 1979; Horowitz and Pearson, 1981). In spite of this research effort, the current status of the subject is one of considerable disagreement, and very different interpretations have been applied to the data, notably from NMR. These various views and their physiological consequences have been described in recent books edited by Keith (1979), Drost-Hansen and Clegg (1979), and Franks (1982).

Only two previous studies have been carried out on the density of cellular water: Po'csik (1967) who used rat skeletal muscle, and Hansson Mild, et al. (1979) and Hansson Mild and Løvtrup (1981), who studied amphibian oocytes.

It was concluded that the average density of cellular water in these systems exceeded that of pure water by about two percent. Such observations are important because they indicate that either all the water in these cells exhibits an increase in density compared with pure water, or that an appreciable fraction must do so.

The present paper will describe density measurements on single *Artemia* cysts, which, in principle, should also provide information on the density of their cell water. The results are very similar to those obtained from muscle and oocytes and, like those studies, can be interpreted to indicate the presence of water whose average density exceeds that of the pure liquid. However, such an interpretation requires that an important assumption be made about the density of the nonaqueous components in these systems; that assumption will be discussed in detail.

MATERIALS AND METHODS

Artemia Cysts

The cysts of *Artemia*, a primitive crustacean known as the brine shrimp, consists of about 4000 cells enclosed within a complex shell. The cells are capable of virtually complete but fully reversible dehydration, and they have been widely used as a model experimental system (Persoone, et al., 1980). When fully hydrated the cysts are spheres, roughly 225 μm in diameter for the population used in this study.

The outer layer of the shell, the tertiary envelope, can be removed by treatment with 0.5% sodium hypochlorite at 0°C. (Nakanishi, et al., 1962) and all studies described here were done on such "decapsulated" cysts. The remaining cyst is essentially an inner cellular mass surrounded by a very thin protein-chitin cuticle which makes up less than 4% of the total volume, and even less of the total mass of the cyst. Therefore, the system being studied here is an individual unit of about 4000 eucaryotic cells and the results can be interpreted in terms of cellular properties.

The cysts were purchased from San Francisco Bay Brand, Newark, California (diploid, bisexual), the probable date of collection being the summer of 1978. They were washed, processed, and stored as previously described (Clegg, 1974, 1978). About 90% of the population is viable based on production of nauplius larvae in sea water for 72 hr. at 23°C. Decapsulated cysts can also be reversibly dehydrated, although the viability is often reduced to about 70%.

Hydrated Cyst Density (ρ_c)

The apparatus used to measure the density of individual cysts is shown in Figure 1. The center tube of the Liebig condenser (≈ 40 cm in length, and ≈ 1 cm inner diameter) was filled with a 0.0750 m NaCl solution. The cysts were equilibrated in that same solution at 0°C prior to use, and their water content determined by a gravimetric method (Clegg, 1974, 1978). Temperature control was maintained at $25 \pm 0.05^\circ\text{C}$ by a Haake heater circulator and a Sargent refrigeration unit. Cyst diameters were measured to ± 0.5 μm by using a Bausch & Lomb micrometer disc (13-16-15) calibrated with an American Optical Company micrometer scale (2mm, in units of 0.01 mm) at 100X magnification. Single cysts were handled under a dissecting microscope with a




Fig. 1. Diagrammatic version of the apparatus used for sedimentation velocity measurements on *Artemia* cysts. The equation at the lower right yields the density of the cyst (ρ_c) using the velocity (v , in cm/s), the radius of the cyst (r , in cm), the force of gravity (g), the density ($\rho_{\text{sol.}}$, in g/cm³) and the viscosity of the solution ($\eta_{\text{sol.}}$, in poise).

wood splinter, to which they adhere. Each cyst was transferred to the condenser inner tube and allowed to fall at least 5 cm through the solution before its sedimentation velocity (v) was measured using a cathetometer with a scale divided in units of 0.001 mm (Gaertner Scientific Corp., Chicago) and a stopwatch accurate to 0.05 s. Sedimentation velocity was shown to be constant as a function of distance over the 20 cm measured (Fig. 1). The density of individual cysts was calculated from the well known equation given in Fig. 1 which relates sedimentation velocity to sphere density. Values for the viscosity

($\eta_{\text{sol.}} = 0.00897$ poise) and density ($\rho_{\text{sol.}} = 1.001$ g/cm³) of the NaCl solution were obtained from published values (Millero, 1970; Stokes and Mills, 1965), and g is the acceleration due to gravity, 980.7 cm sec⁻²). Hence, the hydrated cyst density (ρ_c) was obtained by adding $\Delta \rho$ to the density of NaCl solution (Fig. 1).

Dried Cyst Density (ρ_s)

Following sedimentation in the NaCl solution each cyst was recovered from the bottom of the apparatus (Fig. 1) and dried in a small cup placed in an individual desiccator (scintillation vial containing "anhydrous" CaSO_4). The cysts buckle when they dry, assuming the shape of an irregular, thick-walled soup bowl. The initial dehydration was performed at 2°C , to suppress what little metabolism occurs during water loss, followed by equilibration at 25°C for 3 days. Gravimetric measurements of the water content of 100-mg quantities of these CaSO_4 -dried cysts showed that they contained less than $0.02\text{ g H}_2\text{O/g dry mass}$, referred to as "g/g" for brevity.

Each dried cyst was then sedimented at unit gravity in two organic liquids using the same apparatus as for hydrated cysts. Of the large number of liquids examined, 1-bromooctane and acetophenone proved to be the most suitable. Their densities and viscosities were measured at 25°C using 25-ml pycnometers and Ostwald viscometers, respectively: 1-bromooctane, $\rho = 1.114\text{ g/cm}^3$, $\eta = 1.329\text{ cp}$; acetophenone, $\rho = 1.024\text{ g/cm}^3$, $\eta = 1.661\text{ cp}$. These values are in reasonable agreement with published values (Timmermans, 1956; Weast, 1976). Each cyst was air dried after the first sedimentation run and equilibrated over CaSO_4 as usual. Initial experiments showed that the sequence in which the two liquids were used had negligible effect on the ratio of sedimentation velocities. The data presented here were obtained by initial sedimentation in 1-bromooctane, then in acetophenone. Evidence will be presented to show that these two liquids neither penetrated the cysts nor interacted with them in a way that would alter the velocity of their sedimentation.

The reason for using two liquids is that it allows the geometric terms ($2r^2/9$) to be eliminated from the sedimentation equation (Fig. 1) which, in that form, applies only to spheres. Thus, one obtains two sets of sedimentation values for the same cyst, and it is easy to show that when the sedimentation equation is divided by itself (for these two sets of data) that

$$\frac{V_1}{V_2} = \frac{\eta_2 (\rho_5 - \rho_1)}{\eta_1 (\rho_5 - \rho_2)} \quad (1)$$

where subscripts 1 and 2 refer to the two liquids employed. All terms are known except ρ_s , the dried-cyst density, which is the desired quantity.

RESULTS

Hydrated Cysts

Results from 47 hydrated cysts are shown in Figure 2. Several features are evident: there is no discernible dependence of cyst density on volume, as the density is independent of cyst radius; the densities are surprisingly uniform, the standard deviation being about 0.5% of the mean; the precision of the method is good. The value of 1.087 g/cm^3 for hydrated *Artemia* cysts is well within the range (1.02 to 1.12) observed for the densities of individual mammalian cells (Pretlow and Pretlow, 1974) and almost exactly the same as the density of unfertilized eggs of the amphibian *Ambystoma* at comparable water content (Hansson Mild, et al., 1979). This comparison further justifies the view that studies on decapsulated cysts can be considered to reflect properties of the cells themselves.




Fig. 2. The density of *Artemia* cysts as a function of cyst radius. Each point represents one cyst. At the top are shown the means (\pm their standard deviations) for cyst hydration (X), radius (r), and density (ρ_c).

It should be understood that the size range of the cysts examined in Figure 2 does not represent a random sampling of this cyst population. On the contrary, an effort was made to sample cysts across their entire size range. Thus, very few cysts in these populations exhibit the extremes of radius shown in Fig. 2; unpublished size distribution studies on large populations show that the mean decapsulated cyst radius is $105 \text{ } \mu\text{m}$ at this water content, with a normal distribution about this mean.

Density Measurements on the Same Cyst Hydrated and Dried

Figure 3 shows the outcome of studies in which the densities of hydrated and dried cysts were determined on the same individual. A total of 21 cysts was carried successfully through the three separate determinations required, one sedimentation velocity for the hydrated cyst on 0.075 m NaCl , and two for the dried ones in the organic liquids. The filled circles in Figure 3A and B represent a cyst whose ratio of ρ_s/ρ_c was somewhat high (1.226) and is not represented in Fig. 3C; however, this cyst was included in calculations of the means and standard deviations also provided in this Figure.

Dried-cyst densities are evidently independent of their hydrated radius




Fig. 3. Densities (g/cm^3) of the same dried (ρ_s) and hydrated (ρ_c) cyst as a function of the hydrated cyst radius. The numbers at the top refer to means \pm standard deviations for 21 cysts. The filled circle represents a cyst whose ratio, ρ_s/ρ_c , was 1.226. That point is not shown in part C, but was included in the calculations.

(Fig. 3A). Of some significance is the observation that the ratio of dried to hydrated cyst density (Fig. 3C) is also independent of cyst size (i.e., hydrated radius). This demonstrates that the ratio of water mass to dry mass is much the same in cysts of very different size. For instance, the extremes in hydrated cyst volume for these data are 3.6 to $9.4 \times 10^6 \mu\text{m}^3$, almost a threefold difference, but the ratio ρ_s/ρ_c remains within a few percent of the mean throughout this volume range. Therefore, measurements of water content using large populations (hundreds of milligrams) accurately describe the water content of individual cysts.

DISCUSSION

Because of their small mass—a dried cyst weighs roughly $3 \mu\text{g}$, depending on the population (Clegg, 1974) — it is not usually feasible to measure water contents of individual cysts unless $^3\text{H}_2\text{O}$ of high specific activity is used (Clegg, 1976). The latter study did indicate that measurements of water content on large populations of cysts reflect the hydration behavior of individual cysts, but the errors were large and some uncertainty existed concerning “isotope effects” and variation in the size of individual cysts. The density measurements in the present paper (Fig. 3) remove this uncertainty. The point is not trivial because water contents enter heavily into calculations on the density of water in these cells, which will be considered later in this discussion. Equally important, a number of other studies have been carried out on the physical properties of water in this system using NMR (Seitz, et al., 1980, 1981), sorption isotherms (Clegg, 1978), calorimetry (Clegg, 1979; Crowe, et al., 1981), microwave dielectrics (Clegg, et al., 1982), and quasi-elastic neutron scattering (Trantham, 1981). These studies require the use of large cyst populations, and previous interpretations of the data have essentially assumed uniformity and equivalency of water content in all individuals in the large sample studied. That assumption has been shown to be correct.

It is evident that results on the dried-cyst density (ρ_s) are valid only if the organic liquids do not penetrate the cyst or otherwise interfere with its sedimentation (i.e., cause shape changes). It has been shown that dried whole cysts are impermeable to a wide variety of organic liquids including acetone, ethyl ether, and xylene (Tazawa and Iwanami, 1974), absolute methanol, ethanol, 1-propanol, and 1-butanol (Smith and Siegel, 1975), and *n*-heptane (Clegg, 1981). That conclusion is based on the ability of cysts to remain in these liquids for days or even months without measurable effects on viability. The permeability barrier has been identified to be the cuticle, a thin part of the shell that remains on decapsulated cysts (Morris and Afzelius, 1967; DeChaffoy, et al., 1978). If these liquids penetrated the cells it is inconceivable that they would survive (Clegg, 1981). Further support for lack of penetration was obtained by intentionally abrading dried cysts and then observing their sedimentation behavior in the liquids. In these cases the cyst would begin to fall, but then slow down and swell, often then moving upward in the tube (Fig. 1) indicating penetration of liquid. Indeed, this was also observed in some of the cysts that had not been intentionally damaged; presumably damage had occurred during handling, processing, and drying of the hydrated cyst. In other words, it was quite obvious when liquid penetration had taken place. Microscopic observation of dried cysts in the two liquids showed that no detectable alteration in shape occurred, and that they sedimented similarly in these solvents (i.e., fell with the same morphological orientation). Therefore, we conclude that the data in Fig. 3 are reliable measures of dried-cyst density. These results will now be considered in the context of the density of water in these cells at their maximum hydration.

As mentioned, density measurements have been performed previously on skeletal muscle by Pócsik (1967) and on amphibian oocytes by Hansson Mild, et al. (1979) and Hansson Mild and Løvtrup (1981). In both cases the average density of the water in these systems was estimated to exceed that of pure water by one or two percent. If correct, that could have far-reaching cellular consequences. However, these authors employed calculations that either assumed, additivity of densities between the dry mass of the cells and the water present in them (Pócsik, 1967) or were based on indirect and uncertain values for ρ_s (Hansson Mild and Løvtrup, 1981). Thus, it is well known that in the *absence* of interactions between components of a mixture (that is, when solutes are dissolved, two solutions are mixed, etc.), the following equation relates the densities of the components to that of the mixture (Moelwyn-Hughes, 1965):

$$\rho_c = \rho_s + \left(1 - \frac{\rho_s}{\rho_w}\right) \rho_c W_f \quad (2)$$

We will use notation that applies to *Artemia* cysts: ρ_w = the density of water present, W_f = the weight fraction of total water, and the other densities apply to hydrated (c) and dried (s) cysts. Using experimental values and errors for ρ_s and ρ_c (Fig. 3) and a weight fraction of water 0.626 (= 1.672 g/g) we can calculate ρ_w to be 1.022 g/cm³ (\pm 0.011, accumulated σ). That value is significantly higher than bulk water density at 25° C (0.997 g/cm³) and almost the same as ρ_w for amphibian oocytes at comparable water content, also calculated by use of Eq. 2 (Hansson Mild and Løvtrup, 1981). However, it is clear that the assumption of additivity is open to very serious objections and that such calculations, and conclusions derived from them, are severely constrained as a result. For example, ρ_w obviously depends upon the value of ρ_s that is used in the calculation. If the density of the nonaqueous components changes when the cysts are hydrated then the value of ρ_s used in Eq. 2 will not be correct and will yield the wrong value for ρ_w . In fact, only a 5% increase in ρ_s upon hydration of the cysts will give ρ_w equal to that of pure water at 25° C (0.997 g/cm³). On the other hand, an increase in ρ_s would indicate $\rho_w > 1.022$ g/cm³. Without evidence for either increases or decreases in ρ_s , and in view of the fact that "volumes of mixing" can be positive for some solutes in water (expansion) and negative for others (contraction) and clearly of the order of 5% in both directions (Moelwyn-Hughes, 1965), no unambiguous conclusion can be drawn on the density of cell water in any system using such experimental values of ρ_s and ρ_c .

The dehydration susceptibility of *Artemia* allows us to examine the problem in another way. If ρ_c is measured as a function of water content (weight fraction) then a plot of ρ_c vs. $\rho_c (W_f)$ should be linear over the entire range only if additivity applies (Moelwyn-Hughes, 1965). In that case, Eq. 2 would be valid and the value of ρ_w thus obtained is likely to be correct. On the other hand, any nonlinearity in this relationship will reveal interactions that, for example, could result from hydration of the nonaqueous components. These studies are in progress. For the moment, however, it seems necessary to conclude that firm evidence for the value of the average density of water in any living cell has yet to be obtained.

ACKNOWLEDGMENTS

This research was supported by National Science Foundation grant PCM 79-25609 (J.S.C.). We appreciate comments on this paper by Drs. S. Pócsik, K. Hansson Mild, and S. Løvtrup, and also the review by Professor F. Franks of an

earlier version. The use of nonpenetrating organic liquids to evaluate dried-cyst density was suggested by Professor H. E. Rorschach, Jr., whose help we appreciate.

REFERENCES

- Aubin, M.; Prud'homme, R.E.; Pérolet, M.; Caillé, J-P. 1980. *Biochim. Biophys. Acta* 631, 90.
- Brown, A.D.; Sturtevant, J.M. 1980. *J. Membrane Biol.* 54, 21.
- Burnell, E.E.; Clark, M.E.; Hinke, J.A.M.; Chapman, N.R. 1981. *Biophys. J.* 33, 1.
- Clegg, J.S. 1974. *J. exp. Biol.* 61, 291.
- Clegg, J.S. 1976. *J. exp. Zool.* 198, 267.
- Clegg, J.S. 1978. *J. Cell Physiol.* 94, 123.
- Clegg, J.S. 1979. In *Cell-Associated Water*, W. Drost-Hansen and J.S. Clegg, eds. Academic Press, New York, NY, pp. 362-413.
- Clegg, J.S. 1981. *J. exp. Zool.* 215, 303.
- Clegg, J.S.; Szwarnowski, S.; McClean, V.; Sheppard, R.; Grant, E.H. 1983. *Biochem. Biophys. Acta*, in Press.
- Crowe, J.H.; Crowe, L.M.; O'Dell, S.J. 1981. *Mol Physiol.* 1, 145.
- DeChaffoy, D.; DeMaeyer-Criel, G.; Kondo, M. 1978. *Differentiation* 12, 99.
- Drost-Hansen, W.; Clegg, J.S., eds. 1979. In *Cell Associated Water*, Academic Press, New York, NY, 470 pp.
- Finch, E.D. 1979. In *The Aqueous Cytoplasm*, A.D. Keith, ed., Marcel Dekker, Inc., New York, NY, pp. 61-90.
- Foster, K.R.; Schepp, J.L.; Schwan, H.P. 1980. *Biophys. J.* 29, 271.
- Franks, F. ed. 1982. In *The Biophysics of Water*, Wiley-Interscience, New York, NY, in Press.
- Garlid, K.D. 1979. In *Cell-Associated Water*, W. Drost-Hansen and J.S. Clegg, eds., Academic Press, New York, NY, pp. 293-362.
- Grant, E.H.; Sheppard, R.J.; South, G.P. 1978. In *Dielectric Behavior of Biological Molecules in Solution*, Clarendon Press, Oxford, U.K., 237 pp.
- Hansson Mild, K.; Løvtrup, S.; Forslind, S. 1979. *J. exp. Biol.* 83, 305.
- Hansson Mild, K.; Løvtrup, S. 1981. *J. exp. Biol.* 91, 361.
- Hazlewood, C.F. 1979. In *Cell-Associated Water*, W. Drost-Hansen and J.S. Clegg, eds., Academic Press, New York, NY, pp. 165-260.
- Hempling, H.G.; Cicoria, A.D.; Dupre, A.M.; Thompson, S. 1981. *J. exp. Zool.* 215, 259.
- Horowitz, S.B.; Pearson, T.W. 1981. *Molec. Cell. Biol.* 1, 769.
- Keith, A.D., ed. In *The Aqueous Cytoplasm*, Marcel Dekker, Inc., New York, NY, 230 pp.
- Ling, G.N. 1981. In *International Cell Biology 1980-81*, H.E. Schweiger, ed., Springer-Verlag, Berlin, pp. 904-914.
- Ling, G.N.; Negendank, W. 1980. *Persp. Biol. Med.* 24, 215.
- Mathur-DeVré, R. 1979 *Progr. Biophys. Molec. Biol.* 35, 103.
- Millero, F.J. 1970. *J. Phys. Chem* 74, 356.
- Moelwyn-Hughes, E.A. 1965. In *Physical Chemistry*, 2nd ed. Pergamon Press, New York, NY, pp. 806-808.
- Morris, J.; Afzelius, B.A. 1967. *J. Ultrastr. Res.* 20, 244.
- Nakanishi, Y.H.; Iwasaki, T.; Okigaki, T.; Kato, H. 1962. *Annot. Zool. Jap.* 35, 223.
- Paine, P.L.; Horowitz, S.B. 1980. In *Cell Biology*, Vol. 4, L. Goldstein and D. Prescott, eds., Academic Press, New York, NY, pp. 299-338.
- Persoone, G.; Sorgeloos, P.; Roels, O.; Jaspers, E., eds. 1980. *The Brine Shrimp, Artemia* Vols. 1-3. Universa Press, Wetteren, Belgium, Vol. 1, 382 pp.

- Pethig, R. 1979. In *Dielectric and Electronic Properties of Biological Materials*. John Wiley and Sons, New York, NY, 376 pp.
- Pócsik, S. 1967. *Acta Biochem. Biophys. Acad. Sci. Hung.* 2, 149.
- Pretlow, T.J.; Pretlow, T. P. 1977. In *Methods of Cell Separation*, N. Catsimpoolas, ed., Plenum Press, New York, NY. pp. 171-192.
- Ratković, S. 1981. *Sci. Yugoslav.* 7, 19.
- Schwan, H.P.; Foster, K.R. 1977. *Biophys. J.* 17, 193.
- Seitz, P.K.; Hazlewood, C.F.; Clegg, J.S. 1980. In *The Brine Shrimp, Artemia*, Vol. 2, G. Persoone, P. Sorgeloos, O. Roels, and E. Jaspers, eds., Universa Press, Wetteren, Belgium, pp. 545-556.
- Seitz, P.K.; Chang, D.C.; Hazlewood, C.F.; Rorschach, H.E.; Clegg, J.S. 1981. *Arch. Biochem Biophys.* 210, 517.
- Smith, C.W.; Siegel, S.M. 1975. *J. Histochem, Cytochem.* 23, 80.
- Stokes, R.H.; Mills, R. 1965. In *The International Encyclopedia of Physical Chemistry and Chemical Physics*, Vol. 3, R.H. Stokes, ed., Pergamon Press, London, UK, p. 118.
- Tagawa, T.; Iwanami, Y. 1974. *Zool. Mag. (Japan)* 83, 267.
- Timmermans, J. 1956. In *Physico-Chemical Constants of Pure Organic Compounds*, Vol. 1. Elsevier Publishing Co., Inc., New York, NY, pp. 476-477.
- Trantham, E. C. 1981. Ph.D.dissertation, Rice University, Houston, TX 128 pp.
- Weast, R.C. ed. 1967. In *Handbook of Chemistry and Physics*, 48th ed. The Chemical Rubber Co., Cleveland, Ohio, p. C438.