



Dodge GR, et al: Preferential incorporation of glucosamine into the galactosamine moieties of chondroitin sulfates in articular cartilage explants. *Arthritis Rheum* 2001;44:1089-1095.

3. Gouze JN, Bordji K, Gulberti S, et al: Interleukin-1beta down-regulates the expression of glucuronosyltransferase I, a key enzyme priming glycosaminoglycan biosynthesis. *Arthritis Rheum* 2001;44:351-360.
4. Reginster JY, Deroisy R, Rovati LC, et al: Long-term effects of glucosamine sulphate on osteoarthritis progression: A randomised, placebo-controlled clinical trial. *Lancet* 2001;357:251-256.
5. McAlindon TE, LaValley MP, Gulin JP, Felson DT: Glucosamine and chondroitin for treatment of osteoarthritis: A systematic quality assessment and meta-analysis. *JAMA* 2000;283:1469-1475.
6. American College of Rheumatology Subcommittee on Osteoarthritis Guidelines: Recommendations for the Medical Management of Osteoarthritis of the Hip and Knee: 2000 update. *Arthritis Rheum* 2000;43:1905-1915.

*The Authors Reply:* We thank Drs. Arnold and Arnold for their interest in our article. The purpose of

our article was to provide a synopsis of the basic and clinical science data in the literature regarding the use of glucosamine and chondroitin sulfate in the management of osteoarthritis. This endeavor was motivated by reports and accompanying publicity in the popular media pertaining to the use of these drugs as potential chondroprotective agents in repairing articular cartilage and decelerating the degenerative process.

While we agree with Drs. Arnold and Arnold that glucosamine sulfate "has the potential to contribute to the relief of pain and the improvement of function with a low likelihood of side effects," we feel that many unanswered questions remain regarding their long-term effects, the most effective dosage and route of administration, and product purity. At the time of publication, the lack of substantial and conclusive evidence underscored our inability to reach a definitive conclusion regarding their role in

the treatment of arthritis and, we believe, underlies the refusal of the Arthritis Foundation to support their use. As our goal was to summarize the existing documented studies of the use of glucosamine and chondroitin sulfate, we did not venture beyond an objective survey of the data.

Since our publication, further efforts to elucidate the role of glucosamine and chondroitin sulfate have been initiated. We, as well as the rest of the orthopaedic community, eagerly await the results of clinical trials involving large sample sizes and long-term follow-up. In our opinion, the paucity of concrete data at present continues to preclude a ringing endorsement of these agents.

Andrew A. Brief, MD  
Stephen G. Maurer, MD  
Paul E. Di Cesare, MD

New York University-Hospital for  
Joint Diseases  
New York, New York