

MINIREVIEW

Pathologic Effect of Estradiol on the Hypothalamus

JAMES R. BRAWER,^{1,2} ALAIN BEAUDET,^{2,3} GINA C. DESJARDINS,⁴ and HYMAN M. SCHIPPER³

*Departments of Anatomy and Cell Biology,² Neurology and Neurosurgery,³ and Psychiatry⁴
McGill University, Faculty of Medicine, Montreal, Quebec, Canada H3A 2B2*

ABSTRACT

Estradiol provides physiological signals to the brain throughout life that are indispensable for the development and regulation of reproductive function. In addition to its multiple physiological actions, we have shown that estradiol is also selectively cytotoxic to β -endorphin neurons in the hypothalamic arcuate nucleus. The mechanism underlying this neurotoxic action appears to involve the conversion of estradiol to catechol estrogen and subsequent oxidation to *o*-semiquinone free radicals. The estradiol-induced loss of β -endorphin neurons engenders a compensatory increment in μ opioid binding in the medial preoptic area rendering this region supersensitive to residual β -endorphin or to other endogenous opioids. The consequent persistent opioid inhibition results in a cascade of neuroendocrine deficits that are ultimately expressed as a chronically attenuated plasma LH pattern to which the ovaries respond by becoming anovulatory and polycystic. This neurotoxic action of estradiol may contribute to a number of reproductive disorders in humans and in animals in which aberrant hypothalamic function is a major component.

ESTRADIOL-INDUCED HYPOTHALAMIC PATHOLOGY

Gonadal steroids in general, and estradiol (E_2) in particular, act upon the hypothalamus throughout life. Steroids contribute to the sexual differentiation, development, and maturation of the hypothalamus and provide physiological signals to the hypothalamus and pituitary that are essential for the regulation of the ovulatory cycle. These normal physiological interactions between E_2 and the hypothalamus have been extensively characterized. There is, however, another dimension to the role of estradiol that is less well appreciated. Estradiol exerts a specific pathologic effect on the hypothalamus that may contribute to a variety of female reproductive disorders with a significant hypothalamic component. We first discovered this pathologic action using a single high dose of estradiol valerate (EV) [1]. Subsequently we found that this pathology actually develops in response to physiological levels of estradiol [2, 3]. Indeed, to some extent, the hypothalamic damage occurs with time in untreated animals as a consequence of normal physiological exposure to E_2 [2, 4].

A single large injection of estradiol valerate (EV) given to an adult female normally cycling rat produces (within 4 wk) chronic anovulation, persistent vaginal cornification, and polycystic ovaries [1, 5–7]. This single EV treatment also initiates a progressive multifocal lesion throughout the hypothalamic arcuate nucleus [1, 2]. Other hypothalamic nuclei and limbic structures, such as the amygdala, are unaffected. Within lesion foci, electron microscopy demonstrates the presence of degenerating neuronal elements (often identified as dendrites) as well as reactive microglial

cells and an unusual variety of reactive astrocytes, characterized by numerous, pleomorphic dense inclusions [1]. If the ovaries are removed prior to EV injection, the lesion does not occur [2], indicating that the pathology evolves in response to uninterrupted and unopposed exposure to physiological concentrations of E_2 produced by the ovaries as a result of the initial EV insult to the neuroendocrine axis. Chronic exposure to physiologic concentrations of E_2 by means of silastic implants in ovariectomized rats also generates the arcuate pathology [3]. The lesion does not occur in response to androgens or progestins [3, 8]. Moreover, 5 α -reduced androgen impedes the pathologic action of E_2 on the arcuate nucleus [3].

Two classes of neuroendocrine defects are associated with this arcuate pathology. The first is a blocked gonadotropin surge mechanism. Since gonadotropin surges can be reinitiated by hemiovariectomy [9, 10], this deficit appears to be a reversible consequence of the unchanging steroidal milieu engendered by the noncycling, polycystic ovaries. The second aberration is an intractable ovary-independent impairment in the hypothalamic circuitry regulating tonic LH release, resulting in a chronically suppressed plasma LH pattern. This defect is reflected in retarded depletion of hypothalamic GnRH [11] and in a corresponding deficit in pituitary GnRH receptor induction following bilateral ovariectomy [12]. It also accounts for the inability of hemiovariectomy to restore normal basal plasma LH levels despite the reinitiation of cyclic LH surges. Although these surges are probably also defective, they are sufficient, at least, to produce corpora lutea [9].

Neuronal Circuitry Affected by Estradiol Toxicity

Recently it has been possible to correlate the intractable ovary-independent hypothalamic dysfunction with the EV-induced lesion in the arcuate nucleus. The arcuate nucleus

Accepted May 26, 1993.

Received March 30, 1993.

¹Correspondence: James R. Brawer, Department of Anatomy and Cell Biology, McGill University, Strathcona Anatomy and Dentistry Building, 3640 University Street, Montreal, Quebec Canada H3A 2B2. FAX: (514) 398-5047.

contains neurons that project to the medial preoptic area (MPOA), the region in which most GnRH cells are located. Many of the afferents from the arcuate nucleus have been shown to contribute to the regulation of LH by directly or indirectly influencing the GnRH neuronal system [13]. Some of these inputs, such as those that release neuropeptide Y [14, 15] or neurotensin [16, 17], are excitatory, whereas others such as β -endorphin-containing inputs [15, 18] are inhibitory. Remarkably, although these neuronal populations are densely intermingled in the arcuate nucleus, the pathologic action of estradiol appears selective for the β -endorphin neurons.

Within 8 wk of EV treatment, the hypothalamic concentration of β -endorphin and the number of β -endorphin immunoreactive neurons in the arcuate nucleus are diminished by about 60% [19]. In contrast, EV treatment has no effect on the numbers of arcuate neurons immunoreactive for neurotensin, somatostatin, or tyrosine hydroxylase. Likewise, EV treatment does not appear to affect the hypothalamic concentrations of Met-enkephalin or of neuropeptide Y [19].

Two lines of indirect evidence indicate that β -endorphin neurons are probably destroyed, as opposed to inhibited or otherwise compromised, by the EV treatment. Cell counts performed on Nissl-stained material via unbiased stereological methods reveal a total loss of neurons in the arcuate nucleus in the EV-treated rats; this loss corresponds to the EV-induced decrement in β -endorphin neurons estimated via quantitative immunocytochemistry [19]. Secondly, quantitative analysis of cell shape in the remaining β -endorphin neurons in EV-treated rats indicates numerous distorted immunoreactive perikarya and dendritic processes, suggesting that many of the surviving cells are in the process of degenerating [19].

Beta-endorphin is a powerful inhibitor of LH release. The selective loss of a significant number of β -endorphin afferents to the MPOA, therefore, should result in enhanced pituitary LH secretion. Paradoxically, the opposite is true. The EV-induced destruction of β -endorphin neurons coincides with a marked suppression of the plasma LH pattern [20, 21]. One explanation for this incongruity involves a singular response of β -endorphin target cells in the MPOA to the loss of β -endorphin afferents.

Long before estradiol was found to selectively destroy β -endorphin cells, opioid (naloxone) binding in the anterior hypothalamus of EV-treated rats [22] and of E_2 -implanted rats [23] had been shown to be significantly greater than in normal controls. Radioautographic studies using specific opioid ligands demonstrated that this increment was due to a selective increase in μ opioid binding and that it was confined to the region of the MPOA densely populated with GnRH neurons [24]. This suggested that β -endorphin target elements in the MPOA, possibly the GnRH cells themselves, exhibit a compensatory up-regulation of μ binding sites in response to the partial deafferentation. This would render

these cells supersensitive to residual β -endorphin or to other endogenous μ ligands such as Met-enkephalin. The resultant chronic opioid inhibition would then account for the suppressed plasma LH pattern characterizing the EV-treated rat [20]. The interpretation that the increment in μ binding in the MPOA is a compensatory response to deafferentation is supported by observations in monosodium glutamate-lesioned rats [25]. Treatment of neonates with monosodium glutamate results in a selective injury to the arcuate nucleus with a marked loss of β -endorphin neurons. Animals thus lesioned exhibited significantly more μ opioid binding in the MPOA than nonlesioned controls. Moreover, they showed a strong inverse correlation between the extent of the loss in hypothalamic β -endorphin and the increment in μ binding.

The identity of the opioid target structures in the MPOA that express additional μ opioid receptors in response to the loss of β -endorphin afferents is not yet known. A likely candidate would be the GnRH neurons themselves. β -Endorphin axons synapse directly on GnRH neurons [26, 27] and may account for as much as 10% of the total synaptic input of these cells [27]. There is, however, evidence indicating that opioid suppression of GnRH involves presynaptic inhibition of facilitatory alpha adrenergic inputs to GnRH neurons [28, 29]. Although this evidence is not unequivocal [30], there is a real possibility that the ascending noradrenergic system may be directly involved in events occurring in the MPOA of the EV-treated rat.

In contrast to the putative hypersensitivity to opioids hypothesized above, there is evidence from one study that ovariectomized rats exposed to chronic supraphysiologic levels of estradiol for 17 days acquire a marked insensitivity to naloxone [31]. Because this occurs only after 17 days, it may reflect an early stage at which β -endorphin neurons are just beginning to degenerate. The resultant incipient loss of β -endorphin afferents to target neurons in the MPOA may not yet initiate compensatory up-regulation of μ opioid receptors. Thus the opioid target neurons may simply be deprived of ligand. Alternatively, the naloxone insensitivity seen in this study may result from physiological alterations that are not directly related to β -endorphin cell loss.

In addition to the loss of β -endorphin cells, other changes occur within the arcuate nucleus following EV treatment. There is a significant, albeit transient, reduction in the number of axosomatic and axodendritic shaft synapses [32]. This alteration is accompanied by a decrease in the concentration of small intramembrane particles in membranes of dendritic shafts and perikarya [33]. Although the import of these modifications is not clear, they undoubtedly reflect some sort of synaptic remodeling. Whether this is secondary to the loss of β -endorphin cells or is itself caused by the estradiol exposure is unknown.

Pituitary Deficits

The low mean plasma LH concentration in the EV-treated rat is not much greater than the GnRH-independent nadir




FIG. 1. Pathologic action of E₂ on the hypothalamus. This diagram summarizes our interpretations of the data reviewed in this article and represents a hypothetical model explaining the pathologic effect of E₂ on the hypothalamo-pituitary axis.

[34]. The pituitary LH content, LH responses to GnRH, and LH replenishment following GnRH-stimulated release are severely attenuated [5, 6, 11]. Since all of these LH parameters are, in large measure, regulated by GnRH [35–38], this cluster of deficits is consistent with a long-standing aberrant GnRH signal engendered by the arcuate lesion. Moreover, these deficits persist 28 days following ovariectomy; this is consistent with a physical lesion at the hypothalamic level that ultimately affects the GnRH system. It is also not surprising that the same parameters for FSH either are minimally affected or are normal, since the regulation of LH is far more dependent on GnRH than is that of FSH [42, 43]. The accumulation of hypothalamic and subsequent pituitary defects in the EV-treated rat ultimately translates into an abnormal attenuated plasma LH pattern. The plasma LH pattern in animals in estrus is characterized by large-amplitude LH pulses interspersed among more frequent low-amplitude pulses [20]. Following EV treatment there is a progressive diminution in the number and amplitude of the large pulses such that by Day 16, they disappear altogether [21]. Conversely, there is an equivalent increase in the frequency of small-amplitude pulses, such that the total pulse frequency remains constant. In contrast, the pulsatile FSH pattern is indistinguishable from that characterizing

normal estrus. This accords with observations that pituitary FSH function in the EV-treated rat is relatively unimpaired [6, 11].

Coincident with the striking deficits in pituitary LH content, release, and replenishment is a marked diminution in LH β mRNA. LH/FSH α and FSH β mRNAs are unaffected. The expression of gonadotropin subunit mRNAs is directly, although differentially, related to the stimulatory action of GnRH [39–41]. It is not surprising, therefore, that the steady state concentration of LH β mRNA in the EV-lesioned rat is about 30% that of normally cycling controls (unpublished). Although ovariectomy in the EV-treated rats was seen to induce a significant increase in LH β mRNA, the content was only one-half of that seen in ovariectomized controls.

Two lines of evidence link the cluster of pituitary deficits to sustained opioid inhibition of the GnRH system. The first is the nature of the alterations in the endogenous episodic plasma LH pattern in the EV-treated rat. In the rat, β -endorphin acts to reduce LH secretion by primarily decreasing the amplitude of GnRH pulses [44]. It most often does not modify other parameters such as pulse frequency or duration. Thus, the EV-induced diminution in mean LH pulse amplitude with no change in pulse frequency is consistent with opioid inhibition of the GnRH system [44]. Moreover, it has been shown that specific activation of μ opioid receptors in the MPOA diminishes the amplitude of plasma LH pulses but has no effect on pulse frequency [45]. The low-amplitude, normal-frequency, LH pattern characterizing the EV-treated rat, therefore, is precisely what would be predicted to result from the observed increase in μ binding in the MPOA [24].

Another aspect of pituitary activity that is highly dependent on the pattern of exposure to GnRH is the pituitary concentration of GnRH receptors [46, 47]. In the EV-treated rat, the pituitary concentration of GnRH receptors is chronically low [12]. Although ovariectomy in normal rats results in a marked increase in GnRH receptors, it does not improve the receptor concentration in the EV-treated animals. Opiatergic blockade, however, reinitiates cyclicity, restores normal ovarian morphology, and increases the pituitary concentration of GnRH receptors in EV-treated rats [48], suggesting that the low receptor content in the EV-treated rat reflects chronic opioid suppression of the endogenous GnRH pattern.

Mechanism of Estradiol Neurotoxicity

Until recently, it has not been at all clear how and why estrogen is selectively pathogenic to the arcuate nucleus. Since estradiol target cells in the arcuate nucleus, such as those expressing tyrosine hydroxylase or Met-enkephalin, are spared by the lesion, it is unlikely that the neuronal damage is directly mediated by a conventional E₂-receptor interaction. Moreover, other hypothalamic nuclei, rich in E₂ receptors and/or afferents from E₂-receptive neurons, do not exhibit this pathologic response to E₂ [1, 2]. However,

recent studies by H. Schipper suggest the following hypothesis [49]. A unique population of periventricular astrocytes containing dense Gomori-positive inclusions are concentrated in the arcuate nucleus. The astrocytic inclusions exhibit a nonenzymatic peroxidase activity and therefore stain intensely with diaminobenzidine (DAB) [50, 51]. EV treatment greatly increases the number and size of these inclusions [1, 2]. This peroxidase activity, by catalyzing the formation of highly cytotoxic free radicals, could mediate the local pathologic effect of E_2 on the arcuate nucleus.

The hypothalamus contains the enzymes (estrogen 2/4-hydroxylase) necessary for converting estradiol to 2- or 4-hydroxyestradiol (catechol estrogen, OHE_2) [52]. OHE_2 thus generated can be transformed to reactive *o*-semiquinone radicals by a peroxidase/ H_2O_2 -catalyzed reaction [53]. Indeed, the participation of cultured peroxidase-positive astrocytes in the oxidation of catechol estrogens to their respective *o*-semiquinones has recently been demonstrated through electron spin resonance spectroscopy [54].

We suspect that a reaction of this sort is mediated by the peroxidase-positive astrocytes in the arcuate nucleus of the EV-treated rat. It has been shown that 5α -reduced androgen, a class of steroids shown to inhibit hydroxylase activity [55], blocks estradiol induction of the arcuate lesion [3]. We have also demonstrated that antioxidants such as vitamin E [56], and 21-amino steroids (unpublished results) block the EV-induced loss of hypothalamic β -endorphin and prevent the loss of cyclicity and the formation of polycystic ovaries.

Significance of the Estradiol-Induced Neuropathology

The pathologic action of estradiol on the hypothalamus and the subsequent neuroendocrine aberrations have thus far been studied largely in the EV-treated rat. There are, however, diverse anovulatory conditions in a range of species that share important features with the EV-treated model, suggesting cross-species sensitivity of the hypothalamus to the pathologic effects of estradiol. Estradiol-treated mice, for example, exhibit reproductive and pituitary deficits similar to those characterizing the EV-treated rat [57–59]. Moreover, the EV-induced anovulatory state shares features with the process of normal aging in the rodent. The lesion in the arcuate nucleus [2, 4], as well as reductions in hypothalamic β -endorphin concentrations [60], proopiomelanocortin mRNA [61, 62], and loss of β -endorphin neurons [63], have all been associated with reproductive aging in the female. In addition to these decreases in β -endorphin parameters, aging rodents show decline in LH secretion [62, 64] as well as hypothalamic supersensitivity to endogenous opioid inhibition [65].

Pathologic processes similar to those observed in the EV-treated rat may contribute to a variety of veterinary and human reproductive disorders. Irreversible anovulatory acyclicity characterized by multicystic ovaries and accompanied by hypothalamic pathology also occurs in sheep and cattle as a consequence of ingestion of forage crops containing

nonsteroidal estrogenic compounds [66–69]. In humans, excessive endogenous opioid tone has been implicated in the attenuation of plasma LH characterizing hypothalamic amenorrhea [70, 71] and anorexia nervosa [72]. Excessive endogenous opioid function has also been implicated in the fatigue and depression associated with premenstrual syndrome, symptoms thought to reflect diminished noradrenergic transmission engendered by inappropriate opioid inhibition [73]. The possibility that the pathology described above could contribute or predispose to these or similar dysfunctions is very real, as the human hypothalamus is fully capable of converting E_2 to 2- OHE_2 [74, 75]. Moreover, the unique Gomori glia that are hypothesized to mediate the pathologic action of OHE_2 in the rat have also been identified in the human hypothalamus [76].

REFERENCES

1. Brawer JR, Naftolin F, Martin J, Sonnenschein C. Effects of a single injection of estradiol valerate on the hypothalamic arcuate nucleus and on reproductive function in the female rat. *Endocrinology* 1978; 103:501–512.
2. Brawer JR, Schipper H, Naftolin F. Ovary dependent degeneration in the hypothalamic arcuate nucleus. *Endocrinology* 1980; 107:274–279.
3. Brawer JR, Schipper HM, Robaire B. Effects of long term androgen and estradiol exposure on the hypothalamus. *Endocrinology* 1983; 112:194–199.
4. Schipper HM, Brawer JR, Felicio L, Nelson J, Finch C. The role of the gonads in histological aging of the hypothalamic arcuate nucleus. *Biol Reprod* 1981; 25:413–419.
5. Hemmings R, Farookhi R, Brawer JR. Pituitary and ovarian responses to luteinizing hormone releasing hormone in a rat with polycystic ovaries. *Biol Reprod* 1983; 29:239–248.
6. Schulster A, Farookhi R, Brawer JR. The polycystic ovarian condition in estradiol valerate-treated rats: spontaneous changes in characteristic endocrine features. *Biol Reprod* 1984; 31:587–593.
7. Brawer JR, Munoz M, Farookhi R. Development of the polycystic ovarian condition (PCO) in the estradiol valerate-treated rat. *Biol Reprod* 1986; 35:647–655.
8. Schipper HM, Pottle M, Brawer JR. Effects of progestins on the estradiol-related accumulation of astrocytic granules in the hypothalamic arcuate nucleus. *Brain Res* 1990; 527:176–179.
9. Farookhi R, Hemmings R, Brawer JR. Unilateral ovariectomy restores ovulatory cyclicity in rats with a polycystic ovarian condition. *Biol Reprod* 1985; 32:530–540.
10. Convery M, McCarthy GF, Brawer JR. Remission of the polycystic ovarian condition (PCO) in the rat following hemiovariectomy. *Anat Rec* 1990; 226:328–336.
11. Simard M, Brawer JR, Farookhi R. An intractable ovary-independent impairment in hypothalamo-pituitary function in the estradiol valerate-induced polycystic ovarian condition in the rat. *Biol Reprod* 1987; 36:1229–1237.
12. Carriere P, Brawer JR, Farookhi R. Pituitary GnRH receptor content in rats with polycystic ovaries. *Biol Reprod* 1988; 38:562–567.
13. Hokfelt T, Everitt B, Meister B, Melendrar T, Schalling M, Johansson O, Lundberg JM, Hulting A, Werner S, Cuella C, Hemmings H, Ouimet C, Walaas W, Greengard P, Goldstein M. Neurons with multiple messengers with special reference to neuroendocrine systems. In: Greep RO (ed.), *Recent Progress in Hormone Research*. Orlando, FL: Academic Press; 1986: 1–70.
14. Kalra SP, Sahu A, Kalra PS, Crowley WR. Hypothalamic neuropeptide Y: a circuit in the regulation of gonadotropin secretion and feeding behavior. *Ann New York Acad Sci* 1990; 611:273–283.
15. Kalra SP, Allen LG, Sahu A, Kalra PS, Crowley W. Gonadal Steroids and Neuropeptide Y-opioid-LHRH axis: interactions and diversities. *J Steroid Biochem* 1988; 30:185–193.
16. Akema TC, Praputittaya C, Kimura F. Effects of preoptic microinjection of neurotensin on luteinizing hormone secretion in unanesthetized ovariectomized rats with or without estrogen priming. *Neuroendocrinology* 1987; 46:345–349.
17. Alexander MJ, Mahoney PD, Ferris CF, Carraway RE, Leeman SE. Evidence that neurotensin participates in the central regulation of the preovulatory of luteinizing hormone in the rat. *Endocrinology* 1989; 124:783–788.
18. Bicknell RJ. Endogenous opioid peptides and hypothalamic neuroendocrine neurons. *J Endocrinol* 1985; 107:437–446.

19. Desjardins GC, Brawer JR, Beaudet A. Estradiol is selectively neurotoxic to hypothalamic β -endorphin neurons. *Endocrinology* 1993; 132:89–93.
20. Grosser P, McCarthy GF, Farookhi R, Robaire B, Brawer JR. Plasma patterns of LH, FSH, and prolactin in rats with a polycystic ovarian condition induced by estradiol valerate. *J Endocrinol* 1987; 114:33–39.
21. McCarthy GF, Farookhi R, Brawer JR. Plasma gonadotropin patterns characterizing the development of polycystic ovaries (PCO) in the estradiol valerate (EV)-treated rat. *Can J Physiol Pharmacol* 1990; 67:28–33.
22. Wilkinson M, Bhanot R, Wilkinson DA, Brawer JR. Prolonged estrogen treatment induces changes in opiate, benzodiazepine, and beta adrenergic binding sites. *Brain Res Bull* 1983; 11:279–281.
23. Wilkinson M, Brawer JR, Wilkinson DA. Gonadal steroid-induced modification of opioid binding sites in the anterior hypothalamus of female rats. *Biol Reprod* 1985; 35:501–506.
24. Desjardins GC, Beaudet A, Brawer JR. Alteration in opioid parameters in the hypothalamus of rats with estradiol valerate-induced polycystic ovarian disease. *Endocrinology* 1990; 127:2969–2976.
25. Desjardins GC, Brawer JR, Beaudet A. Monosodium glutamate-induced reductions in hypothalamic β -endorphin content result in μ opioid receptor upregulation in medial preoptic area. *Neuroendocrinology* 1992; 56:378–384.
26. Leranth C, MacLuskey N, Shanabrough M, Naftolin F. Immunohistochemical evidence for synaptic connections between pro-opiomelanocortin-immunoreactive axons and LHRH neurons in the preoptic area of the rat. *Brain Res* 1988; 167:176.
27. Chen W, Witkin J, Silverman A. Beta-endorphin and gonadotropin-releasing hormone synaptic input to gonadotropin-releasing hormone neurosecretory cells in the male rat. *J Comp Neurol* 1989; 286:85–95.
28. Kalra SP, Simpkins JW. Evidence for noradrenergic mediation of opioid effects on luteinizing hormone secretion. *Endocrinology* 1981; 109:776–782.
29. Dyer RG, Grossman R, Mansfield S, Diez-Guerra FJ, Bicknell RJ, Hollingsworth S. Opioid peptides inhibit noradrenergic transmission in the preoptic area to block LH secretion: evidence from neonatally androgenized rats. *Brain Res Bull* 1988; AUS; volume no., please 721–727.
30. Clough RW, Hoffman GE, Sladek CD. Synergistic interaction between opioid receptor blockade and alpha-adrenergic stimulation on luteinizing hormone-releasing hormone (LHRH) secretion in vitro. *Neuroendocrinology* 1990; 51:131–138.
31. Fuentes M, Sahu A, Kalra SP. Evidence that long-term estrogen treatment disrupts opioid involvement in the induction of pituitary LH surge. *Brain Res* 1992; 583:183–188.
32. Garcia-Segura LM, Baetens D, Naftolin F. Synaptic remodeling in arcuate nucleus after injection with estradiol valerate in adult female rats. *Brain Res* 1986; 366:131–136.
33. Olmos G, Aguilera P, Tranque P, Naftolin F, Garcia-Segura LM. Estrogen-induced synaptic remodeling in adult rat brain is accompanied by the reorganization of neuronal membranes. *Brain Res* 1987; 425:57–64.
34. Carriere P, Brawer JR, Farookhi R. Alterations in GnRH-dependent gonadotropin secretion in rats with polycystic ovaries (PCO). *Biol Reprod* 1991; 45:685–690.
35. Lewis CE, Morris JF, Fink G, Johnson M. Changes in the granule population of gonadotrophs of hypogonadal (*bpg*) and normal female mice associated with the priming effect of LH-releasing hormone *in vitro*. *J Endocrinol* 1985; 109:35–44.
36. Starzec A, Counis R, Justiz M. Gonadotropin-releasing hormone stimulates synthesis of the polypeptide chains of luteinizing hormone. *Endocrinology* 1986; 119:561–565.
37. Ramey JW, Highsmith RF, Wilfinger WW, Baldwin DM. The effects of gonadotropin-releasing hormone and estradiol on luteinizing hormone biosynthesis in cultured rat anterior pituitary cells. *Endocrinology* 1987; 120:1503–1513.
38. Hamernik DJ, Nett TM. Gonadotropin-releasing hormone increases the amount of messenger ribonucleic acid for gonadotropins in ovariectomized ewes after hypothalamic-pituitary disconnection. *Endocrinology* 1988; 122:959–966.
39. Haisenlender DJ, Ortolano GA, Yasin M, Dalkin AC, Marshall JC. Regulation of gonadotropin subunit messenger ribonucleic acid expression by gonadotropin releasing hormone pulse amplitude *in vitro*. *Endocrinology* 1993; 132:1292–1296.
40. Shupnik MA. Effects of gonadotropin-releasing hormone on rat gonadotropin gene transcription *in vitro*: requirement for pulsatile administration for luteinizing hormone- β gene stimulation. *Mol Endocrinol* 1990; 4:1444–1450.
41. Gharib SD, Wierman ME, Shupnik MA, Chin W. Molecular biology of pituitary gonadotropins. *Endocr Rev* 1990; 11:177–199.
42. De Paolo LV. Differential regulation of pulsatile luteinizing hormone (LH) and follicle stimulating hormone secretion in ovariectomized rats disclosed by treatment with a LH-releasing hormone antagonist and phenobarbital. *Endocrinology* 1985; 117:1826–1833.
43. Culler MD, Negro-Vilar A. Pulsatile follicle-stimulating hormone secretion is independent of luteinizing hormone-releasing hormone (LHRH): pulsatile replacement of LHRH bioactivity in LHRH-immunoneutralized rats. *Endocrinology* 1987; 120:2011–2021.
44. Levine JE, Bauer-Dantoin AC, Besecke LM, Conaghan LA, Legan SJ, Meredith JM, Strobl FJ, Urban JH, Vogelsong KM, Wolfe AM. Neuroendocrine regulation of the luteinizing hormone-releasing hormone pulse generator in the rat. In: Bardin CW (ed.), *Recent Progress in Hormone Research*, Orlando, FL: Academic Press; 1991: 97–153.
45. Mallory DS, Gallo RV. Medial preoptic-anterior hypothalamic area involvement in the suppression of pulsatile LH release by a Mu-opioid agonist in the ovariectomized rat. *Brain Res Bull* 1990; 25:251–257.
46. Clayton RN, Popkin RM, Fraser HM. Hypothalamic regulation of pituitary gonadotropin releasing hormone receptors: effects of gonadotropin-releasing hormone immunoneutralization. *Endocrinology* 1982; 110:1116–1123.
47. Clayton RN, Catt KJ. Gonadotropin-releasing hormone receptors: characterization, physiological regulation, and relationship to reproductive function. *Endocr Rev* 1981; 2:186–207.
48. Carriere PD, Farookhi R, Brawer JR. The role of aberrant hypothalamic opiate function in generating polycystic ovaries in the rat. *Can J Physiol Pharmacol* 1989; 67:896–901.
49. Schipper HM, Lechan RM, Reichlin S. Glial peroxidase activity in the hypothalamic arcuate nucleus: effects of estradiol valerate-induced persistent estrus. *Brain Res* 1990; 507:200–207.
50. Srebro Z, Cichocki T. A system of periventricular glia in brain characterized by large peroxisome-like organelles. *Acta Histochem* 1971; 41:108–114.
51. Schipper HM, Mateescu-Cantunari A. Identification of peroxidase-positive astrocytes by combined histochemical and immunolabeling techniques in situ and in cell culture. *J Histochem Cytochem* 1991; 39:1009–1016.
52. MacLusky NJ, Barnea ER, Clark CR, Naftolin F. Catechol estrogens and estrogen receptors. In: Merriam GR, Lipsett MB (eds.), *Catechol Estrogens*. New York: Raven Press; 1983: 151–165.
53. Kalyanaraman B, Felix CC, Sealy RC. Semiquinone anion radicals of catechol(amine)s, catechol estrogens, and their metal ion complexes. *Environ Health Perspect* 1985; 64:185–198.
54. Schipper HM, Kotake Y, Janzen EG. Catechol oxidation by peroxidase-positive astrocytes in primary culture: an electron spin resonance study. *J Neurosci* 1991; 11:2170–2176.
55. Brueggemeier RW, Bannan RA. Androgens inhibit the formation of catechol estrogens by estrogen 2-hydroxylase present in rat liver microsomal preparations. *J Steroid Biochem* 1986; 25:351–354.
56. Desjardins GC, Beaudet A, Schipper H, Brawer JR. Vitamin-E protects hypothalamic beta-endorphin neurons from estradiol neurotoxicity. *Endocrinology* 1992; 131:2482–2484.
57. Mobbs CV, Flurkey K, Gee DM, Yamamoto K, Sinha YN, Finch CE. Estradiol-induced anovulatory syndrome in female C57BL/6J mice: age-like neuroendocrine, but not ovarian, impairments. *Biol Reprod* 1984; 30:556–563.
58. Kohama SG, Anderson CP, Osterburg HH, May CP, Finch CE. Oral administration of estradiol to young C57BL/6J mice induces age-like neuroendocrine dysfunctions in the regulation of estrous cycles. *Biol Reprod* 1989; 41:227–232.
59. Forman LJ, Marquis DE, Stevens R. The effect of chronic estrogen treatment on immunoreactive beta-endorphin levels in intact female rats. *Proc Soc Exp Biol Med* 1985; 179:365–372.
60. Barden N, Dupont A, Labrie F, Merand Y, Rouleau D, Vauchy H, Roissier JR. Age-dependent changes in the beta-endorphin content of discrete rat brain nuclei. *Brain Res* 1981; 208:209–212.
61. Nelson JF, Bender M, Schachter BS. Age-related changes in proopiomelanocortin messenger ribonucleic acid levels in hypothalamus and pituitary of female C57BL/6J mice. *Endocrinology* 1988; 123:340–344.
62. Wise PM, Scarbrough K, Larson GH, Lloyd JM, Weiland NG, Chiu S. Neuroendocrine influences on aging of the female reproductive system. *Front Neuroendocrinol* 1991; 12:239–244.
63. Miller MM, Joshi D, Billiar RB, Nelson J. Loss during aging of beta-endorphinergic neurons in the hypothalamus of female C57BL/6J mice. *Neurobiol Aging* 1991; 12:239–244.
64. Finch CE, Felicio LS, Mobbs CV, Nelson JF. Ovarian and steroidal influences on neuroendocrine aging processes in female rodents. *Endocr Rev* 1984; 5:467–497.
65. Field EA, Kuhn CM. Opiate antagonist treatment reinstates estrous cycles in middle-aged persistent estrous rats. *Biol Reprod* 1989; 40:714–719.
66. Leavitt WW, Wright PA. The plant estrogen, coumestrol, as an agent effecting hypophysial gonadotropic function. *J Exp Zool* 1965; 160:319–328.
67. Adams NJ. Pathological changes in the tissues of infertile ewes with clover disease. *J Comp Pathol* 1976; 86:29–35.

68. Adams NR. Morphological changes in organs of ewes grazing oestrogenic subterranean clover. *Res Vet Sci* 1977; 22:216–221.
69. Kelly RW, Adams NR, Lindsay DR. Effect of coumestans on reproduction in the ewe. *Aust J Agric Res* 1976; 27:253–259.
70. Khoury SA, Reame NE, Kelcj R, Marshall JC. Diurnal patterns of pulsatile luteinizing hormone secretion in hypothalamic amenorrhea: reproducibility and responses to opiate blockade and an alpha adrenergic agonist. *J Clin Endocrinol & Metab* 1987; 64:755–762.
71. Wildt L, Leyendecker G. Induction of ovulation by chronic administration of naltrexone in hypothalamic amenorrhea. *J Clin Endocrinol & Metab* 1987; 64:1334–1335.
72. Yen SSC. Chronic anovulation due to CNS-hypothalamic-pituitary function. In: Yen SSC, Jaffe RB (eds.), *Reproductive Endocrinology*. Philadelphia: W.B. Saunders Company; 1986: 500–545.
73. Yen SSC. The human menstrual cycle. In: Yen SSC, Jaffe RB (eds.), *Reproductive Endocrinology*. Philadelphia: W.B. Saunders Company; 1986: 200–263.
74. Fishman J, Naftolin F, Davies IJ, Ryan KJ, Petro Z. Catechol estrogen formation by the human fetal brain and pituitary. *J Clin Endocrinol & Metab* 1976; 42:177–180.
75. Ball P, Knuppen R. Formation of 2- and 4-hydroxyestrogens by brain, pituitary, and liver of the human fetus. *J Clin Endocrinol & Metab* 1978; 47:732–737.
76. Schipper H. Gomori-positive astrocytes: biological properties and implications for neurologic and neuroendocrine disorders. *Glia* 1991; 4:365–377.