

*Menstrual cycle-related exacerbation of common medical conditions such as migraine, epilepsy, asthma, irritable bowel syndrome, and diabetes, is a well-recognized phenomenon. Accurate documentation of symptoms on a menstrual calendar allows identification of women with cyclic alterations in disease activity.*

# Menstrual Cycle Effects on Common Medical Conditions

ALLISON M. CASE, MD, FRCS(C)  
Department of Obstetrics and Gynaecology  
Royal University Hospital  
University of Saskatchewan  
Saskatoon, Saskatchewan Canada

ROBERT L. REID, MD, FRCS(C)  
Department of Obstetrics and Gynaecology  
Queen's University  
Kingston General Hospital  
Kingston, Ontario Canada

## INTRODUCTION

The menstrual cycle is an event which punctuates the lives of most women, and may be associated with diverse physical, psychological, and behavioural changes. Not surprisingly, it plays a significant role in women's health and disease. Conversely, menstrual cyclicity can be easily disrupted by disease, both physical and psychological.

There are many examples of psychological and behavioural changes associated with the menstrual cycle. Medical hospitalizations and consultations, accidents, work absenteeism, criminal behaviour, and suicides are all more common during the second half of the cycle.<sup>1,2</sup> Mothers also report minor illnesses in their children more frequently during the luteal and menstrual phases.<sup>3</sup> Premenstrual syndrome (PMS) is another well-documented example of menstrual cycle-related psychological and behavioural effects.

Exacerbation of certain medical conditions at specific menstrual cycle phases is a well recognized phenomenon. Diseases most often affected are those characterized by relapsing and remitting courses, and those which are easily triggered by external factors; for example, migraine, asthma, and epilepsy. The majority of these effects occur during the luteal and menstrual phases of the cycle. Accurate documentation of symptoms on a menstrual calendar allows identification of women with cyclic alterations in disease activity.

Several theories exist to explain these menstrual cycle-related effects on existing disease processes. These include fluctuations in levels of sex steroids, cyclic alterations in the immune system, and changing perceptions of disease severity brought about by premenstrual alterations in mood, as seen in premenstrual syndrome.

## REPRINTS

Allison M. Case, M.D. FRCS(C), Department Obs/Gyn, Rm 4544, Royal University Hospital, 103 Hospital Drive, Saskatoon, Saskatchewan, CANADA S7N 0W8

**Submitted for publication:** November 19, 1999. **Accepted:** April 26, 2000.

## THE MENSTRUAL CYCLE

Normal menstrual cyclicity requires coordination of the hypothalamus, pituitary gland, and ovaries. Gonadotropin-releasing hormone is secreted in a pulsatile fashion from the hypothalamus, stimulating the release of follicle-stimulating hormone (FSH) and luteinizing hormone (LH) from the anterior pituitary.<sup>4</sup>

During the follicular phase, there is progressive growth of an ovarian follicle. Estrogen secretion increases, at first gradually, and then exponentially in the 5–6 days leading up to ovulation. The luteal phase is characterized by production of both estrogen and progesterone from the corpus luteum. The lifespan of the corpus luteum is fixed at approximately 12 days. If fertilization of the ovum does not occur, the corpus luteum involutes, levels of estrogen and progesterone fall dramatically, and menstruation occurs. Falling levels of ovarian hormones remove the negative feedback from the pituitary and hypothalamus, and a new cycle of ovarian stimulation begins.<sup>4</sup>

## GONADOTROPIN-RELEASING HORMONE (GNRH) AGONISTS

Gonadotropin-releasing hormone (GnRH) agonists, by binding to GnRH receptors in the anterior pituitary, initially cause an increase in FSH and LH secretion, the so-called "flare" effect. However, after about one week, down-regulation and desensitization of the pituitary produces a hypogonadal state, sometimes likened to a "medical oophorectomy".<sup>5</sup>

Side effects of GnRH agonist treatment are related to hypoestrogenism. They include hot flashes, headache, vaginal dryness, and sleep disturbances. The side effect of greatest concern is accelerated bone loss. Prolonged administration of GnRH agonists (i.e., > 6 months) increases the patient's risk for osteoporosis. This effect, and the hypoestrogenic symptoms, can be prevented by the addition of steroid "add-back" therapy, very similar to postmenopausal hormone replacement therapy. Patients on GnRH agonists take a small dose of estrogen daily (0.625–1.25 mg conjugated estrogens or 1–2 mg estradiol). A progestational agent (medroxyprogesterone acetate 5 mg or oral micronized progesterone 200 mg) is administered on a cyclic schedule for 12–14 days every 1–3 months to induce a withdrawal bleed and minimize the risk of endometrial hyperplasia.<sup>5,6</sup> Sometimes the addition of a progestin will create symptoms attributable to the steroid itself, or may exacerbate the pre-existing medical condition. Since endometrial hyperplasia would be very unlikely to develop with six months of unop-

posed estrogen, estrogen add-back may be used on its own during a brief (3–6 months) diagnostic trial.

In gynecology, GnRH agonists are used to temporarily suppress ovarian steroid secretion in order to treat such conditions as endometriosis, precocious puberty, uterine leiomyomas, PMS,<sup>7</sup> and for prevention of menstruation in specific clinical situations (e.g. thrombocytopenia, leukemia).<sup>6</sup> They can also be used as diagnostic tools to confirm the relationship of the menstrual cycle to conditions such as PMS and menstrual migraine.<sup>6</sup> If symptoms or cyclic exacerbations resolve with elimination of the ovarian cycle, a link to the menstrual cycle is confirmed. These patients generally have excellent symptomatic relief with continued use of GnRH agonists or after surgical oophorectomy. GnRH agonists have also been used to treat a variety of medical conditions with severe, potentially life-threatening, menstrual cycle-related exacerbations, examples of which will be discussed in this review.

## MENSTRUAL MIGRAINE

Migraine headaches are two to three times more common in women than men,<sup>8</sup> and their frequency increases considerably after menarche.<sup>9</sup> While 60% of migrainous women link attacks to menstruation,<sup>10</sup> only 7% of women have true "menstrual migraine," defined as "attacks of migraine without aura that occur regularly on day one of menstruation,  $\pm$  2 days, and at no other time," implicating the existence of an exclusive hormonal trigger for migraine in these particular women.<sup>11</sup>

Migraines are vascular headaches, associated with a vasoconstrictive phase followed by vasodilatation.<sup>12</sup> Estrogen withdrawal is likely responsible for initiating some or all of these vascular effects on intracranial vessels.<sup>9,13–15</sup> In women with menstrual migraine, the onset of headache appears to be triggered by an abrupt decline in serum estrogen levels, rather than by any absolute level. Estrogen administration can preclude the expected migraine attack until the estrogen is discontinued.<sup>16</sup>

Several clinical observations support the association between menstrual migraine and estrogen withdrawal. Many women taking the combined oral contraceptive experience migraine headaches during the pill-free or placebo week.<sup>17</sup> Migraine tends to improve during pregnancy, but may recur postpartum associated with falling estrogen levels.<sup>9,18</sup> Migraine also tends to improve following menopause when menstrual cyclicity ceases, resulting in low estrogen levels with little fluctuation.<sup>19</sup>

Effective treatment of menstrual migraine depends first on accurate diagnosis of migraine, and

establishing a link between attacks and menstruation. This is best accomplished by prospective recording of migraine attacks and menstrual periods by the patient herself for at least three months. During this time, symptomatic relief of acute migraine attacks with the usual migraine therapies (ergotamine, analgesics, antiemetics) can be offered.

Prophylactic therapy is indicated for women with menstrual migraine who do not experience adequate relief from symptomatic treatment. Hormonal prophylaxis to stabilize estrogen levels may be useful.<sup>21-24</sup> Estrogen can be administered either orally, or percutaneously as a 25 µg or 50 µg transdermal patch or 1.5 mg of percutaneous estradiol gel.<sup>25</sup> Percutaneous estrogen may be more effective as there is less variation in serum estrogen levels compared to oral preparations.<sup>26</sup> Estrogen should be started in the late luteal phase, at least 48 hours prior to the anticipated onset of migraine, and continued for four to six days. Estradiol (1 mg) taken sublingually immediately at the onset of the headache may interrupt the usual progression to migraine.<sup>27</sup>

Medications which suppress the hypothalamic-pituitary-ovarian (H-P-O) axis have been successful for treatment of women whose menstrual migraines are refractory to the usual therapies. The combined oral contraceptive (OC) can be given continuously for three to four months, followed by a withdrawal bleed and symptomatic treatment of any resulting migraine attack. A new 28-day OC which maintains continuous, albeit lower, estrogen administration, rather than placebo for the last seven days is currently being developed, and may be very useful for women experiencing menstrual migraine, or migraine on the OC during the usual pill-free or placebo week.

GnRH agonists are a useful tool for both diagnosis and treatment of menstrual migraine, but should be tried only after all other possible treatments have been exhausted. A recent report demonstrated dramatic success in treating menstrual migraine with GnRH agonists, combined with continuous low-dose estrogen replacement therapy.<sup>28</sup> Women who respond to this treatment typically can expect to experience long-term relief following surgical oophorectomy with low dose estrogen replacement therapy. Concomitant hysterectomy, while unnecessary for migraine prevention, simplifies subsequent hormone replacement therapy, allowing replacement with estrogen alone and avoiding the need for progestin, which at times triggers recrudescence of migraine. This "definitive" therapy should only be offered to the woman with recalcitrant menstrual migraine who has completed her childbearing. Indi-

vidualized counselling regarding the potential risks of a permanent menopausal state, and the consequent need for estrogen replacement therapy (ERT) to maintain long-term health is essential, as is long-term postoperative follow-up to maintain patient compliance with ERT. For the few women who experience exacerbation of migraine headaches on ERT, several strategies have been suggested including reduction of estrogen dose, continuous rather than cyclic estrogen administration, and changing the type or route of administration of estrogen.<sup>20</sup>

## CATAMENIAL EPILEPSY

Catamenial epilepsy is observed in 10%–70% of epileptic women.<sup>29,30</sup> As in menstrual migraine, the wide range in reported incidence is due to the lack of a generally accepted definition.<sup>30</sup> Strictly defined, catamenial epilepsy is epilepsy which occurs at or worsens around menstruation. While up to 70% of epileptic women claim that their seizures are exacerbated by menstruation,<sup>30</sup> true catamenial epilepsy can be objectively demonstrated in approximately 12%.<sup>30</sup> Menstrual exacerbations occur with all types of seizures.<sup>31</sup> Accurate prospective documentation of seizures in relation to menstrual periods is essential for diagnosis, and prior to specific treatment.

Catamenial epilepsy is believed to result from cyclic alterations in both ovarian hormone levels and drug metabolism. Seizure threshold is increased by progesterone and decreased by estrogen.<sup>32</sup> A decrease in the progesterone level, or in the progesterone:estrogen ratio, correlates with increased seizure activity.<sup>33</sup> Seizure frequency has been shown to increase during two specific times in the menstrual cycle. The first corresponds to the rapid decrease in progesterone just prior to menses, and the second to the elevation of estrogen prior to ovulation.<sup>32,34</sup> An increase in seizure frequency has also been demonstrated during anovulatory cycles when progesterone levels are relatively low.<sup>35</sup> Menopause or oophorectomy may lead to significant improvement in epilepsy.<sup>2</sup>

Altered metabolism of anticonvulsants at different times in the cycle is well established.<sup>36</sup> The decrease in estrogen and progesterone at menstruation is believed to stimulate the release of hepatic monooxygenase enzymes, which accelerates anticonvulsant metabolism, and increases the risk for breakthrough seizures.<sup>36</sup> Treatment of catamenial epilepsy should include measurement of serum levels of anticonvulsants during times of seizure exacerbation.<sup>33</sup> A supplemental daily dose of seizure medication at the time of exacerbation may improve seizure control.<sup>37</sup>

TABLE 1

Mechanisms and Management of Menstrual Cycle-related Effects on Common Medical Conditions

| Medical Condition   | Mechanism   | Management  |
|---|---|---|
| Menstrual Migraine<br><i>Migraines occurring only on day 1 of menstruation, <math>\pm</math> 2 days</i> | Estrogen withdrawal   | <ol style="list-style-type: none"> <li>Symptomatic <ul style="list-style-type: none"> <li>Ergotamine, analgesics, antiemetics</li> </ul> </li> <li>Prophylactic <ul style="list-style-type: none"> <li>Estrogen (percutaneous patch or gel; oral)</li> <li>Cycle suppression (continuous OC, GnRH agonist)</li> </ul> </li> </ol>       |
| Catamenial Epilepsy<br><i>Epilepsy that worsens, or occurs exclusively at menstruation</i>              | <ol style="list-style-type: none"> <li><math>\downarrow</math> progesterone = <math>\downarrow</math> seizure threshold</li> <li>Altered drug metabolism</li> </ol> | <ol style="list-style-type: none"> <li>Symptomatic <ul style="list-style-type: none"> <li>Measure anticonvulsant levels and alter dose accordingly</li> </ul> </li> <li>Prophylactic <ul style="list-style-type: none"> <li>Progesterone (p.o., i.m.)</li> <li>Continuous combined OC</li> <li>Progestin-only OC</li> </ul> </li> </ol> |
| Premenstrual Asthma<br><i>Premenstrual &amp;/or menstrual asthma exacerbations</i>                      | Progesterone-induced hyper-ventilation and alterations in airway responsiveness   | <ol style="list-style-type: none"> <li>Symptomatic <ul style="list-style-type: none"> <li>Optimize usual asthma medications</li> </ul> </li> <li>Prophylactic <ul style="list-style-type: none"> <li>GnRH agonist (if life-threatening)</li> </ul> </li> </ol>  |
| Irritable Bowel Syndrome<br><i>Premenstrual &amp;/or menstrual exacerbations</i>                        | Progesterone-induced changes in gut motility  | <ol style="list-style-type: none"> <li>Symptomatic <ul style="list-style-type: none"> <li>Antispasmodics, promotility agents, bulk-forming laxatives</li> </ul> </li> <li>Prophylactic <ul style="list-style-type: none"> <li>GnRH agonist (severe cases)</li> </ul> </li> </ol>  |
| Diabetes<br><i>Premenstrual &amp;/or menstrual alterations in glycemic control</i>                      | <ol style="list-style-type: none"> <li>Changes in carbohydrate metabolism—? mechanism</li> <li>Altered eating control/patterns</li> </ol>                           | <ol style="list-style-type: none"> <li>Symptomatic <ul style="list-style-type: none"> <li>Adjust insulin accordingly</li> </ul> </li> <li>Prophylactic <ul style="list-style-type: none"> <li>Awareness of altered control; adjust insulin accordingly</li> </ul> </li> </ol>   |
| OC, oral contraceptive; GnRH, gonadotropin-releasing hormone  |   |   |

Many investigators have shown progesterone therapy to be beneficial.<sup>33-35,38-40</sup> Medroxyprogesterone acetate given orally (10–40 mg o.d.) or intramuscularly (150 mg q 6–12 weeks) has been the most widely tested.<sup>33,35</sup> In addition to its antiepileptic properties, progesterone given in this fashion may also reduce seizure frequency by suppressing gonadotropin release which, in turn, lowers estrogen levels.<sup>34,39</sup> The effects of combined OCs on seizure frequency have been inconsistent, with seizure exacerbation occurring on the pill-free days in some reports.<sup>33</sup> Uninterrupted combined OCs or the progestin-only OC may be preferable in women

with epilepsy as they result in continuous progesterone exposure.<sup>41</sup>

The use of ovarian suppressive agents such as danazol and GnRH agonists with steroid add-back for seizure control has not been described in the literature. It may be worthwhile to consider these medications in patients refractory to the anticonvulsant and hormonal therapies previously discussed.

## ASTHMA

There are numerous reports of increased frequency and severity of asthma attacks occurring premenstrually or at menstruation in some women.<sup>42,43</sup>

Gibbs<sup>44</sup> objectively confirmed the worsening of asthmatic symptoms premenstrually by documenting significant decreases in peak expiratory flow rates (PEFR). The precise etiology may be related to changing levels of progesterone or prostaglandins. Progesterone increases steadily after ovulation, and falls abruptly in the days prior to menstruation. Its relaxant effect on smooth muscle contractility may contribute to cyclic changes in airway responsiveness.<sup>29</sup> Progesterone-stimulated hyperventilation may further influence asthma leading to symptomatic deterioration and dyspnea. Menstrual cycle-related alterations in immune mechanisms have also been suggested.<sup>45</sup>

Treatment for premenstrual asthma consists of the usual asthma medications:  $\beta$ -adrenergic agonists, anticholinergics and corticosteroids.<sup>29</sup> Intramuscular progesterone was shown to be effective in three women with severe, refractory premenstrual asthma, eliminating the decrease in peak flow rate, as well as reducing total corticosteroid requirement.<sup>46</sup> There is one report of the use of a GnRH agonist in a woman with recurrent menstrually-related status asthmaticus.<sup>47</sup>

## IRRITABLE BOWEL SYNDROME

Irritable bowel syndrome (IBS) is a common gastrointestinal (GI) disorder, diagnosed clinically by the triad of chronic or recurrent abdominal pain, altered bowel habits, and the absence of a structural or biochemical abnormality.<sup>48,49</sup> In women, symptoms tend to recur and become cyclic, with exacerbation during the postovulatory and premenstrual phases of the menstrual cycle, suggesting a hormonal influence.<sup>50,51</sup> Ovulatory women frequently report constipation in the progesterone dominant luteal phase, with loose stool or diarrhea at, or immediately preceding, the onset of menstruation.

Progesterone has well-documented effects on the GI system, including a reduction in lower esophageal sphincter tone and delayed gastric emptying.<sup>52</sup> Delayed GI transit time in women, particularly during the luteal phase, has also been demonstrated.<sup>53</sup> Abrupt progesterone withdrawal may trigger an increase in bowel activity.

Fifty percent of women with IBS report an increase in GI symptoms such as abdominal pain, diarrhea, and constipation in association with the menstrual cycle.<sup>54-56</sup> The most dramatic changes in bowel symptoms occur at the start of menstrual flow, a time when the levels of progesterone fall, and prostaglandin E<sub>2</sub> and F<sub>2 $\alpha$</sub> , powerful stimulants of colonic contractility, rise.<sup>50</sup> Patients with IBS, in whom the colon is hyperresponsive to a variety of

stimuli, may have an exaggerated colonic response to prostaglandins released during menstruation.<sup>50</sup>

Treatment of IBS is generally symptomatic, including antispasmodic and promotility agents, and bulk-forming laxatives. There have also been several reports of successful treatment of severe, menstrually-related IBS with GnRH agonists.<sup>51</sup>

## DIABETES

Menstrual cycle-related alterations in glycemic control during the luteal and premenstrual phases in some insulin-dependent diabetics have been reported.<sup>2,57,58</sup> Deterioration in glycemic control, diabetic ketoacidosis, severe insulin reactions, and hypoglycemic episodes may occur more frequently around menstruation.<sup>59</sup> In one exceptional case, insulin resistance and ketoacidosis recurred exclusively during menstruation.<sup>60</sup>

Menstrual cycle-related changes in carbohydrate metabolism have been attributed to a variety of factors, including altered insulin receptor binding and affinity,<sup>61</sup> and impaired glucose tolerance.<sup>62</sup> Loss of eating control, such as bingeing or increased intake of sweets, is described by many women in the luteal phase and premenstrually,<sup>63</sup> and is another possible cause for loss of glycemic control. Diabetic women should be counseled regarding the possibility of altered control, and the need to recognize changes in their eating patterns and adjust insulin dosages accordingly.

## MISCELLANEOUS DISORDERS

The previous discussion describes the effects of the menstrual cycle on several relatively common disorders. Several rare conditions show menstrual variation in severity. There are numerous case reports of catamenial pneumothorax, the occurrence of recurrent spontaneous pneumothorax exclusively associated with menses, possibly the result of pleural or diaphragmatic endometriosis.<sup>64,65</sup> This condition has been successfully treated with GnRH agonists.<sup>66</sup>

There is some evidence that acute appendicitis presents more frequently in the luteal phase,<sup>67,68</sup> although this could be due to the misdiagnosis of right lower quadrant pain resulting from corpus luteal cysts, leading to unnecessary appendectomy. Other disorders exacerbated by the post-ovulatory and premenstrual phases of the menstrual cycle include acne, endocrine allergy and anaphylaxis, hereditary angioedema, erythema multiforme, urticaria, aphthous ulcers, Behçet syndrome, acute intermittent porphyria, paroxysmal supraventricular tachycardia, glaucoma, and multiple sclerosis.<sup>29,69-73</sup> GnRH agonists have been used for some of these



conditions, in particular for recurrent anaphylaxis and acute intermittent porphyria, when symptoms are severe or disabling.<sup>74-76</sup> In contrast to these disorders, myasthenia gravis generally improves premenstrually.<sup>77</sup> Although one early report indicated better outcomes if breast surgery was performed in the luteal phase,<sup>78</sup> subsequent investigators have been unable to confirm this finding.<sup>79</sup>

## AN APPROACH TO MENSTRUAL CYCLE-RELATED EXACERBATIONS OF MEDICAL CONDITIONS

Exacerbation of certain medical conditions at specific times during the menstrual cycle is a well-recognized phenomenon. The importance of prospective recording of symptoms in relation to the menstrual cycle to facilitate the diagnosis and treatment of menstrual cycle-related exacerbations cannot be overestimated. Where appropriate, cyclic alteration of symptomatic therapy may be sufficient for treatment. If such measures prove ineffectual, a trial of medical ovulation suppression may be warranted. If a dramatic benefit results, consideration should be given to ongoing therapy with a GnRH agonist and steroid add-back, or to a cheaper alternative such as medroxyprogesterone acetate or danazol. If these medical approaches are impractical due to cost or side effects, and the woman's fertility aspirations have been met, consideration of hysterectomy and bilateral oophorectomy with ongoing estrogen replacement therapy is a final option.

## CONCLUSION

The recognition of recurrent menstrual cycle-related exacerbations of a medical disorder allows consideration of innovative treatments that alter or suppress ovarian hormone production. Elimination of ovarian cyclicity may provide dramatic relief for some women in whom standard therapies prove less than ideal. **CT**

## REFERENCES

1. Dalton K. The influence of menstruation on health and disease. *Proc R Soc Med*. 1964;57:18.
2. Magos A, Studd J. Effects of the menstrual cycle on medical disorders. *Br J Hosp Med*. 1985;33:68-77.
3. Tuch RH. The relationship between a mother's menstrual status and her response to illness in her child. *Psychosom Med*. 1975;37:388-394.
4. Reid RL, Van Vugt DA. Neuroendocrine events that regulate the menstrual cycle. *Contemp Obstet Gynecol*. 1987;30:147-155.
5. Speroff L, Glass RH, Kase NG: *Clinical Gynecologic Endocrinology and Infertility*, 5th ed. Baltimore: Williams and Wilkins. 1994.
6. Reid RL. Use of GnRH agonists for menstrual cycle-related disorders. *J SOGC*. 1992;14(8):59-65.
7. Mortola JF, Girtan L, Fischer U. Successful treatment of severe premenstrual syndrome by combined use of gonadotropin-releasing hormone agonist and estrogen/progesterone. *J Clin Endocrinol Metab*. 1991;72:252A-252F.
8. Linet MS, Stewart WF. Migraine headache: epidemiologic perspectives. *Epidemiol Rev*. 1984;6:107-139.
9. Digre K, Damasio H. Menstrual migraine: differential diagnosis, evaluation, and treatment. *Clin Obstet Gynecol*. 1987;30:417-430.
10. Marcus DA. Focus on primary diagnosis and management of headache in women. *Obstet Gynecol Surv*. 1999;54:395-402.
11. MacGregor EA, Chia H, Vohrah RC, Wilkinson M. Migraine and menstruation: a pilot study. *Cephalalgia*. 1990;10:305-310.
12. Dalessio DJ. In: *Wolff's Headache and Other Head Pain*. New York: Oxford University Press. 1980.
13. Salvatore CA. Arteriole supply in normal and hyperplastic endometrium. *Am J Obstet Gynecol*. 1968;102:493-500.
14. Grant ECG. Relation of arterioles in the endometrium to headache from oral contraceptives. *Lancet*. 1965;1:1143-1144.
15. Grant ECG. Relation between headache from oral contraceptives and development of endometrial arterioles. *Br Med J*. 1968;2:402-405.
16. Somerville BW. The role of estradiol withdrawal in the etiology of menstrual migraine. *Neurology*. 1972;22:355-365.
17. Whitty CWM, Hockaday JM, Whitty MM. The effect of oral contraceptive on migraine. *Lancet*. 1966;1:856-859.
18. Stein GS. Headaches in the first post partum week and their relationship to migraine. *Headache*. 1981;21:201-205.
19. Neri I, Granella F, Nappi R, Manzoni GC, Facchinetti F, Genazaani AR. Characteristics of headache at menopause: a clinico-epidemiologic study. *Maturitas*. 1993;17:31-37.
20. Silberman SD, Merriam GR. Estrogens, progestins, and headache. *Neurology*. 1991;41:786-793.
21. DeLignières B, Vincens M, Mauvais-Jarvis P, Mas JL, Touboul PJ, Bousser MG. Prevention of menstrual migraine by percutaneous oestradiol. *Br Med J*. 1986;293:1540.
22. Magos AL, Zikha KJ, Studd JWW. Treatment of menstrual migraine by oestradiol implants. *J Neurol Neurosurg Psychiatry*. 1983;46:1044-1046.
23. Somerville BW. Estrogen-withdrawal migraine: I. Duration of exposure required and attempted prophylaxis by premenstrual estrogen administration. *Neurology*. 1975;25:239-244.
24. Somerville BW. Estrogen-withdrawal migraine. II. Attempted prophylaxis by continuous estradiol administration. *Neurology*. 1975;25:245-250.
25. Pfaffenrath V. Efficacy and safety of percutaneous estradiol vs placebo in menstrual migraine. *Cephalalgia*. 1993;13 [Suppl 13]:244.
26. MacGregor EA. Menstruation, sex hormones, and migraine. *Neurol Clin*. 1997;15:125-141.
27. Sarrell PM. Blood flow and ovarian secretions. In: Naftolin F, DeCherney A, Gutmann JN, Sarrell PM, eds. *Ovarian Secretions and Cardiovascular and Neurological Function*. Vol 80. New York, NY: Raven Press; 1990:81-89.
28. Murray SC, Muse KN. Effective treatment of severe menstrual migraine headaches with gonadotropin-releasing hormone agonist and "add-back" therapy. *Fertil Steril*. 1997;67:390-393.
29. Boggess KA, Williamson HO, Homm RJ. Influence of the menstrual cycle on systemic disease. *Obstet Gynecol Clin North Am*. 1990;17:321-342.
30. Duncan S, Read CL, Brodie MJ. How common is catamenial epilepsy? *Epilepsia*. 1993;34:827-831.
31. Newmark ME, Penry JK. Catamenial epilepsy: a review. *Epilepsia*. 1980;21:281-300.
32. Bäckström T. Epileptic seizures in women related to plasma estrogen and progesterone during the menstrual cycle. *Acta Neurol Scand*. 1976;54:321-347.

33. Zimmerman AW. Hormones and epilepsy. *Neurol Clin.* 1986;4:853–861.
34. Bäckström T, Zetterlund B, Blom S, Romano M. Effects of intravenous progesterone infusions on the epileptic discharge frequency in women with partial epilepsy. *Acta Neurol Scand.* 1984;69:240–248.
35. Mattson RH, Kamer JA, Cramer JA, Caldwell BV. Seizure frequency and the menstrual cycle: a clinical study. *Epilepsia.* 1981;22:242.
36. Kumar N, Behari M, Aruja GK, Jailkhani BL. Phenytoin levels in catamenial epilepsy. *Epilepsia.* 1988;29:155–158.
37. Liporace JD. Women's issues in epilepsy. Menses, child-bearing and more. *Postgrad Med.* 1997;102:123–124, 127–129, 133–135.
38. Dana-Haeri J, Richens A. Effect of norethisterone on seizures associated with menstruation. *Epilepsia.* 1983;24:377–381.
39. Hall SM. Treatment of menstrual epilepsy with a progesterone-only oral contraceptive. *Epilepsia.* 1977;18:235–236.
40. Herzog AG. Intermittent progesterone therapy and frequency of complex partial seizures in women with menstrual disorders. *Neurology.* 1986;36:1607–1610.
41. Holmes GL. Effects of menstruation and pregnancy on epilepsy. *Semin Neurol.* 1988;8:234–239.
42. Settupane RA, Simon RA. Menstrual cycle and asthma. *Ann Allergy.* 1989;63:373–378.
43. Hanley SP. Asthma variation with menstruation. *Br J Dis Chest.* 1981;75:306–308.
44. Gibbs CJ, Counts II, Lock R, Finnegan OC, White RJ. Premenstrual exacerbations of asthma. *Thorax.* 1984;39:833–836.
45. Ozkaragoz K, Cakin F. The effect of menstruation on immediate skin reactions in patients with respiratory allergy. *J Asthma Res.* 1970;7:171–175.
46. Beynon HLC, Garbett ND, Barnes PJ. Severe premenstrual exacerbations of asthma: effect of intramuscular progesterone. *Lancet.* 1988;2:370–372.
47. Blumenfeld Z, Bentur L, Yoffe N, Alroy G, Rubin AE. Menstrual asthma: use of a gonadotropin-releasing hormone analogue for the treatment of cyclic aggravation of bronchial asthma. *Fertil Steril.* 1994;62:197–200.
48. Sandler RS. Epidemiology of irritable bowel syndrome in the United States. *Gastroenterology.* 1990;99:409–412.
49. Schuster M. Diagnostic evaluation of the irritable bowel syndrome. *Gastroenterol Clin North Am.* 1991;20:269–278.
50. Whitehead WE, Cheskin LJ, Heller BR, et al. Evidence for exacerbation of irritable bowel syndrome during menses. *Gastroenterology.* 1990;98:1485–1489.
51. Mathias JR, Clench MH, Reeves-Darby VG, et al. Effect of leuprolide acetate in patients with moderate to severe functional bowel disease. Double-blind, placebo controlled study. *Dig Dis Sci.* 1994;39:1155–1162.
52. Fisher RS, Roberts GS, Grabowski CJ, Cohen S. Inhibition of lower esophageal sphincter circular muscle by female sex hormones. *Am J Physiol.* 1978;234:E243–E247.
53. Wald A, Van Thiel DH, Hoechstetter L, et al. Gastrointestinal transit: the effect of the menstrual cycle. *Gastroenterology.* 1981;80:1497–1500.
54. Turnbull GK, Thompson DG, Day S, Martin J, Walker E, Lennard-Jones JE. Relationships between symptoms, menstrual cycle and orocaecal transit in normal and constipated women. *Gut.* 1989;30:30–34.
55. Heitkemper M, Jarrett M. Pattern of gastrointestinal and somatic symptoms across the menstrual cycle. *Gastroenterology.* 1992;102:505–513.
56. Moore J, Barlow D, Jewell D, Kennedy S. Do gastrointestinal symptoms vary with the menstrual cycle? *Br J Obstet Gynaecol.* 1998;105:1322–1325.
57. Widom B, Diamond MP, Simonson DC. Alterations in glucose metabolism during menstrual cycle in women with IDDM. *Diabetes Care.* 1992;15:213–220.
58. Letterie GS, Fredlund PN. Catamenial insulin reactions treated with a long-acting gonadotropin releasing hormone agonist. *Arch Intern Med.* 1994;154:1868–1870.
59. Walsh CH, Malins JM. Menstruation and control of diabetes. *Br Med J.* 1977;2:177–179.
60. Hubble D. Insulin resistance. *Br Med J.* 1954;2:1022.
61. Bertoli A, De Pirro R, Fusco A, Greco AV, Magnatta R, Lauro R. Differences in insulin receptors between men and menstruating women and influence of sex hormones on insulin binding during the menstrual cycle. *J Clin Endocrinol Metab.* 1980;50:246–250.
62. Jarrett RJ, Graver HJ. Changes in oral glucose tolerance during the menstrual cycle. *Br Med J.* 1968;2:528–529.
63. Tomelleri R, Gruewald K. Menstrual cycle and food cravings in young college women. *J Am Diet Assoc.* 1987;87:311–315.
64. Maurer ER, Schaal FA, Mendez FL. Chronic recurring spontaneous pneumothorax due to endometriosis of the diaphragm. *JAMA.* 1958;168:2013–2024.
65. Markham SM, Carpenter SE, Rock JA. Extrapelvic endometriosis. *Obstet Gynecol Clin North Am.* 1989;16:193–207.
66. Slabbynck H, Laureys M, Impens N, De Vroey P, Schandevyl W. Recurring catamenial pneumothorax treated with a GnRH analogue. *Chest.* 1991;100:851.
67. Arnbjornsson E. Acute appendicitis risk in various phases of the menstrual cycle. *Acta Chir Scand.* 1983;149:603–605.
68. Arnbjornsson E. The influence of oral contraceptives on the frequency of acute appendicitis in different phases of the menstrual cycle. *Surg Gynecol Obstet.* 1984;158:464–466.
69. Dalton K. Influence of menstruation on glaucoma. *Br J Ophthalmol.* 1967;51:692–695.
70. Smith R, Studd JW. A pilot study of the effect upon multiple sclerosis of the menopause, hormone replacement therapy and the menstrual cycle. *J R Soc Med.* 1992;85:612–613.
71. Caughey SC, Margesson LJ, Reid RL. Erythema multiforme with ulcerative stomatitis at menstruation. *Can J Obstet Gynecol.* 1990;112–113.
72. Howard RE. Premenstrual urticaria and dermatographism. *JAMA.* 1981;245:1068.
73. Rosano GMC, Leonardo F, Sarrel P, Beale CM, De Luca F, Collins P. Cyclical variation in paroxysmal supraventricular tachycardia in women. *Lancet.* 1996;347:786–88.
74. Anderson KE, Spitz IM, Sassa S, Bardin W, Kappas A. Prevention of cyclical attacks of acute intermittent porphyria with a long-acting agonist of luteinizing hormone-releasing hormone. *N Eng J Med.* 1984;311:643–645.
75. Meggs WJ, Pescovitz OH, Metcalfe D, Loriaux DL, Cutler G Jr., Kaliner M. Progesterone sensitivity as a cause of recurrent anaphylaxis. *N Eng J Med.* 1984;311:1236–1238.
76. Slater JE, Raphael G, Cutler GB Jr., Loriaux DL, Meggs WJ, Kaliner N. Recurrent anaphylaxis in menstruating women: treatment with a luteinizing hormone-releasing hormone agonist—a preliminary report. *Obstet Gynecol.* 1987;70:542–546.
77. Vijayan N, Vijayan VK, Dreyfuss PM. Acetylcholinesterase activity and menstrual remissions in myasthenia gravis. *J Neurol Neurosurg Psychiatry.* 1977;40:1060–1065.
78. Badwe RA, Gregory WM, Chaudary MA, et al. Timing of surgery during menstrual cycle and survival of premenopausal women with operable breast cancer. *Lancet.* 1991;337:1261–1264.
79. Wobbes T, Thomas CM, Segers MF, Peer PG, Bruggink ED, Beex LV. The phase of the menstrual cycle has no influence on the disease-free survival of patients with mammary carcinoma. *Br J Cancer.* 1994;69:599–600.