

ORIGINAL ARTICLE

Meta-analysis reveals an association of *PTPN22* C1858T with autoimmune diseases, which depends on the localization of the affected tissue

J Zheng¹, S Ibrahim², F Petersen³ and X Yu^{1,3}

Protein tyrosine phosphatase non-receptor type 22 (*PTPN22*) is a strong susceptibility gene shared by many autoimmune diseases. The aim of this study was to explore the mechanisms underlying this relationship. We performed a comprehensive analysis of the association between *PTPN22* polymorphism C1858T and autoimmune diseases. The results showed a remarkable pattern; *PTPN22* C1858T was strongly associated with type I diabetes, rheumatoid arthritis, immune thrombocytopenia, generalized vitiligo with concomitant autoimmune diseases, idiopathic inflammatory myopathies, Graves' disease, juvenile idiopathic arthritis, myasthenia gravis, systemic lupus erythematosus, anti-neutrophil cytoplasmic antibody-associated vasculitis and Addison's disease. By contrast, *PTPN22* C1858T showed a negligible association with systemic sclerosis, celiac disease, multiple sclerosis, psoriasis, ankylosing spondylitis, pemphigus vulgaris, ulcerative colitis, primary sclerosing cholangitis, primary biliary cirrhosis, Crohn's disease and acute anterior uveitis. Further analysis revealed a clear distinction between the two groups of diseases with regard to their targeted tissues: most autoimmune diseases showing an insignificant association with *PTPN22* C1858T manifest in skin, the gastrointestinal tract or in immune privileged sites. These results showed that the association of *PTPN22* polymorphism with autoimmune diseases depends on the localization of the affected tissue, suggesting a role of targeted organ variation in the disease manifestations.

Genes and Immunity (2012) 13, 641–652; doi:10.1038/gene.2012.46; published online 18 October 2012

Keywords: protein tyrosine phosphatase non-receptor type 22; autoimmune diseases; meta-analysis; susceptibility gene; target tissue; association

INTRODUCTION

Autoimmune diseases are influenced by a combination of genetic predisposition and environmental factors and are mostly poly-genic disorders to which many genes contribute. Human leukocyte antigen (HLA) loci are associated with all autoimmune diseases and are the strongest genetic factors for individual predisposition.¹ Of the non-HLA susceptibility genes, variants of protein tyrosine phosphatase non-receptor type 22 (*PTPN22*) show the strongest associations with autoimmune diseases. Initially identified as a susceptibility gene for type 1 diabetes (T1D),² variants of *PTPN22*, especially C1858T, have subsequently been shown to be associated with many other autoimmune diseases such as rheumatoid arthritis (RA), systemic lupus erythematosus (SLE) and Graves' disease.^{3–5}

Given the important role of *PTPN22* in many autoimmune diseases, exploration of the mechanism(s) behind this association will help us to better understand the pathogenesis of disease. The human *PTPN22* gene is located on chromosome 1 and encodes a protein (called Lyp) comprising 807 amino-acid residues,⁶ which contains a catalytic domain and four SH3 binding sites. The R620W mutation, resulting from the C1858T polymorphism, is located within the first SH3 binding domain of Lyp, indicating that it exerts its effects by affecting intermolecular interactions.² Indeed, binding of Lyp to C-terminal Src tyrosine kinase (Csk) is partially or completely disrupted, and is associated with SLE, T1D and RA.^{2,3} Vang *et al.*⁷ reported that R620W is a gain-of-function mutation,

increasing catalytic activity and acting as a potent negative regulator of T lymphocyte activation. Further studies showed that that the R620W mutation also inhibited activation of NK cell and B cell.^{8,9} Although considerable progress has been achieved in examining the function of the R620W mutation, the mechanisms underlying the association between *PTPN22* and autoimmune diseases are still largely unknown. Studies in mice provided only limited information to elucidate these regulatory processes. Murine *PTPN8*, which represents the ortholog of human *PTPN22* in mice, was successfully knocked out but the corresponding animal did not develop spontaneous autoimmune diseases.¹⁰ There may be two reasons for this, first, R620W is a gain-of-function mutation and knockout mice (essentially a loss-of-function model) are of no help. Second, a high divergence between the human *PTPN22* and mouse *PTPN8* genes exists, which may result in functional differences of the molecule in both species.

With regard to the strong association between *PTPN22* polymorphisms and certain autoimmune diseases, extensive studies have been undertaken to evaluate the association between *PTPN22* variants and more than 20 such diseases. Although *PTPN22* is the one of the strongest non-HLA susceptibility genes for T1D, RA, SLE and Graves' disease,^{2–5} no association has been observed between *PTPN22* variants and other autoimmune diseases such as systemic sclerosis (SSc), Crohn's disease, ulcerative colitis (UC) and multiple sclerosis (MS).^{11–13}

¹Laboratory of Autoimmunity, The Medical College of Xiamen University, Xiamen University, Xiamen, China; ²Department of Dermatology, University of Lübeck, Lübeck, Germany and ³Division of Biochemical Immunology, Department of Asthma & Allergy, Research Center Borstel, Borstel, Germany. Correspondence: Dr X Yu, Laboratory of Autoimmunity, The Medical College of Xiamen University, Xiamen University, 361005 Xiamen, China.
E-mail: xinhuayu@fz-borstel.de

Received 13 April 2012; revised 23 August 2012; accepted 24 August 2012; published online 18 October 2012

Examining the association pattern between *PTPN22* variants and autoimmune diseases will help us to understand the pathogenesis of individual diseases. It may also help us to understand the mechanism(s) underlying this association. Therefore, we for the first time performed a comprehensive meta-analysis to identify the association pattern between *PTPN22* and autoimmune diseases. We then evaluated this association pattern in detail with the aim of shedding new light on the role of *PTPN22* in autoimmune disease.

RESULTS

Frequency of the *PTPN22* minor allele

The frequency of the *PTPN22* C1858T minor allele varies considerably across the world (Figure 1). In brief, European populations or populations of European descent showed a higher frequency than other populations, and the frequency in northern Europe was higher than in the south. The highest T allele frequency (17.5%) was reported in northeast Russia,¹⁴ followed by Finland (15.4%),¹⁵ Ukraine (14.1%)¹⁶ and Estonia (13.9%).¹⁷ The frequency decreased toward the west, showing 11–13% in Poland,^{18–20} 10–12% in Sweden,^{21–23} Norway^{24–26} and Germany,^{24,27–29} 11.7% in Croatia,³⁰ 10–11% in the Czech Republic,^{30,31} 10.7% in Slovakia,³² 9.2% in Denmark,³³ and 8–10% in the UK^{34–39} and the Netherlands.^{24,40–42} The decrease in the frequency toward the south was dramatic, with 7.75% in Hungary,⁴³ 7.23% in Belgium, 7–8% in France,^{44,45} 5–7% in Spain^{46–48} and 4.1% in Romania.⁴⁹ The frequency in Italy varied from 2.1 to 5.9%.^{2,50–53} The lowest frequency for the *PTPN22* T allele was reported in two island populations: Sassari, Italy (2.1%)² and Crete, Greece (2.9%).⁵⁴ Outside Europe, the T allele frequency in populations of European descent in New Zealand ranged from 9.9 to 10.7%,^{24,55} and in America it ranged from 3.8 to 10.0%.^{49,56–67} Americans of non-European descent showed a very low T allele frequency, as low as 1.1% in Mexico.⁶⁸ North Africa had a similar T allele frequency to Turkey and the Middle East, which ranged from 2 to 3.5%;^{69–75} however, *PTPN22* is not polymorphic in the black

African population.⁷⁶ *PTPN22* is also non-polymorphic or almost non-polymorphic in some Asian populations, including Japanese, Korean, Indian and some Chinese populations.^{77–81} Two Chinese populations, Uygers and Kazaks, showed relatively high T allele frequencies of 5.3% and 7.0%, respectively; both are Caucasian-related populations.⁸¹

Diseases showing a strong association with *PTPN22*

Although not proven, it is believed that the *PTPN22* C1858T polymorphism is the main causal mutation for previously described associations of *PTPN22* with autoimmune diseases.⁸² Therefore, we focused this meta-analysis only on the C1858T polymorphism. Due to dramatic differences in the frequency of the T allele between populations, most case-control studies have been conducted in European populations and populations of European descent. Only a few studies have been conducted in other populations such as North African, Turkish, Mexican and Indian, and then with only a small number of samples.^{70–72,74,75,83,84} Therefore, to avoid any possible heterogeneity, we excluded these studies and focused on studies in European or European-descended populations.

The association between *PTPN22* C1858T and autoimmune diseases was first identified in individuals with T1D in 2004.² Numerous studies have been performed across the world to confirm this association. Twenty-three case-control studies were recruited for this meta-analysis with odds ratios (ORs) ranging from 1.04 to 3.15.^{2,5,14,16,17,30,33,42,51,52,57,60,63,65,85–91} Only 5/23 studies failed to reach a significant threshold. As expected, the meta-analysis showed a strong association between the T allele and T1D (OR = 1.84; 95% confidence intervals (CI) = 1.72–1.96, $P < 1.0 \times 10^{-16}$) (Figure 2a).

As the most common autoimmune disease, RA is also the most extensively investigated in terms of its association with *PTPN22* C1858T. In total, 36 case-control studies were recruited for the present study with ORs ranging from 0.99 to 2.56.^{3,14,15,18,22,24,26,28,32,35,39–42,54,60,67,92–103} Thirty-one out of thirty-six showed a significant association between *PTPN22* and




Figure 1. Heat map of the allele frequency of the *PTPN22* minor allele across the world. Each color-filled circle represents a single population. Only populations with a clearly defined location are presented. All populations from USA, Canada, South America and New Zealand are of European descent.




Figure 2. Forest plot of case-control studies of nine autoimmune diseases showing a strong association with *PTPN22*. The nine autoimmune diseases include (a) type 1 diabetes, (b) rheumatoid arthritis, (c) systemic lupus erythematosus, (d) Graves' disease, (e) Addison's disease, (f) ANCA-associated vasculitis, (g) juvenile idiopathic arthritis, (h) generalised vitiligo, and (i) myasthenia gravis. The information regarding individual case-control studies, including the first author name and year of publication, as well as the number of cases/number of controls, are indicated. OR and 95% CI of individual studies are represented by squares and horizontal lines, respectively. The size of the square represents the weight of the individual study in the meta-analysis. The diamonds represent the OR (center line of the diamond) and 95% CI (lateral tips of the diamond) of the fixed-effect meta-analysis. The OR, 95% CI and *P*-values for the meta-analysis are indicated by bold text.

RA. Meta-analysis showed a strong association between the T allele and RA (OR = 1.65; 95% CI = 1.58–1.71, $P < 1.00 \times 10^{-16}$) (Figure 2b).

Another common autoimmune disease, SLE has also been extensively studied. Of the 14 case-control studies recruited for this study,^{4,54,60,103–108} 9 showed a significant association between *PTPN22* C1858T and SLE. Meta-analysis showed a strong association between the T allele and SLE (OR = 1.46; 95%

CI = 1.31–1.62, $P = 7.33 \times 10^{-13}$), although this was not as strong as that between T1D and *PTPN22* C1858T or RA and *PTPN22* C1858T (Figure 2c).

Even though it has not been not extensively evaluated, a strong association between Graves' disease and *PTPN22* C1858T has also been observed. All three case-control studies recruited for this meta-analysis showed a significant association between *PTPN22* C1858T and Graves' disease with ORs ranging from 1.43

to 1.88.^{5,20,109} Meta-analysis showed a strong association between the T allele and Graves' disease (OR = 1.59; 95% CI = 1.37–1.85, $P = 1.01 \times 10^{-9}$) (Figure 2d).

Six case-control studies evaluated the association between *PTPN22* C1858T and Addison's disease. Three showed a significant association, whereas the other three studies showed a tendency toward a positive association but failed to reach the significance threshold.^{25,87,109,110} Meta-analysis showed a significant association between the T allele and Addison's disease (OR = 1.43; 95% CI = 1.21–1.68, $P = 2.36 \times 10^{-5}$) (Figure 2e).

Anti-neutrophil cytoplasmic antibody (ANCA)-associated vasculitis includes Wegener's granulomatosis, microscopic polyangiitis and Churg-Strauss syndrome. Two case-control studies were recruited for this meta-analysis.^{111,112} Meta-analysis showed a significant association between the T allele and ANCA-associated vasculitis (OR = 1.45; 95% CI = 1.24–1.69, $P = 3.07 \times 10^{-6}$) (Figure 2f).

Seven case-control studies were undertaken to investigate the association between *PTPN22* C1858T and juvenile idiopathic arthritis (JIA), five of which revealed a significant association.^{15,30,43,66,103,113} Meta-analysis showed a highly significant association between the T allele and JIA (OR = 1.54; 95% CI = 1.40–1.70, $P < 1.00 \times 10^{-16}$) (Figure 2g).

Only two case-control studies evaluated the association between generalized vitiligo (GV) and *PTPN22* C1858T and both were recruited for the meta-analysis. Both studies suggested an association between GV and *PTPN22* C1858T.^{34,49} Meta-analysis showed a strong association between the T allele and GV (OR = 1.98; 95% CI = 1.35–2.88, $P = 3.70 \times 10^{-4}$) (Figure 2h). This association is supported by a recent genome-wide association studies (GWAS).¹¹⁴

Case-control studies have also identified a strong association between *PTPN22* C1858T and myasthenia gravis (MG). Meta-analysis of five studies^{23,45,115,116} showed a significant association between the T allele and MG (OR = 1.53; 95% CI = 1.31–1.80, $P = 1.09 \times 10^{-7}$) (Figure 2i).

Diseases showing no or weak association with *PTPN22*

After the identification of *PTPN22* C1858T as a susceptibility gene for autoimmune disease, the association between *PTPN22* C1858T and inflammatory bowel disease was extensively investigated, including Crohn's disease and UC. Of the 11 case-control studies in Crohn's disease patients, only 2 identified an association with *PTPN22*.^{12,21,29,31,37,50,55,58,67} Meta-analysis showed a very




Figure 3. Forest plot of case-control studies of nine autoimmune diseases showing a weak or no association with *PTPN22*. The nine autoimmune diseases include (a) Crohn's disease, (b) ulcerative colitis, (c) systemic sclerosis, (d) multiple sclerosis, (e) celiac disease, (f) psoriasis, (g) psoriatic arthritis, (h) alopecia areata, and (i) primary Sjögren's syndrome. The detailed information in the forest plot is explained in the legend to Figure 2.

weak association between the T allele and a reduced risk of Crohn's disease (OR = 0.84; 95% CI = 0.76–0.94, $P = 1.89 \times 10^{-3}$) (Figure 3a). Seven case–control studies on UC patients were recruited, six of which showed no association between *PTPN22* and UC.^{12,21,29,37,50,67,117} Meta-analysis showed no association between the T allele and UC (OR = 0.96; 95% CI = 0.87–1.05, $P = 0.347$) (Figure 3b).

Another extensively studied disease is SSc, an autoimmune disease causing extensive fibrosis and vascular alterations in the skin.¹³ Eleven case–control studies have been performed, but only one identified a significant association between *PTPN22* C1858T and SSc.^{13,46,61,118} Meta-analysis showed a very weak association between the T allele and SSc (OR = 1.16; 95% CI = 1.02–1.31, $P = 0.015$) (Figure 3c). This is consistent with the findings of a GWAS, which failed to identify an association between *PTPN22* C1858T and SSc.¹¹⁹

MS is a common autoimmune disease that targets the central nervous system. A study of family-based samples showed no association between *PTPN22* C1858T and MS.¹¹ This lack of association was confirmed by a GWAS.¹²⁰ Meta-analysis of two case–control studies^{96,121} did not identify any association between *PTPN22* C1858T and MS (OR = 1.11; 95% CI = 0.85–1.44, $P = 0.442$) (Figure 3d).

Another autoimmune disease, celiac disease, affects the gastrointestinal tract. No association with *PTPN22* has been identified. Of the four studies recruited for this meta-analysis, three showed no association between *PTPN22* C1858T and celiac disease.^{42,47,48,92} Meta-analysis also showed no association between *PTPN22* C1858T and celiac disease (OR = 1.12; 95% CI = 0.97–1.29, $P = 0.125$) (Figure 3e).

Psoriasis also showed no association with *PTPN22* C1858T according to our meta-analysis. Meta-analysis of three case–control studies with psoriasis^{96,122,123} failed to identify any association with the T allele (OR = 1.08; 95% CI = 0.93–1.27, $P = 0.306$) (Figure 3g), which is consistent with the findings of a GWAS.¹²⁴ A psoriasis-related autoimmune disease, called psoriasis arthritis, is weakly associated with *PTPN22* C1858T. Meta-analysis of four case–control studies identified a weak association between the T allele and psoriasis arthritis (OR = 1.22; 95% CI = 1.02–1.44, $P = 0.023$) (Figure 3e).^{56,96,125}

Two case–control studies have been performed for alopecia areata (AA). One suggested a weak association with *PTPN22* C1858T, but the other did not.^{27,36} Meta-analysis showed a weak association between AA and the T allele (OR = 1.38; 95% CI = 1.11–1.72, $P = 0.003$) (Figure 3f). This weak association is in line with the findings of a GWAS conducted in a Caucasian population.¹²⁶

The association between *PTPN22* C1858T and primary Sjogren's syndrome (pSS) is not clear. One study from Columbia (70 cases and 308 controls) identified a strong association (OR = 2.42; 95% CI = 1.23–4.75, $P = 0.01$); this was similar to that for SLE, but stronger than that for RA or T1D in the same population.⁶⁰ However, another study from France (183 cases and 172 controls) failed to show any association.⁴⁴ Meta-analysis of both these studies showed a relatively high OR for the T allele (OR = 1.40; 95% CI = 0.91–2.14), but it failed to reach the significance threshold ($P = 0.12$). As no GWAS has been performed for pSS, more case–control studies are required to clarify any association between *PTPN22* C1858T and pSS.

Stratified analyses based on autoantibodies

The first study to report an association between RA and *PTPN22* C1858T showed that it is associated with rheumatoid factor positive (RF+) RA but not with RF– RA,³ raising the possibility that *PTPN22* C1858T affects autoantibody production. This stratified analysis, based on autoantibody status, was performed in some of the case–control studies, allowing us to conduct a

stratified meta-analysis to explore the role of *PTPN22* C1858T in autoantibody production. To better evaluate the association between *PTPN22* C1858T and autoantibody status, we performed stratified meta-analysis using three comparisons: antibody-positive cases vs controls, antibody-negative cases vs controls and antibody-positive cases vs antibody-negative cases (Figure 3). Stratified analyses have been extensively evaluated in patients with RA based on the levels of RF or anti-cyclic citrullinated peptide (anti-CCP) antibodies. Meta-analysis of 11 RA case–control studies^{18,22,28,40,41,95,97–99,101,102} according to RF status showed a strong association between the T allele and RF+ RA (OR = 1.63; 95% CI = 1.51–1.75, $P < 1.00 \times 10^{-16}$) and a weak association between the T allele and RF– RA (OR = 1.35; 95% CI = 1.22–1.49, $P = 9.04 \times 10^{-9}$). A comparison between RF+ RA and RF– RA showed that the T allele was more common in RF+ RA (OR = 1.21; 95% CI = 1.09–1.35, $P = 2.79 \times 10^{-4}$) (Supplementary Figure 1a). Five RA case–control studies based on anti-CCP antibody status were recruited for the meta-analysis.^{18,22,28,40,101} The results showed a strong association between the T allele and anti-CCP+ RA (OR = 1.78; 95% CI = 1.59–2.01, $P < 1.00 \times 10^{-16}$) but no association between the T allele and anti-CCP– RA (OR = 1.15; 95% CI = 0.98–1.35, $P = 0.081$). Also the frequency of T allele was significantly higher in anti-CCP+ RA than in anti-CCP– RA (OR = 1.45; 95% CI = 1.24–1.70, $P = 4.09 \times 10^{-6}$) (Supplementary Figure 1b).

SSc is an autoimmune disease characterized by high levels of autoantibodies against various cellular antigens including anti-topoisomerase I antibodies (ATA) and anticentromere antibodies (ACA).¹¹⁸ Stratified analyses have been performed based on ATA and ACA levels in 11 case–control studies.^{13,46,61,118} The present meta-analysis identified a marginal association between SSc and *PTPN22*. Stratified meta-analysis based on ATA status showed a marginal association between both ATA+ SSc (OR = 1.25; 95% CI = 1.03–1.54, $P = 0.037$) and ATA– SSc (OR = 1.16; 95% CI = 1.00–1.33, $P = 0.019$) and *PTPN22* (Figure 4d). Comparison between ATA+ SSc and ATA– SSc did not reveal significant differences in the T allele frequency (OR = 1.11; 95% CI = 0.90–1.37, $P = 0.323$) (Supplementary Figure 1c). Similar results were observed in the stratified meta-analysis based on ACA status, where the T allele was marginally associated with both ACA+ SSc (OR = 1.28; 95% CI = 1.07–1.52, $P = 0.0058$) and ACA– SSc (OR = 1.14; 95% CI = 1.00–1.31, $P = 0.053$). There was no difference in the frequency of the T allele between ACA+ SSc and ACA– SSc (OR = 1.15; 95% CI = 0.96–1.38, $P = 0.127$) (Supplementary Figure 1d).

Association between *PTPN22* and other autoimmune diseases

Besides the autoimmune diseases mentioned above, an association between *PTPN22* and several other autoimmune diseases has been investigated, including ankylosing spondylitis (AS), primary sclerosing cholangitis (PSC), primary biliary cirrhosis (PBC), idiopathic inflammatory myopathies (IIM), acute anterior uveitis (AAU), immune thrombocytopenia and pemphigus vulgaris (PV).^{64,103,127–131} The results are summarized in Supplementary Table S1. Only one case–control study was conducted for each disease in European populations or populations of European descent, making meta-analysis impossible. However, evidence from GWAS or case–control studies conducted in other populations provides additional evidence for an association.

A case–control study performed by Orozco *et al.*¹³¹ in 2005 did not find an association between *PTPN22* C1858T and AS, a highly HLA B27-associated autoimmune disease. This lack of association between *PTPN22* C1858T and AS was confirmed in a GWAS.¹³² In addition, no association with *PTPN22* C1858T was reported for another HLA B27-associated autoimmune disease, AAU.¹²⁹ PBC is an autoimmune disease characterized by the slow progressive destruction of the small bile ducts within the liver.¹³³ In 2006, Milkiewicz *et al.*¹³⁰ reported that *PTPN22* was not associated with




Figure 4. The pattern of association between *PTPN22* and autoimmune diseases. **(a)** Association between *PTPN22* and 26 autoimmune diseases. OR with 95% CI for the *PTPN22* T allele are presented. **(b)** Analysis of the target tissue specificity of 12 autoimmune diseases showing a strong association with *PTPN22* and 12 autoimmune diseases showing a negligible association with *PTPN22*. The lines connecting diseases and tissues indicate the target tissue for the associated disease, whereas the solid lines connect diseases targeting one tissue and dashed lines connect diseases targeting multiple tissues.

PBC in Canada. This result was confirmed by a GWAS in 2009 with samples from both the USA and Canada.¹³³ PSC is another autoimmune disease targeting the small bile ducts within the liver, and most patients with PSC also have UC.¹³⁴ As is the case for UC and PBC, *PTPN22* is not associated with PSC.¹⁰³ PV is caused by antibodies against desmoglein 1 or desmoglein 3, resulting in the loss of cohesion between keratinocytes and subsequent skin blisters.¹³⁵ In 2011, Sachdev *et al.*⁶⁴ reported no association between *PTPN22* C1858T and PV, confirming a previous observation in a Tunisian population.¹³⁶

A strong association between *PTPN22* C1858T and two other autoimmune diseases has been reported. Immune thrombocytopenia is characterized by antibody-mediated platelet destruction. D'silva *et al.*¹²⁸ reported a strong association between ITP and the *PTPN22* T allele in a caucasian US population. This association was confirmed in an Egyptian population.⁷⁰ Finally, a strong association between the *PTPN22* T allele and IIM was reported in a British Caucasian population.¹²⁷

Association pattern between *PTPN22* and autoimmune diseases

So far, the association between *PTPN22* C1858T and 25 autoimmune diseases has been evaluated. Figure 4a summarizes the ORs for the T allele of the different diseases in a ranked order. The *PTPN22* T allele is strongly associated with GV, ITP, T1D, RA, GD, JIA, MG, SLE, ANCA-associated vasculitis and Addison disease.

The association between pSS and *PTPN22* needs further evaluation. A weak association was observed between *PTPN22* and psoriasis arthritis and AA, and a negligible association was observed between *PTPN22* and SSc, celiac disease, MS, psoriasis, AS, PV, UC, PSC, PBC, Crohn's disease and AAU.

To further examine the association pattern of *PTPN22* C1858T with autoimmune diseases, we evaluated the relationship between association status and the target tissues affected by these autoimmune diseases. We focused on diseases showing a strong association with *PTPN22* and diseases showing a negligible association with *PTPN22*. As a GWAS showed that *PTPN22* is only associated with GV in patients also suffering from other autoimmune diseases, but not associated with GV in patients with GV alone,¹¹⁴ we divided GV into two subsets, including GV with concomitant autoimmune diseases and GV without concomitant autoimmune diseases. This analysis identified a fine target tissue specificity between diseases showing a strong association with *PTPN22* C1858T and diseases showing a negligible association with *PTPN22* (Figure 4b). The target tissues of diseases showing a strong association with *PTPN22* C1858T include skin (SLE and GV with other concomitant autoimmune diseases), blood (immune thrombocytopenia), pancreas (T1D and GV with other concomitant autoimmune diseases), kidney (SLE), connective tissue (RA), thyroid (GD), joint (JIA, RA, SLE and GV with other concomitant autoimmune diseases), muscle (MG and IIM), blood vessels (SLE and ANCA-associated vasculitis) and adrenal

glands (Addison's disease and GV with other concomitant autoimmune diseases). The target tissues of diseases showing a negligible association with *PTPN22* C1858T include skin (GV without any concomitant autoimmune diseases, SSc, psoriasis and PV), gastrointestinal tract (celiac disease, UC, Crohn's disease), bile duct (PSC and PBC), spine (AS) and two immune privileged tissues; the central nervous system (MS) and eye (AAU). Interestingly, although some tissues are the target for several autoimmune diseases, there is a clear distinction between those diseases showing a strong association with *PTPN22* C1858T and those diseases showing a weak or negative association with *PTPN22* C1858T. There is only one exception: skin is the target tissue for four autoimmune diseases (GV without concomitant autoimmune diseases, SSc, psoriasis and PV) that show a negligible association with *PTPN22* C1858T, and two autoimmune diseases showing a strong association with *PTPN22* C1858T (SLE and GV with concomitant autoimmune diseases). Notably, both SLE and GV with other concomitant autoimmune diseases target many tissues, indicating that skin might not be their specific target tissue. Therefore, when we exclude those autoimmune diseases that target several tissues, we see a clear distinction between autoimmune diseases showing a strong association with *PTPN22* C1858T and diseases showing a negligible association with *PTPN22* C1858T.

DISCUSSION

The comprehensive meta-analysis carried out in the present study showed that *PTPN22* C1858T is strongly associated with nine autoimmune diseases: T1D, RA, SLE, Graves' disease, Addison's disease, ANCA-associated vasculitis, JIA, MG and GV with other concomitant autoimmune diseases. Together with 2 other *PTPN22* C1858T-associated diseases (ITP and IIM), which are not included in present meta-analysis, 11 autoimmune diseases are strongly associated with *PTPN22* C1858T, with the T allele increasing the risk of disease. The association suggests that there may be a common mechanism shared by these diseases. Exploration of these mechanisms will greatly help us to understand the pathogenesis of the diseases.

Based on two observations, a possible underlying mechanism affecting autoantibody production by *PTPN22* variants has been suggested.⁸² First, some primary T-cell-mediated autoimmune diseases, for example, Crohn's disease, UC and MS, are not associated with *PTPN22* C1858T,^{11,12} whereas autoantibody-associated diseases, for example, T1D, RA and SLE, are associated.²⁻⁴ Second, several stratified analyses show that *PTPN22* C1858T is only associated with RF+ RA or anti-CCP RA+ but not with RF- RA or anti-CCP- RA.^{3,40,101} However, although our analysis could indeed confirm this hypothesis for some diseases others did not fit into this model. Stratified meta-analysis for RA showed that *PTPN22* C1858T is associated with anti-CCP+ RA but not with anti-CCP- RA, and the association between *PTPN22* C1858T and RF+ RA is much stronger than that between *PTPN22* C1858T and RF- RA, suggesting that the C1858T mutation might affect the production of anti-CCP antibodies and RF. By contrast, the stratified meta-analysis for SSc did not confirm an association between *PTPN22* and ATA or ACA autoantibodies. Moreover, *PTPN22* C1858T is not associated with autoantibody status in T1D patients.¹³⁷ Finally, PV, which is an autoimmune disease mediated by autoantibodies against desmoglein 1 and desmoglein 3, is not associated with *PTPN22* C1858T; refuting the idea that *PTPN22* C1858T is associated with autoantibody status. Taken together, we have to conclude that the *PTPN22* C1858T polymorphism is in some but not all cases associated with the presence autoantibodies and that development of self-reacting antibodies is not direct consequence of this gene variation.

According to a second hypothesis, the variations C1858T could lead to enhanced generation of autoreactive T cells escaping

thymic selection.¹³⁷ As this gain-of-function mutation weakens TCR signaling,^{2,3,7} it is conceivable that it will result in some moderately autoreactive T cells escaping the thymic selection process. Furthermore, the C1858T variation also weakens BCR signaling in B cells⁸ and, therefore, this model can be expanded to B cells as well. As a consequence, increased numbers of autoreactive T and B cells could be responsible for an enhanced risk of developing autoimmune diseases. This new model is supported by experimental evidence from a very recent breakthrough by Menard *et al.*¹³⁸ In this study, Menard *et al.*¹³⁸ showed that the R620W variation interferes with the removal of self-reacting B cells in humans by affecting both central and peripheral B-cell tolerance check points. This study clearly shows that the R620W mutation increases the generation of autoreactive B cells, which supports the above proposed model. However, because *PTPN22* as a cell signaling regulator is expressed in many immune cells, it is unlikely that *PTPN22* should affect only the generation of autoimmune T and B cells. Therefore, the mechanism behind the association could be much more complex than what this model proposed. Very recently, Zhang *et al.*¹³⁹ investigated knock-in mice and showed that the C1858T allele affects T-cell activation and positive selection, as well as the activation of dendritic- and B-cell activation. These results provide strong evidence that *PTPN22* C1858T could affect autoimmune response on several levels, including generation and activation of autoimmune cells, and in many different cell types, including T and B cells and dendritic cells. Therefore, this model need to be further revised as: *PTPN22* appears to have a role in autoimmune disease manifestations by affecting the generation and activation of immune cells.

Given a role of *PTPN22* in generation and activation of immune cells, the question arises why several autoimmune diseases lack any association to *PTPN22*. Our analysis indicates that the C1858T mutation has a negligible effect on 12 autoimmune diseases: Crohn's disease, UC, celiac disease, SSc, MS, psoriasis, PSC, PBC, AS, PV, AAU and GV without other concomitant autoimmune diseases. Together with the 11 autoimmune diseases that show a strong association with *PTPN22* C1858T, a remarkable distribution pattern becomes visible that suggests a novel mechanism how some diseases could be related to *PTPN22* variations. If associations are considered with regard to the targeted tissues, a clear distinction between diseases showing a strong association with *PTPN22* C1858T and those showing a negligible association was found. Interestingly, the target tissues for most diseases showing a negligible association with *PTPN22* are skin (in GV without concomitant other autoimmune diseases, SSc, psoriasis and PV), the gastrointestinal tract (in celiac disease, UC and Crohn's disease) and some immune privileged sites such as the central nervous system (in MS) and the eye (in AAU). Compared with other tissues, both skin and the gastrointestinal tract have exposed surfaces and therefore may be more affected by environmental factors. This raises the possibility that autoimmune diseases associated with these tissues might be triggered by some environmental factor(s), for example, bacterial infection. One typical example is Crohn's disease. Nucleotide-binding oligomerization domain-containing protein 2 (*NOD2*), a receptor of bacterial peptidoglycan, has been identified as the strongest non-HLA susceptibility gene for Crohn's disease,^{140,141} indicating a role for bacteria in the development of this disease. Regarding immune privileged tissues, some proteins responsible for triggering the deletion of autoreactive T and B cells are not expressed in the central immune system, allowing autoreactive lymphocytes to escape from central selection.^{142,143} Under normal physiological conditions, these autoreactive T and B cells are inactive because they are unable to enter immune privileged tissues; however, when immune privilege is disturbed, these autoreactive T and B cells may gain access to these tissues and cause an autoimmune response.




Figure 5. A model of the mechanism underlying the association between *PTPN22* and autoimmune diseases. In healthy individuals, autoimmune response is a tolerant state and does not attack target tissue. However, the break of such tolerance can be triggered by at least two group of factors. The first group consists of factors acting on immune cells, and the other group consists of factors acting on target tissues. *PTPN22* belongs to the first group.

Based on previous knowledge that *PTPN22* affects the generation and activation of immune cells and our own findings regarding the association pattern and target tissue specificity, we propose a new model explaining the association between *PTPN22* C1858T and autoimmune diseases (Figure 5). In a healthy state, some autoreactive immune cells may escape from the central immune system. However, they are not active because they show low autoreactivity, lack access to the target antigens or are suppressed by regulatory T cells. Therefore, the autoimmune response will not manifest in the organ. Autoimmune disease (as the result of a misdirected autoimmune response) may be triggered in two ways. First, autoimmune diseases can be triggered by factors acting on immune cells, resulting in the generation of autoimmune cells or activation of immune cells. This may cause unwanted autoimmune responses and a consequent development of autoimmune diseases. The second way involves factors acting on the target tissues. For example, bacterial infection or environmental noxa may lead to a proinflammatory microenvironment in the target tissue resulting in an activation of self-reactive T and B cells with principal low autoreactivity. Such factors could also be involved in the disturbance of immune privilege, for example, by changing the permeability of the brain–blood barrier, enabling the entrance of autoreactive lymphocytes into these tissues and initiating autoimmune disease. Within this view, the main effect of *PTPN22* in autoimmunity maybe its capacity to affect immune cells. This could explain our observation that *PTPN22* C1858T appears to be strongly associated with those autoimmune diseases, which are primarily triggered by factors dysregulating immune cells, whereas no or only low association is observed in autoimmune diseases primarily triggered by factors acting on target tissues.

In summary, our comprehensive analysis reveals an extraordinary association between *PTPN22* polymorphism and a defined set of autoimmune diseases, which correlates with specific target tissues, where the corresponding disease manifests. Thus, we propose a new model for the mechanism underlying the association between *PTPN22* and autoimmune disease.

MATERIALS AND METHODS

Identification of eligible studies

An exhaustive search of the Medline database was performed to identify eligible studies (the last search update was August 2011). First, the key word 'PTPN22' was used for the search without any limitations, and retrieved abstracts for 498 articles. Second, we reviewed these 498 abstracts and identified relevant articles that studied the association between the *PTPN22* C1858T polymorphism and autoimmune disease.

Finally, we reviewed the full text of all the relevant articles and identified eligible studies based on following criteria: (a) the study population was European or of European-descent, (b) the study was original and (c) the patients were sporadic cases. We excluded the following: (a) studies containing overlapping data, (b) studies in which the allele frequency could not be ascertained, (c) studies in which family members had been studied and (d) GWAS. Supplementary Table S2 lists all studies used for meta-analysis.

Data extraction

Data extraction was conducted independently by two authors and consensus was achieved regarding the data. The following information was collected from each study: the first author's name, the year of publication, the population of origin, the type of autoimmune disease, the number of cases and controls, the different genotypes of the cases and controls, and the number of T and C alleles in both cases and controls. For studies including subjects from different ethnic groups, only samples from Europeans or those of European descent were extracted. For studies including several independent case–control populations, each case–control population was extracted separately.

Data evaluation and statistical analysis

For the control group in each study, we assessed departures from the Hardy–Weinberg equilibrium using the χ^2 -test. The allele frequencies for the *PTPN22* C1858T polymorphism were determined in each study using the allele-counting method. We then examined the allelic effect of T (minor allele) vs C (major allele). The C allele was considered as the reference allele and the T allele was considered the variant allele. The OR, 95% CI and *P*-values were estimated for each study. Meta-analysis was performed to calculate the pooled OR, 95% CI and *P*-values using the fixed-effects model.¹⁴⁴ Statistical analysis was performed using the Comprehensive Meta-Analysis computer program (Biosta, Englewood, NJ, USA).

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We would like to thank Dr Peter K Gregersen and Dr Stefan Niemann for their critical reading of the manuscript. This work was supported by the start-up packages from the Medical College of Xiamen University and was also supported by Deutsche Forschungsgemeinschaft, Cluster of Excellence 'Inflammation at Interfaces' (EXC 306/1).

REFERENCES

- 1 Fernando MM, Stevens CR, Walsh EC, De Jager PL, Goyette P, Plenge RM *et al*. Defining the role of the MHC in autoimmunity: a review and pooled analysis. *PLoS Genet* 2008; **4**: e1000024.

- 2 Bottini N, Musumeci L, Alonso A, Rahmouni S, Nika K, Rostamkhani M *et al*. A functional variant of lymphoid tyrosine phosphatase is associated with type 1 diabetes. *Nat Genet* 2004; **36**: 337–338.
- 3 Begovich AB, Carlton VE, Honigberg LA, Schrodri SJ, Chokkalingam AP, Alexander HC *et al*. A missense single-nucleotide polymorphism in a gene encoding a protein tyrosine phosphatase (PTPN22) is associated with rheumatoid arthritis. *Am J Hum Genet* 2004; **75**: 330–337.
- 4 Kyogoku C, Langefeld CD, Ortmann WA, Lee A, Selby S, Carlton VE *et al*. Genetic association of the R620W polymorphism of protein tyrosine phosphatase PTPN22 with human SLE. *Am J Hum Genet* 2004; **75**: 504–507.
- 5 Smyth D, Cooper JD, Collins JE, Heward JM, Franklyn JA, Howson JM *et al*. Replication of an association between the lymphoid tyrosine phosphatase locus (LYP/PTPN22) with type 1 diabetes, and evidence for its role as a general autoimmunity locus. *Diabetes* 2004; **53**: 3020–3023.
- 6 Cohen S, Dadi H, Shaoul E, Sharfe N, Roifman CM. Cloning and characterization of a lymphoid-specific, inducible human protein tyrosine phosphatase, Lyp. *Blood* 1999; **93**: 2013–2024.
- 7 Vang T, Congia M, Macis MD, Musumeci L, Orru V, Zavattari P *et al*. Autoimmune-associated lymphoid tyrosine phosphatase is a gain-of-function variant. *Nat Genet* 2005; **37**: 1317–1319.
- 8 Archiga AF, Habib T, He Y, Zhang X, Zhang ZY, Funk A *et al*. Cutting edge: the PTPN22 allelic variant associated with autoimmunity impairs B cell signaling. *J Immunol* 2009; **182**: 3343–3347.
- 9 Douroudis K, Shcherbakova A, Everaus H, Aints A. PTPN22 gene regulates natural killer cell proliferation during *in vitro* expansion. *Tissue Antigens* 2010; **76**: 315–318.
- 10 Hasegawa K, Martin F, Huang G, Tumas D, Diehl L, Chan AC. PEST domain-enriched tyrosine phosphatase (PEP) regulation of effector/memory T cells. *Science* 2004; **303**: 685–689.
- 11 Begovich AB, Caillier SJ, Alexander HC, Penko JM, Hauser SL, Barcellos LF *et al*. The R620W polymorphism of the protein tyrosine phosphatase PTPN22 is not associated with multiple sclerosis. *Am J Hum Genet* 2005; **76**: 184–187.
- 12 Martin MC, Oliver J, Urcelay E, Orozco G, Gomez-Garcia M, Lopez-Nevot MA *et al*. The functional genetic variation in the PTPN22 gene has a negligible effect on the susceptibility to develop inflammatory bowel disease. *Tissue Antigens* 2005; **66**: 314–317.
- 13 Wipff J, Allanore Y, Kahan A, Meyer O, Mouthon L, Guillemin L *et al*. Lack of association between the protein tyrosine phosphatase non-receptor 22 (PTPN22)*620W allele and systemic sclerosis in the French Caucasian population. *Ann Rheum Dis* 2006; **65**: 1230–1232.
- 14 Zhebrun D, Kudryashova Y, Babenko A, Maslyansky A, Kunitskaya N, Popcova D *et al*. Association of PTPN22 1858T/T genotype with type 1 diabetes, Graves' disease but not with rheumatoid arthritis in Russian population. *Aging (Albany NY)* 2011; **3**: 368–373.
- 15 Seldin MF, Shigeta R, Laiho K, Li H, Salla H, Savolainen A *et al*. Finnish case-control and family studies support PTPN22 R620W polymorphism as a risk factor in rheumatoid arthritis, but suggest only minimal or no effect in juvenile idiopathic arthritis. *Genes Immun* 2005; **6**: 720–722.
- 16 Fedetz M, Matesanz F, Caro-Maldonado A, Smirnov II, Chvorostinka VN, Moiseenko TA *et al*. The 1858T PTPN22 gene variant contributes to a genetic risk of type 1 diabetes in a Ukrainian population. *Tissue Antigens* 2006; **67**: 430–433.
- 17 Douroudis K, Prans E, Haller K, Nemvalts V, Rajasalu T, Tillmann V *et al*. Protein tyrosine phosphatase non-receptor type 22 gene variants at position 1858 are associated with type 1 and type 2 diabetes in Estonian population. *Tissue Antigens* 2008; **72**: 425–430.
- 18 Majorczyk E, Jasek M, Ploski R, Wagner M, Kosior A, Pawlik A *et al*. Association of PTPN22 single nucleotide polymorphism with rheumatoid arthritis but not with allergic asthma. *Eur J Hum Genet* 2007; **15**: 1043–1048.
- 19 Ploski R, Dziunycz P, Kostrowski G, Roszkowski PI, Barcz E, Zabek J *et al*. PTPN22/LYP 1858C>T gene polymorphism and susceptibility to endometriosis in a Polish population. *J Reprod Immunol* 2009; **79**: 196–200.
- 20 Skorka A, Bednarczyk T, Bar-Andziak E, Nauman J, Ploski R. Lymphoid tyrosine phosphatase (PTPN22/LYP) variant and Graves' disease in a Polish population: association and gene dose-dependent correlation with age of onset. *Clin Endocrinol (Oxford)* 2005; **62**: 679–682.
- 21 az-Gallo LM, Espino-Paisan L, Franssen K, Gomez-Garcia M, van SS, Cardena C *et al*. Differential association of two PTPN22 coding variants with Crohn's disease and ulcerative colitis. *Inflamm Bowel Dis* 2011; **17**: 2287–2294.
- 22 Kokkonen H, Johansson M, Innala L, Jidell E, Rantapaa-Dahlqvist S. The PTPN22 1858C/T polymorphism is associated with anti-cyclic citrullinated peptide antibody-positive early rheumatoid arthritis in northern Sweden. *Arthritis Res Ther* 2007; **9**: R56.
- 23 Lefvert AK, Zhao Y, Ramanujam R, Yu S, Pirsanen R, Hammarstrom L. PTPN22 R620W promotes production of anti-AChR autoantibodies and IL-2 in myasthenia gravis. *J Neuroimmunol* 2008; **197**: 110–113.
- 24 Rodriguez-Rodriguez L, Taib WR, Topless R, Steer S, Gonzalez-Escribano MF, Balsa A *et al*. The PTPN22 R263Q polymorphism is a risk factor for rheumatoid arthritis in Caucasian case-control samples. *Arthritis Rheum* 2011; **63**: 365–372.
- 25 Skinningsrud B, Husebye ES, Gervin K, Lovas K, Blomhoff A, Wolff AB *et al*. Mutation screening of PTPN22: association of the 1858T-allele with Addison's disease. *Eur J Hum Genet* 2008; **16**: 977–982.
- 26 Viken MK, Olsson M, Flam ST, Forre O, Kvien TK, Thorsby E *et al*. The PTPN22 promoter polymorphism -1123G>C association cannot be distinguished from the 1858C>T association in a Norwegian rheumatoid arthritis material. *Tissue Antigens* 2007; **70**: 190–197.
- 27 Betz RC, Konig K, Flaquer A, Redler S, Eigelshoven S, Kortum AK *et al*. The R620W polymorphism in PTPN22 confers general susceptibility for the development of alopecia areata. *Br J Dermatol* 2008; **158**: 389–391.
- 28 Pierer M, Kaltenhauser S, Arnold S, Wahle M, Baerwald C, Hantzschel H *et al*. Association of PTPN22 1858 single-nucleotide polymorphism with rheumatoid arthritis in a German cohort: higher frequency of the risk allele in male compared to female patients. *Arthritis Res Ther* 2006; **8**: R75.
- 29 Wagenleiter SE, Klein W, Griga T, Schmiegell W, Epplen JT, Jagiello P. A case-control study of tyrosine phosphatase (PTPN22) confirms the lack of association with Crohn's disease. *Int J Immunogenet* 2005; **32**: 323–324.
- 30 Cinek O, Hradsky O, Ahmedov G, Slavcev A, Kolouskova S, Kulich M *et al*. No independent role of the -1123 G>C and +2740 A>G variants in the association of PTPN22 with type 1 diabetes and juvenile idiopathic arthritis in two Caucasian populations. *Diabetes Res Clin Pract* 2007; **76**: 297–303.
- 31 Hradsky O, Lenicek M, Dusatkova P, Bronsky J, Nevala J, Valtrova V *et al*. Variants of CARD15, TNFA and PTPN22 and susceptibility to Crohn's disease in the Czech population: high frequency of the CARD15 1007fs. *Tissue Antigens* 2008; **71**: 538–547.
- 32 Stark K, Rovinsky J, Blazickova S, Grosse-Wilde H, Ferencik S, Hengstenberg C *et al*. Association of common polymorphisms in known susceptibility genes with rheumatoid arthritis in a Slovak population using osteoarthritis patients as controls. *Arthritis Res Ther* 2009; **11**: R70.
- 33 Nielsen C, Hansen D, Husby S, Lillevang ST. Sex-specific association of the human PTPN22 1858T-allele with type 1 diabetes. *Int J Immunogenet* 2007; **34**: 469–473.
- 34 Canton I, Akhtar S, Galvalas NG, Gawkrödger DJ, Blomhoff A, Watson PF *et al*. A single-nucleotide polymorphism in the gene encoding lymphoid protein tyrosine phosphatase (PTPN22) confers susceptibility to generalised vitiligo. *Genes Immun* 2005; **6**: 584–587.
- 35 Hinks A, Eyre S, Barton A, Thomson W, Worthington J. Investigation of genetic variation across the protein tyrosine phosphatase gene in patients with rheumatoid arthritis in the UK. *Ann Rheum Dis* 2007; **66**: 683–686.
- 36 Kemp EH, McDonagh AJ, Wengraf DA, Messenger AG, Gawkrödger DJ, Cork MJ *et al*. The non-synonymous C1858T substitution in the PTPN22 gene is associated with susceptibility to the severe forms of alopecia areata. *Hum Immunol* 2006; **67**: 535–539.
- 37 Prescott NJ, Fisher SA, Onnie C, Pattni R, Steer S, Sanderson J *et al*. A general autoimmunity gene (PTPN22) is not associated with inflammatory bowel disease in a British population. *Tissue Antigens* 2005; **66**: 318–320.
- 38 Smyth DJ, Plagnol V, Walker NM, Cooper JD, Downes K, Yang JH *et al*. Shared and distinct genetic variants in type 1 diabetes and celiac disease. *N Engl J Med* 2008; **359**: 2767–2777.
- 39 Steer S, Lad B, Grumley JA, Kingsley GH, Fisher SA. Association of R602W in a protein tyrosine phosphatase gene with a high risk of rheumatoid arthritis in a British population: evidence for an early onset/disease severity effect. *Arthritis Rheum* 2005; **52**: 358–360.
- 40 Wesoly J, van der Helm-van Mil AH, Toes RE, Chokkalingam AP, Carlton VE, Begovich AB *et al*. Association of the PTPN22 C1858T single-nucleotide polymorphism with rheumatoid arthritis phenotypes in an inception cohort. *Arthritis Rheum* 2005; **52**: 2948–2950.
- 41 Wesoly J, Hu X, Thabet MM, Chang M, Uh H, Allaart CF *et al*. The 620W allele is the PTPN22 genetic variant conferring susceptibility to RA in a Dutch population. *Rheumatology (Oxford)* 2007; **46**: 617–621.
- 42 Zhernakova A, Eerligh P, Wijmenga C, Barrera P, Roep BO, Koeleman BP. Differential association of the PTPN22 coding variant with autoimmune diseases in a Dutch population. *Genes Immun* 2005; **6**: 459–461.
- 43 Pazar B, Gergely Jr P, Nagy ZB, Gombos T, Pozsonyi E, Rajczyk K *et al*. Role of HLA-DRB1 and PTPN22 genes in susceptibility to juvenile idiopathic arthritis in Hungarian patients. *Clin Exp Rheumatol* 2008; **26**: 1146–1152.
- 44 Ittah M, Gottenberg JE, Proust A, Hachulla E, Puechal X, Loeiseau P *et al*. No evidence for association between 1858 C/T single-nucleotide polymorphism of PTPN22 gene and primary Sjogren's syndrome. *Genes Immun* 2005; **6**: 457–458.
- 45 Vandiedonck C, Capdevielle C, Giraud M, Krumeich S, Jais JP, Eymard B *et al*. Association of the PTPN22*R620W polymorphism with autoimmune myasthenia gravis. *Ann Neurol* 2006; **59**: 404–407.

- 46 Balada E, Simeon-Aznar CP, Serrano-Acedo S, Martinez-Lostao L, Selva-O'Callaghan A, Fonollosa-Pla V *et al*. Lack of association of the PTPN22 gene polymorphism R620W with systemic sclerosis. *Clin Exp Rheumatol* 2006; **24**: 321–324.
- 47 Rueda B, Nunez C, Orozco G, Lopez-Nevot MA, de la Concha EG, Martin J *et al*. C1858T functional variant of PTPN22 gene is not associated with celiac disease genetic predisposition. *Hum Immunol* 2005; **66**: 848–852.
- 48 Santin I, Castellanos-Rubio A, Aransay AM, Castano L, Vitoria JC, Bilbao JR. The functional R620W variant of the PTPN22 gene is associated with celiac disease. *Tissue Antigens* 2008; **71**: 247–249.
- 49 Laberge GS, Birlea SA, Fain PR, Spritz RA. The PTPN22-1858C>T (R620W) functional polymorphism is associated with generalized vitiligo in the Romanian population. *Pigment Cell Melanoma Res* 2008; **21**: 206–208.
- 50 Latiano A, Palmieri O, Valvano MR, Bossa F, Latiano T, Corritore G *et al*. Evaluating the role of the genetic variations of PTPN22, NFKB1, and FcGR1IA genes in inflammatory bowel disease: a meta-analysis. *Inflamm Bowel Dis* 2007; **13**: 1212–1219.
- 51 Petrone A, Suraci C, Capizzi M, Giaccari A, Bosi E, Tiberti C *et al*. The protein tyrosine phosphatase nonreceptor 22 (PTPN22) is associated with high GAD antibody titer in latent autoimmune diabetes in adults: Non Insulin Requiring Autoimmune Diabetes (NIRAD) Study 3. *Diabetes Care* 2008; **31**: 534–538.
- 52 Saccucci P, Del DE, Rapini N, Verrotti A, Piccinini S, Maccari A *et al*. Association between PTPN22 C1858T and type 1 diabetes: a replication in continental Italy. *Tissue Antigens* 2008; **71**: 234–237.
- 53 Saccucci P, Banci M, Cozzoli E, Neri A, Magrini A, Bottini E *et al*. Atherosclerosis and PTPN22: a study in coronary artery disease. *Cardiology* 2011; **119**: 54–56.
- 54 Eliopoulos E, Zervou MI, Andreou A, Dimopoulou K, Cosmidis N, Voloudakis G *et al*. Association of the PTPN22 R620W polymorphism with increased risk for SLE in the genetically homogeneous population of Crete. *Lupus* 2011; **20**: 501–506.
- 55 Morgan AR, Han DY, Huebner C, Lam WJ, Fraser AG, Ferguson LR. PTPN22 but not PTPN22 is associated with Crohn's disease in a New Zealand population. *Tissue Antigens* 2010; **76**: 119–125.
- 56 Butt C, Peddle L, Greenwood C, Hamilton S, Gladman D, Rahman P. Association of functional variants of PTPN22 and tp53 in psoriatic arthritis: a case-control study. *Arthritis Res Ther* 2006; **8**: R27.
- 57 Chagastelles PC, Romitti M, Trein MR, Bandinelli E, Tschiedel B, Nardi NB. Association between the 1858T allele of the protein tyrosine phosphatase nonreceptor type 22 and type 1 diabetes in a Brazilian population. *Tissue Antigens* 2010; **76**: 144–148.
- 58 De Jager PL, Sawcer S, Waliszewska A, Farwell L, Wild G, Cohen A *et al*. Evaluating the role of the 620W allele of protein tyrosine phosphatase PTPN22 in Crohn's disease and multiple sclerosis. *Eur J Hum Genet* 2006; **14**: 317–321.
- 59 Gomes FM, Bianco B, Teles JS, Christofolini DM, de Souza AM, Guedes AD *et al*. PTPN22 C1858T polymorphism in women with endometriosis. *Am J Reprod Immunol* 2010; **63**: 227–232.
- 60 Gomez LM, Anaya JM, Gonzalez CI, Pineda-Tamayo R, Otero W, Arango A *et al*. PTPN22 C1858T polymorphism in Colombian patients with autoimmune diseases. *Genes Immun* 2005; **6**: 628–631.
- 61 Gourh P, Tan FK, Assassi S, Ahn CW, McNearney TA, Fischbach M *et al*. Association of the PTPN22 R620W polymorphism with anti-topoisomerase I- and anticentromere antibody-positive systemic sclerosis. *Arthritis Rheum* 2006; **54**: 3945–3953.
- 62 Gourh P, Agarwal SK, Martin E, Divecha D, Rueda B, Bunting H *et al*. Association of the C8orf13-BLK region with systemic sclerosis in North-American and European populations. *J Autoimmun* 2010; **34**: 155–162.
- 63 Reddy MV, Wang H, Liu S, Bode B, Reed JC, Steed RD *et al*. Association between type 1 diabetes and GWAS SNPs in the southeast US Caucasian population. *Genes Immun* 2011; **12**: 208–212.
- 64 Sachdev A, Bhanusali DG, Patterson KC, Zamora MB, Ghuman A, Gerlach JA *et al*. PTPN22 1858T is not a risk factor for North American pemphigus vulgaris. *Exp Dermatol* 2011; **20**: 514–519.
- 65 Steck AK, Liu SY, McFann K, Barriga KJ, Babu SR, Eisenbarth GS *et al*. Association of the PTPN22/LYP gene with type 1 diabetes. *Pediatr Diabetes* 2006; **7**: 274–278.
- 66 Thompson SD, Sudman M, Ramos PS, Marion MC, Ryan M, Tsoras M *et al*. The susceptibility loci juvenile idiopathic arthritis shares with other autoimmune diseases extend to PTPN22, COG6, and ANGPT1. *Arthritis Rheum* 2010; **62**: 3265–3276.
- 67 van OM, Wintle RF, Liu X, Yazdanpanah M, Gu X, Newman B *et al*. Association of the lymphoid tyrosine phosphatase R620W variant with rheumatoid arthritis, but not Crohn's disease, in Canadian populations. *Arthritis Rheum* 2005; **52**: 1993–1998.
- 68 Baca V, Velazquez-Cruz R, Salas-Martinez G, Espinosa-Rosales F, Saldana-Alvarez Y, Orozco L. Association analysis of the PTPN22 gene in childhood-onset systemic lupus erythematosus in Mexican population. *Genes Immun* 2006; **7**: 693–695.
- 69 Alkhateeb A, Qarqaz F, Al-Sabah J, Al RT. Clinical characteristics and PTPN22 1858C/T variant analysis in Jordanian Arab vitiligo patients. *Mol Diagn Ther* 2010; **14**: 179–184.
- 70 Anis SK, bdel Ghany EA, Mostafa NO, Ali AA. The role of PTPN22 gene polymorphism in childhood immune thrombocytopenic purpura. *Blood Coagul Fibrinolysis* 2011; **22**: 521–525.
- 71 Ates A, Karaaslan Y, Karatayli E, Ertugrul E, Aksaray S, Turkiylmaz A *et al*. Association of the PTPN22 gene polymorphism with autoantibody positivity in Turkish rheumatoid arthritis patients. *Tissue Antigens* 2011; **78**: 56–59.
- 72 Chabchoub G, Teixeira EP, Maalej A, Ben HM, Bahloul Z, Cornelis F *et al*. The R620W polymorphism of the protein tyrosine phosphatase 22 gene in autoimmune thyroid diseases and rheumatoid arthritis in the Tunisian population. *Ann Hum Biol* 2009; **36**: 342–349.
- 73 Lamsyah H, Rueda B, Baassi L, Elaoud R, Bottini N, Sadki K *et al*. Association of PTPN22 gene functional variants with development of pulmonary tuberculosis in Moroccan population. *Tissue Antigens* 2009; **74**: 228–232.
- 74 Sahin N, Gunduz F, Inanc N, Direskeneli H, Saruhan-Direskeneli G. No association of PTPN22 gene polymorphism with rheumatoid arthritis in Turkey. *Rheumatol Int* 2009; **30**: 81–83.
- 75 Sfar I, Ben AW, Mouelhi L, Aouadi H, Ben RT, Makhoul M *et al*. Lymphoid tyrosine phosphatase R620W variant and inflammatory bowel disease in Tunisia. *World J Gastroenterol* 2010; **16**: 479–483.
- 76 Tikly M, Govind N, Frost J, Ramsay M. The PTPN22 R620W polymorphism is not associated with systemic rheumatic diseases in South Africans. *Rheumatology (Oxford)* 2010; **49**: 820–821.
- 77 Ikari K, Momohara S, Inoue E, Tomatsu T, Hara M, Yamanaka H *et al*. Haplotype analysis revealed no association between the PTPN22 gene and RA in a Japanese population. *Rheumatology (Oxford)* 2006; **45**: 1345–1348.
- 78 Kawasaki E, Awata T, Ikegami H, Kobayashi T, Maruyama T, Nakanishi K *et al*. Systematic search for single nucleotide polymorphisms in a lymphoid tyrosine phosphatase gene (PTPN22): association between a promoter polymorphism and type 1 diabetes in Asian populations. *Am J Med Genet A* 2006; **140**: 586–593.
- 79 Lee HS, Korman BD, Le JM, Kastner DL, Remmers EF, Gregersen PK *et al*. Genetic risk factors for rheumatoid arthritis differ in Caucasian and Korean populations. *Arthritis Rheum* 2009; **60**: 364–371.
- 80 Pemberton TJ, Mehta NU, Witonsky D, Di RA, Allayee H, Conti DV *et al*. Prevalence of common disease-associated variants in Asian Indians. *BMC Genet* 2008; **9**: 13.
- 81 Zhang ZH, Chen F, Zhang XL, Jin Y, Bai J, Fu SB. PTPN22 allele polymorphisms in 15 Chinese populations. *Int J Immunogenet* 2008; **35**: 433–437.
- 82 Gregersen PK, Olson LM. Recent advances in the genetics of autoimmune disease. *Annu Rev Immunol* 2009; **27**: 363–391.
- 83 Laddha NC, Dwivedi M, Shajil EM, Prajapati H, Marfatia YS, Begum R. Association of PTPN22 1858C/T polymorphism with vitiligo susceptibility in Gujarat population. *J Dermatol Sci* 2008; **49**: 260–262.
- 84 Sfar I, Dhaoui T, Habibi I, Abdelmoula L, Makhoul M, Ben RT *et al*. Functional polymorphisms of PTPN22 and FcγR genes in Tunisian patients with rheumatoid arthritis. *Arch Inst Pasteur Tunis* 2009; **86**: 51–62.
- 85 Dultz G, Matheis N, Dittmar M, Rohrig B, Bender K, Kahaly GJ. The protein tyrosine phosphatase non-receptor type 22 C1858T polymorphism is a joint susceptibility locus for immunthyroiditis and autoimmune diabetes. *Thyroid* 2009; **19**: 143–148.
- 86 Fichna M, Zurawek M, Januszkievicz-Lewandowska D, Fichna P, Nowak J. PTPN22, PDCD1 and CYP27B1 polymorphisms and susceptibility to type 1 diabetes in Polish patients. *Int J Immunogenet* 2010; **37**: 367–372.
- 87 Kahles H, Ramos-Lopez E, Lange B, Zwermann O, Reincke M, Badenhoop K. Sex-specific association of PTPN22 1858T with type 1 diabetes but not with Hashimoto's thyroiditis or Addison's disease in the German population. *Eur J Endocrinol* 2005; **153**: 895–899.
- 88 Kordonouri O, Hartmann R, Charpentier N, Knip M, Danne T, Ilonen J. Genetic risk markers related to diabetes-associated autoantibodies in young patients with type 1 diabetes in Berlin, Germany. *Exp Clin Endocrinol Diabetes* 2010; **118**: 245–249.
- 89 Korolija M, Renar IP, Hadzija M, Medvidovic EP, Pavkovic P, Jokic M *et al*. Association of PTPN22 C1858T and CTLA-4 A49G polymorphisms with type 1 diabetes in Croats. *Diabetes Res Clin Pract* 2009; **86**: e54–e57.
- 90 Maziarz M, Janer M, Roach JC, Hagopian W, Palmer JP, Deutsch K *et al*. The association between the PTPN22 1858C>T variant and type 1 diabetes depends on HLA risk and GAD65 autoantibodies. *Genes Immun* 2010; **11**: 406–415.
- 91 Santiago JL, Martinez A, de la CH, Fernandez-Arguero M, Figueredo MA, de la Concha EG *et al*. Susceptibility to type 1 diabetes conferred by the PTPN22 C1858T polymorphism in the Spanish population. *BMC Med Genet* 2007; **8**: 54.
- 92 Coenen MJ, Trynka G, Heskamp S, Franke B, van Diemen CC, Smolonska J *et al*. Common and different genetic background for rheumatoid arthritis and coeliac disease. *Hum Mol Genet* 2009; **18**: 4195–4203.

- 93 Dieude P, Teixeira VH, Pierlot C, Cornelis F, Petit-Teixeira E. Testing for linkage and association with rheumatoid arthritis a ptpn22 promoter polymorphism reported to be associated and linked with type 1 diabetes in the Caucasian population. *Ann Rheum Dis* 2008; **67**: 900–901.
- 94 Farago B, Talian GC, Komlosi K, Nagy G, Berki T, Gyetvai A *et al*. Protein tyrosine phosphatase gene C1858T allele confers risk for rheumatoid arthritis in Hungarian subjects. *Rheumatol Int* 2009; **29**: 793–796.
- 95 Harrison P, Pointon JJ, Farrar C, Brown MA, Wordsworth BP. Effects of PTPN22 C1858T polymorphism on susceptibility and clinical characteristics of British Caucasian rheumatoid arthritis patients. *Rheumatology (Oxford)* 2006; **45**: 1009–1011.
- 96 Hinks A, Barton A, John S, Bruce I, Hawkins C, Griffiths CE *et al*. Association between the PTPN22 gene and rheumatoid arthritis and juvenile idiopathic arthritis in a UK population: further support that PTPN22 is an autoimmunity gene. *Arthritis Rheum* 2005; **52**: 1694–1699.
- 97 Lee AT, Li W, Liew A, Bombardier C, Weisman M, Massarotti EM *et al*. The PTPN22 R620W polymorphism associates with RF positive rheumatoid arthritis in a dose-dependent manner but not with HLA-SE status. *Genes Immun* 2005; **6**: 129–133.
- 98 Majorczyk E, Pawlik A, Kusnierczyk P. PTPN22 1858C>T polymorphism is strongly associated with rheumatoid arthritis but not with a response to methotrexate therapy. *Int Immunopharmacol* 2010; **10**: 1626–1629.
- 99 Morgan AW, Robinson JI, Conaghan PG, Martin SG, Hensor EM, Morgan MD *et al*. Evaluation of the rheumatoid arthritis susceptibility loci HLA-DRB1, PTPN22, OLIG3/TNFAIP3, STAT4 and TRAF1/C5 in an inception cohort. *Arthritis Res Ther* 2010; **12**: R57.
- 100 Newman WG, Zhang Q, Liu X, Walker E, Ternan H, Owen J *et al*. Rheumatoid arthritis association with the FCRL3 -169C polymorphism is restricted to PTPN22 1858T-homozygous individuals in a Canadian population. *Arthritis Rheum* 2006; **54**: 3820–3827.
- 101 Plenge RM, Padyukov L, Remmers EF, Purcell S, Lee AT, Karlson EW *et al*. Replication of putative candidate-gene associations with rheumatoid arthritis in >4,000 samples from North America and Sweden: association of susceptibility with PTPN22, CTLA4, and PADI4. *Am J Hum Genet* 2005; **77**: 1044–1060.
- 102 Simkins HM, Merriman ME, Highton J, Chapman PT, O'Donnell JL, Jones PB *et al*. Association of the PTPN22 locus with rheumatoid arthritis in a New Zealand Caucasian cohort. *Arthritis Rheum* 2005; **52**: 2222–2225.
- 103 Viken MK, Amundsen SS, Kvien TK, Boberg KM, Gilboe IM, Lilleby V *et al*. Association analysis of the 1858C>T polymorphism in the PTPN22 gene in juvenile idiopathic arthritis and other autoimmune diseases. *Genes Immun* 2005; **6**: 271–273.
- 104 Aksoy R, Duman T, Keskin O, Duzgun N. No association of PTPN22 R620 W gene polymorphism with rheumatic heart disease and systemic lupus erythematosus. *Mol Biol Rep* 2011; **38**: 5393–5396.
- 105 Kaufman KM, Kelly JA, Herring BJ, Adler AJ, Glenn SB, Namjou B *et al*. Evaluation of the genetic association of the PTPN22 R620W polymorphism in familial and sporadic systemic lupus erythematosus. *Arthritis Rheum* 2006; **54**: 2533–2540.
- 106 Orozco G, Sanchez E, Gonzalez-Gay MA, Lopez-Nevot MA, Torres B, Caliz R *et al*. Association of a functional single-nucleotide polymorphism of PTPN22, encoding lymphoid protein phosphatase, with rheumatoid arthritis and systemic lupus erythematosus. *Arthritis Rheum* 2005; **52**: 219–224.
- 107 Piotrowski P, Lianeri M, Wudarski M, Lacki JK, Jagodzinski PP. Contribution of the R620W polymorphism of protein tyrosine phosphatase non-receptor 22 to systemic lupus erythematosus in Poland. *Clin Exp Rheumatol* 2008; **26**: 1099–1102.
- 108 Reddy MV, Johansson M, Sturfelt G, Jonsen A, Gunnarsson I, Svenungsson E *et al*. The R620W C/T polymorphism of the gene PTPN22 is associated with SLE independently of the association of PDCD1. *Genes Immun* 2005; **6**: 658–662.
- 109 Velaga MR, Wilson V, Jennings CE, Owen CJ, Herington S, Donaldson PT *et al*. The codon 620 tryptophan allele of the lymphoid tyrosine phosphatase (LYP) gene is a major determinant of Graves' disease. *J Clin Endocrinol Metab* 2004; **89**: 5862–5865.
- 110 Roycroft M, Fichna M, McDonald D, Owen K, Zurawek M, Gryczynska M *et al*. The tryptophan 620 allele of the lymphoid tyrosine phosphatase (PTPN22) gene predisposes to autoimmune Addison's disease. *Clin Endocrinol (Oxford)* 2009; **70**: 358–362.
- 111 Carr EJ, Niederer HA, Williams J, Harper L, Watts RA, Lyons PA *et al*. Confirmation of the genetic association of CTLA4 and PTPN22 with ANCA-associated vasculitis. *BMC Med Genet* 2009; **10**: 121.
- 112 Jagiello P, Aries P, Arning L, Wagenleiter SE, Csernok E, Hellmich B *et al*. The PTPN22 620W allele is a risk factor for Wegener's granulomatosis. *Arthritis Rheum* 2005; **52**: 4039–4043.
- 113 Hinks A, Eyre S, Ke X, Barton A, Martin P, Flynn E *et al*. Overlap of disease susceptibility loci for rheumatoid arthritis and juvenile idiopathic arthritis. *Ann Rheum Dis* 2010; **69**: 1049–1053.
- 114 Jin Y, Birlea SA, Fain PR, Gowan K, Riccardi SL, Holland PJ *et al*. Variant of TYR and autoimmunity susceptibility loci in generalized vitiligo. *N Engl J Med* 2010; **362**: 1686–1697.
- 115 Chuang WY, Strobel P, Belharazem D, Rieckmann P, Toyka KV, Nix W *et al*. The PTPN22 gain-of-function +1858T(+) genotypes correlate with low IL-2 expression in thymomas and predispose to myasthenia gravis. *Genes Immun* 2009; **10**: 667–672.
- 116 Greve B, Hoffmann P, Illes Z, Rozsa C, Berger K, Weissert R *et al*. The autoimmunity-related polymorphism PTPN22 1858C/T is associated with anti-titin antibody-positive myasthenia gravis. *Hum Immunol* 2009; **70**: 540–542.
- 117 Anderson CA, Massey DC, Barrett JC, Prescott NJ, Tremelling M, Fisher SA *et al*. Investigation of Crohn's disease risk loci in ulcerative colitis further defines their molecular relationship. *Gastroenterology* 2009; **136**: 523–529.
- 118 az-Gallo LM, Gourh P, Broen J, Simeon C, Fonollosa V, Ortego-Centeno N *et al*. Analysis of the influence of PTPN22 gene polymorphisms in systemic sclerosis. *Ann Rheum Dis* 2011; **70**: 454–462.
- 119 Radstake TR, Gorlova O, Rueda B, Martin JE, Alizadeh BZ, Palomino-Morales R *et al*. Genome-wide association study of systemic sclerosis identifies CD247 as a new susceptibility locus. *Nat Genet* 2010; **42**: 426–429.
- 120 Sawcer S, Henthall G, Pirinen M, Spencer CC, Patsopoulos NA, Moutsianas L *et al*. Genetic risk and a primary role for cell-mediated immune mechanisms in multiple sclerosis. *Nature* 2011; **476**: 214–219.
- 121 Matesanz F, Rueda B, Orozco G, Fernandez O, Leyva L, Alcina A *et al*. Protein tyrosine phosphatase gene (PTPN22) polymorphism in multiple sclerosis. *J Neurol* 2005; **252**: 994–995.
- 122 Huffmeier U, Reis A, Steffens M, Lascorz J, Bohm B, Lohmann J *et al*. Male restricted genetic association of variant R620W in PTPN22 with psoriatic arthritis. *J Invest Dermatol* 2006; **126**: 932–935.
- 123 Smith RL, Warren RB, Eyre S, Ke X, Young HS, Allen M *et al*. Polymorphisms in the PTPN22 region are associated with psoriasis of early onset. *Br J Dermatol* 2008; **158**: 962–968.
- 124 Stuart PE, Nair RP, Ellinghaus E, Ding J, Tejasvi T, Gudjonsson JE *et al*. Genome-wide association analysis identifies three psoriasis susceptibility loci. *Nat Genet* 2010; **42**: 1000–1004.
- 125 Juneblad K, Johansson M, Rantapaa-Dahlqvist S, Alenius GM. Association between the PTPN22 +1858 C/T polymorphism and psoriatic arthritis. *Arthritis Res Ther* 2011; **13**: R45.
- 126 Petukhova L, Duvic M, Hordinsky M, Norris D, Price V, Shimomura Y *et al*. Genome-wide association study in alopecia areata implicates both innate and adaptive immunity. *Nature* 2010; **466**: 113–117.
- 127 Chinoy H, Platt H, Lamb JA, Betteridge Z, Gunawardena H, Fertig N *et al*. The protein tyrosine phosphatase N22 gene is associated with juvenile and adult idiopathic inflammatory myopathy independent of the HLA 8.1 haplotype in British Caucasian patients. *Arthritis Rheum* 2008; **58**: 3247–3254.
- 128 D'Silva KJ, Zamora MB, Gerlach J, Schwartz KA. Increased representation of the PTPN22 mutation in patients with immune thrombocytopenia. *J Thromb Haemost* 2010; **8**: 2076–2078.
- 129 Martin TM, Bye L, Modi N, Stanford MR, Vaughan R, Smith JR *et al*. Genotype analysis of polymorphisms in autoimmune susceptibility genes, CTLA-4 and PTPN22, in an acute anterior uveitis cohort. *Mol Vis* 2009; **15**: 208–212.
- 130 Milkiewicz P, Pache I, Buwaneswaran H, Liu X, Coltescu C, Heathcote EJ *et al*. The PTPN22 1858T variant is not associated with primary biliary cirrhosis. *Tissue Antigens* 2006; **67**: 434–437.
- 131 Orozco G, Garcia-Porrúa C, Lopez-Nevot MA, Raya E, Gonzalez-Gay MA, Martin J. Lack of association between ankylosing spondylitis and a functional polymorphism of PTPN22 proposed as a general susceptibility marker for autoimmunity. *Ann Rheum Dis* 2006; **65**: 687–688.
- 132 Reveille JD, Sims AM, Danoy P, Evans DM, Leo P, Pointon JJ *et al*. Genome-wide association study of ankylosing spondylitis identifies non-MHC susceptibility loci. *Nat Genet* 2010; **42**: 123–127.
- 133 Hirschfield GM, Liu X, Xu C, Lu Y, Xie G, Lu Y *et al*. Primary biliary cirrhosis associated with HLA, IL12A, and IL12RB2 variants. *N Engl J Med* 2009; **360**: 2544–2555.
- 134 Charatcharoenwitthaya P, Lindor KD. Primary sclerosing cholangitis: diagnosis and management. *Curr Gastroenterol Rep* 2006; **8**: 75–82.
- 135 Amagai M. Pemphigus vulgaris and its active disease mouse model. *Curr Dir Autoimmun* 2008; **10**: 167–181.
- 136 Meiri K, Kallel-Sellami M, Petit-Teixeira E, Abida O, Mbarek H, Zitouni M *et al*. PTPN22 R620W polymorphism is not associated with pemphigus. *Br J Dermatol* 2007; **157**: 1068–1069.
- 137 Vang T, Miletic AV, Bottini N, Mustelin T. Protein tyrosine phosphatase PTPN22 in human autoimmunity. *Autoimmunity* 2007; **40**: 453–461.
- 138 Menard L, Saadoun D, Isnardi I, Ng YS, Meyers G, Massad C *et al*. The PTPN22 allele encoding an R620W variant interferes with the removal of developing autoreactive B cells in humans. *J Clin Invest* 2011; **121**: 3635–3644.
- 139 Zhang J, Zahir N, Jiang Q, Miliotis H, Heyraud S, Meng X *et al*. The autoimmune disease-associated PTPN22 variant promotes calpain-mediated Lyp/Pep degradation associated with lymphocyte and dendritic cell hyperresponsiveness. *Nat Genet* 2011; **43**: 902–907.

- 140 Hugot JP, Chamaillard M, Zouali H, Lesage S, Cezard JP, Belaiche J *et al*. Association of NOD2 leucine-rich repeat variants with susceptibility to Crohn's disease. *Nature* 2001; **411**: 599–603.
- 141 Ogura Y, Bonen DK, Inohara N, Nicolae DL, Chen FF, Ramos R *et al*. A frameshift mutation in NOD2 associated with susceptibility to Crohn's disease. *Nature* 2001; **411**: 603–606.
- 142 Anderson AC, Nicholson LB, Legge KL, Turchin V, Zaghoulani H, Kuchroo VK. High frequency of autoreactive myelin proteolipid protein-specific T cells in the periphery of naive mice: mechanisms of selection of the self-reactive repertoire. *J Exp Med* 2000; **191**: 761–770.
- 143 Bruno R, Sabater L, Sospedra M, Ferrer-Francesch X, Escudero D, Martinez-Caceres E *et al*. Multiple sclerosis candidate autoantigens except myelin oligodendrocyte glycoprotein are transcribed in human thymus. *Eur J Immunol* 2002; **32**: 2737–2747.
- 144 Mantel N, Haenszel W. Statistical aspects of the analysis of data from retrospective studies of disease. *J Natl Cancer Inst* 1959; **22**: 719–748.

Supplementary Information accompanies the paper on Genes and Immunity website (<http://www.nature.com/gene>).