

NITRIC OXIDE SYNTHASE INHIBITORS UNMASK ACETYLCHOLINE-MEDIATED CONSTRICTION OF CEREBRAL VESSELS IN THE *IN VITRO* ISOLATED GUINEA-PIG BRAIN

L. LIBRIZZI,* G. FOLCO† and M. DE CURTIS*‡

*Dipartimento di Neurofisiologia Sperimentale, Istituto Nazionale Neurologico, via Celoria 11, 20133 Milan, Italy

†Centro di Farmacologia Cardiopolmonare Sperimentale, Istituto di Scienze Farmacologiche, via Balzaretti 9, Milan, Italy

Abstract—The control of arterial vascular tone by acetylcholine contributes to the regulation of cerebral blood flow. We analysed the effects of intraluminal application of acetylcholine (1 μ M) on the cerebral vascular tone by measuring changes in resistance to perfusion pressure in an isolated guinea-pig brain preparation maintained *in vitro* by arterial perfusion under constant flow. Acetylcholine induced a reproducible, fast-onset dilation that was prevented by the nitric oxide scavenger Methylene Blue (10 μ M) and by the muscarinic receptor antagonist atropine (0.1 μ M). Prolonged arterial perfusion with the nitric oxide synthase inhibitors *N*-nitro-L-arginine (1 mM) and *N*-nitro-L-arginine methyl ester (30–100 μ M) induced a slowly developing increase of 25.9 ± 13.44 mmHg in vascular tone and blocked the acetylcholine-induced vasodilation. In these experimental conditions, the dilation determined by the nitric oxide donor nitroprusside (0.1 μ M) was unaffected. In five experiments, the blockade of dilation unmasked a slow acetylcholine-mediated vasoconstriction (14.40 ± 3.85 mmHg) that was antagonized by atropine.

The results demonstrate that acetylcholine exerts two simultaneous and opposite effects on guinea-pig cerebral vessels, characterized by a slow direct constriction concealed in physiological conditions by a fast vasodilation mediated through the release of nitric oxide by endothelial cells. Acetylcholine-mediated increase in vascular tone may play a role in aggravating cerebral perfusion when endothelial cell damage occurs during brain ischemia. © 2000 IBRO. Published by Elsevier Science Ltd. All rights reserved.

Key words: acetylcholine, guinea-pig, isolated brain preparation, nitric oxide, nitric oxide synthase.

Acetylcholine (ACh) contributes to the regulation of blood flow by promoting changes in vascular resistance. Several studies demonstrated that ACh increases cerebral blood flow^{23,32,38} by inducing vasodilation of intracerebral arterioles¹² and pial arterioles⁹ through the stimulation of nitric oxide (NO) release from endothelial cells.^{3,20} The increase in cerebral blood flow mediated by ACh is indeed abolished by endothelial cell damage³⁹ and is prevented by nitric oxide synthase (NOS) inhibitors^{26–29} that induce *per se* an increase in vascular tone.^{20,40}

Cholinergic regulation of cerebrovascular activity is operated by local cholinergic interneurons¹¹ and via cholinergic fibers that originate from neurons in the basal forebrain and terminate either on the basal lamina of arterioles and capillaries^{4,5,42} or on NOS-immunoreactive neurons.⁴⁴ As for the direct effect of ACh applied on brain vessels, the increase in cortical blood flow induced by stimulation of the cholinergic basal forebrain nuclei^{8,13,29,43} has been shown to be mediated by NO release from both NOS-positive neurons and endothelial cells.^{1,35,44,47}

We demonstrate that, in the intact vasculature of the isolated guinea-pig brain preparation,^{14,16,34} a steady-state, NO-independent mechanism controls vessel tone; after prolonged exposure to NOS inhibitors, a slow, persistent increase in tone takes place, ACh-mediated vasodilation is abolished and a contraction due to muscarinic receptor

activation is unmasked. Preliminary results have been published in abstract form.³⁰

EXPERIMENTAL PROCEDURES

The experiments were carried out on brains isolated from young adult guinea-pigs (150–200 g) maintained *in vitro* by arterial perfusion. The procedure for brain isolation has been described extensively elsewhere.^{14,16,31,34} In brief, during anesthesia with pentobarbital (20 mg/kg, i.p.), the brain was extracted under hypothermic conditions and transferred to an incubation chamber, where a micro-cannula was inserted in the basilar artery to ensure arterial perfusion with a complex saline solution (composition, in mM: NaCl, 126; KCl, 3; KH₂PO₄, 1.2; MgSO₄, 1.3; CaCl₂, 2.4; NaHCO₃, 26; glucose, 15; HEPES, 2.1; 3% dextran, mol. wt 70,000) oxygenated with a 95% O₂–5% CO₂ gas mixture. Arterial perfusion at 5.5 ml/min was performed via a peristaltic pump. Experiments were performed at 32°C. All efforts were made to minimize the suffering and the number of animals used. The experimental procedure was approved by the Ethical Committee of the Istituto Nazionale Neurologico in accordance with the regulations on animal research. A pressure transducer (Model 800, Bentley) was inserted between the peristaltic pump and the micro-cannula to monitor the arterial pressure. The changes in vascular tone were measured as increases or decreases in the resistance to flow opposed by the entire cerebral arterial system. The voltage signal recorded at the pressure transducer was amplified and digitized via a National AD board (AT-MIO32) for off-line analysis with Labview software developed in our laboratory by Dr Gerardo Biella. The isolated guinea-pig brain has been utilized in the last decade for electrophysiological studies of brain activity.^{7,17} More recently, we demonstrated that in this preparation the vascular system is functionally and structurally preserved.^{15,16}

Drugs diluted in the perfusate were applied intraluminally at the following concentrations: *N*-nitro-L-arginine methyl ester (L-NAME), 30–100 μ M; *N*-nitro-L-arginine (L-NNA), 1 mM; ACh, 1 μ M; nitroprusside (NP), 0.1 μ M; U46619, 30–100 nM; atropine, 0.1 μ M; Methylene Blue, 10 μ M. With the exception of U46619 (provided by Cayman Chemical Company, USA), all drugs were obtained from Sigma. The results are expressed as mean values \pm S.D. of *n* replications.

‡To whom correspondence should be addressed. Tel.: +39-02-2394280; fax: +39-02-70600775.

E-mail address: decurtis@istituto-besta.it (M. de Curtis).

Abbreviations: ACh, acetylcholine; HEPES, *N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulfonic acid; L-NAME, *N*-nitro-L-arginine methyl ester; L-NNA, *N*-nitro-L-arginine; NO, nitric oxide; NOS, nitric oxide synthase; NP, nitroprusside.




Fig. 1. Intraluminal application of ACh induces a muscarinic receptor-dependent vasodilation. A rapid vasodilation is determined by arterial perfusion of 1 μ M ACh during the constriction exerted by U46619 (0.1 μ M; A). (B) The ACh-induced dilation is prevented by co-perfusion with atropine (0.1 μ M; B). The effect of atropine is reversible (C). Upward and downward deflections correspond to constriction and dilation of the cerebral vessels, respectively. In this and in the following figures, the timing of the drug perfusions is indicated by the bars below the traces. Calibration bars: 10 min and 10 mmHg.

RESULTS

The experiments were performed on 21 isolated guinea-pig brains. The arterial basal resistance to perfusion pressure was 34.26 ± 8.15 mmHg ($n=7$) and 33.36 ± 10.80 mmHg ($n=5$) at 15 and 32°C, respectively. No changes in vascular resistance were observed in untreated brains up to 6–7 h ($n=5$). At the onset of the experiment, the functional viability of the vessels was tested by intraluminal perfusion with U46619 (30–100 nM), a potent vasoconstrictor that activates the thromboxane A₂ receptors on cerebral arterioles.² The choice of U46619 was also justified by the fact that this agonist mimics a platelet-dependent vasoconstriction, and therefore better reflects a cerebrovascular pathogenic event. At the plateau of the constriction (21.89 ± 8.7 mmHg above basal; $n=13$), ACh (1 μ M) was added to the perfusion buffer, in the presence of U46619, to test the ability of the endothelial cells to release NO and induce vasodilation (Fig. 1A). ACh caused a fast-onset dilation that reduced by 16.62 ± 4.59 mmHg ($\sim 76\%$) the U46619-dependent increase vessel tone. The ACh-induced dilation was antagonized by simultaneous arterial perfusion with the muscarinic receptor antagonist atropine (0.1 μ M; $n=11$; Fig. 1B). The effect of atropine was fully reverted upon washing of the preparation with control solution (Fig. 1C). The fall in vascular resistance to perfusion pressure determined by ACh was functionally antagonized by the NO scavenger Methylene Blue (10 μ M; $n=3$; Fig. 2B), suggesting that NO release mediates the ACh effect in the isolated brain vasculature of the guinea-pig. The inhibitory effect of Methylene Blue was not reversible, even after repeated washings of the preparation with control buffer. To confirm, in our experimental conditions, the involvement of NO in the ACh-induced vasodilation, the preparation was intraluminally perfused with either one of the NOS inhibitors, L-NNA ($n=3$) and L-NAME ($n=11$). In eight of 14 experiments, treatment with the NOS inhibitors increased vascular tone (25.9 ± 13.44 mmHg above basal values). In six of 14 experiments, no changes in the perfusion pressure occurred

even after long-lasting perfusion (approximately 3 h) with the NOS inhibitors. In seven of eight experiments, we observed an impairment of the ACh-dependent vasodilation, which required at least 3.5 h of continuous perfusion with the NOS inhibitor to be revealed. In Fig. 3A and B, the effect of ACh before and during L-NNA is shown. The NO donor NP (0.1 μ M) was briefly perfused after abolishing the ACh-induced vasodilation, in order to test the vasodilating capacity of the brain vasculature. In seven tests with NP during NOS inhibitors, a prompt vasodilation was obtained (Fig. 3B).

The ACh-dependent vasodilation observed during the early stages of perfusion with NOS inhibitors was followed by a slowly increasing constriction (Figs 4, 5A). In five experiments in which NOS inhibitors increased vessel tone, the disappearance of the vasodilation unmasked a 14.40 ± 3.85 mmHg vasoconstriction upon ACh perfusion (Fig. 4Ad). As illustrated in Fig. 5, the full development of vasoconstrictor efficacy occurred according to a slower time-course (190.2 ± 115.34 s; $n=5$) in comparison with the pre-NOS inhibitor ACh dilation (18.4 ± 15.88 s; $n=5$). The constriction was abolished by co-administration of 0.1 μ M atropine (Fig. 5C; $n=2$). Atropine alone perfused for 3–4 h did not change the vascular tone ($n=2$; not shown).

DISCUSSION

The present study demonstrates that ACh influences cerebral vessel tone via two, possibly simultaneous, effects, both mediated by muscarinic receptors: a fast NO-dependent dilation and a slower vasoconstriction. The vasodilatory effect of ACh is dependent on the release of NO from the endothelium, as demonstrated by its abolition following arterial perfusion with NOS inhibitors and the NO scavenger Methylene Blue. Recently, Methylene Blue has been shown to possess additional pharmacological activities.³⁷ However, previous studies by one of the authors demonstrated the efficacy of Methylene Blue as an NO scavenger by comparison with




Fig. 2. The NO scavenger Methylene Blue (10 μ M; B) blocks the vasodilation induced by ACh (1 μ M) on the U46619-induced constriction. Calibration bars: 10 min and 20 mmHg.




Fig. 3. The ACh-induced vasodilation (A) is blocked by arterial perfusion with 1 mM L-NNA (B). In these conditions, perfusion with the NO donor NP (0.1 μM) induced a dramatic vasodilation, indicating that the ability of the endothelium to react to NO was functionally preserved. The baseline difference between the two traces reflects the tonic vasoconstriction determined by L-NNA. Resting arterial resistance to perfusion pressure = 28 mmHg (dotted line). Calibration bars: 10 min and 5 mmHg.




Fig. 4. Effect of perfusion with L-NAME (30 μM) at slow time resolution (A). The events outlined in A are shown on a faster time-scale in B. In control conditions, a brief ACh (1 μM) application induced vasodilation on the U46619 constriction (a). After 3 h of perfusion with L-NAME, the ACh-induced dilation progressively vanished (b, c) and a constrictory response was isolated (d). Calibration bars: 5 min and 20 mmHg.




Fig. 5. Responses to 1 μM ACh before (A) and during perfusion with L-NAME (30 μM; B). The ACh-induced constriction observed during L-NAME was prevented by simultaneous atropine perfusion (C). Resting arterial resistance to perfusion pressure = 31.5 mmHg (dotted line). Calibration bars: 10 min and 10 mmHg.

oxyhaemoglobin at equimolar concentrations.²¹ The dilatory action of ACh mediated by NO has been demonstrated previously in cats, rats and other vertebrates.^{3,18,19,25,26,28,39,45} Cerebral cholinergic vasodilation is mediated by intracerebral arterioles¹² and pial arterioles,⁹ as well as by isolated capacitance arteries.²⁰ Accordingly, we propose that the changes in vascular resistance to constant flow observed in our experimental conditions reflect the modification in the vascular district of higher resistance represented by small arterioles and capillaries. Further experiments that allow direct measurements of vessel diameter will be necessary to elucidate the issue.

A constricting effect of ACh cerebral arteries was reported previously on isolated cat pial arteries.¹⁹ In this study, constriction was induced by very high concentrations of ACh (10–1000 μM), whereas more physiological dosages,

similar to those utilized in our experiments (1 μM), induced dilation exclusively. The vasoconstriction revealed by the blockade of the ACh-mediated relaxation may represent the effect of the activation of muscarinic receptors located on arteriolar smooth muscle cells, postsynaptic to the perivascular cholinergic terminals.^{4,5,43} The slower onset of vasoconstriction in comparison to the fast NO-dependent vasodilation could be accounted for by the time required for ACh to cross the blood–brain barrier and attain the necessary bioavailability to explicate its direct effect on muscle cells. An ACh-mediated increase in vascular tone has also been demonstrated when the endothelium is damaged in peripheral vessels²² and in isolated cerebral arterioles.³⁹ In line with these observations, Rosenblum³⁶ recently showed that NO-mediated vasodilation was selectively impaired during the period that follows ischemia and reperfusion, probably

because of the reduced ability of the endothelium to synthesize NO after the ischemic insult. According to our results and to the findings discussed above, it can be speculated that the ACh-mediated vasoconstriction unmasked by the functional impairment of the endothelium could worsen the ischemic damage by decreasing blood flow in a condition that requires an enhanced oxygen supply.

L-NAME, but not other NOS inhibitors such as L-NNA, has been demonstrated to exert an anti-cholinergic action by blocking the muscarinic receptors in pulmonary vessels,³³ hindlimb⁴⁶ and coronary arteries.¹⁰ The anti-muscarinic action of L-NAME has never been demonstrated in the cerebral vascular system, and has recently been questioned both by binding studies in the rat brain²⁴ and by functional studies in the small intestine.⁶ The slow time-course of the L-NAME effect observed in our experiments (onset after 3 h of direct arterial perfusion) is compatible with an enzymatic inhibition rather than a receptor antagonism. The demonstrations that (i) L-NNA and L-NAME block ACh-mediated dilation with a similar time-course, (ii) atropine-sensitive, ACh-induced vasoconstriction is still observed after prolonged perfusions with L-NAME (up to 4 h), and (iii) intraluminal perfusion with atropine does not lead to vasoconstriction, suggest that the increase in vascular tone mediated by L-NAME should not be ascribed to receptor antagonism of ACh on cerebral vessels, i.e. to the release of a cholinergic dilatory tone.

In a recent study, cholinergic denervation of cortical microvessels in Alzheimer's disease was found to be coupled with

an increase in the caliber of the cortical capillaries, interpreted as a compromised ability to adapt cortical perfusion via cholinergic modulation.⁴¹ These results might represent indirect evidence that cholinergic activation exerts a tonic vasoconstriction on the cerebral vascular system. Our observation of an ACh-mediated vasoconstriction could speculatively be called upon to explain these results.

CONCLUSIONS

The present study demonstrates, for the first time, a direct constrictory effect of physiological concentrations of ACh on undamaged cerebral vessels in which the NO-dependent dilation has been functionally impaired. The experiments confirm that the *in vitro* isolated guinea-pig brain is a suitable preparation to study the brain vascular system in a condition of preservation of the neuronal activity, vascular responsiveness and endothelial function. The combined evaluation of the interactions between the neuronal and vascular compartments in this preparation could be useful to understand the complex mechanisms that regulate brain excitability in physiological and pathological conditions.

Acknowledgements—The study was partially supported by Schering-Plough (Milan, Italy) and by the Italian Health Ministry. We would like to thank Dr Carmen Balsamo for the technical contribution in the early phase of the study.

REFERENCES

- Adachi T., Inanami O. and Sato A. (1992) NO is involved in increased cerebral cortical blood flow following stimulation of the nucleus basalis of Meynert in anesthetized rats. *Neurosci. Lett.* **139**, 201–204.
- Akopov S., Sercombe R. and Seylaz J. (1996) Cerebrovascular reactivity: role of endothelium/platelet/leukocyte interactions. *Cerebrovasc. Brain Metab. Rev.* **8**, 11–94.
- Alonso M. J., Salaices M., Sanchez-Ferrer C. F. and Marin J. (1992) Predominant role of NO in the relaxation induced by acetylcholine in cat cerebral arteries. *J. Pharmac. exp. Ther.* **261**, 12–20.
- Armstrong D. M. (1986) Ultrastructural characterization of choline acetyltransferase-containing neurons in the basal forebrain innervation of intracerebral blood vessels. *J. comp. Neurol.* **250**, 81–92.
- Arneric S. P., Honig M. A., Milner T. A., Greco S., Iadecola C. and Reis D. J. (1988) Neuronal and endothelial sites of acetylcholine synthesis and release associated with microvessels in rat cerebral cortex: ultrastructural and neurochemical studies. *Brain Res.* **454**, 11–30.
- Bartho L., Horvath G. and Lenard L., Jr (1999) Lack of anticholinergic effect of L-NAME in the small intestine. *Eur. J. Pharmac.* **16**, 279–282.
- Biella G. and de Curtis M. (2000) Olfactory inputs activate the medial entorhinal cortex via the hippocampus. *J. Neurophysiol.* **83**, 1924–1931.
- Biesold D., Inanami O. and Sato Y. (1989) Stimulation of nucleus basalis of Meynert increases cerebral blood flow in rats. *Neurosci. Lett.* **98**, 39–44.
- Busija D. W. and Heistad D. D. (1982) Atropine does not attenuate cerebral vasodilation during hypercapnia. *Am. J. Physiol.* **242**, H683–H687.
- Buxton I. L., Cheel D. J., Eckman D., Westfall D. P., Sanders K. M. and Keef K. D. (1993) *N*-Nitro-L-arginine methyl ester and other alkyl esters of arginine are muscarinic receptor antagonists. *Circulation Res.* **72**, 387–395.
- Chédotal A., Cozzari C., Faure M.-P., Hartman B. K. and Hamel E. (1994) Distinct ChAT and VIP bipolar neurons project to local blood vessels in the rat cerebral cortex. *Brain Res.* **646**, 181–193.
- Dacey R. G., Jr and Basset J. E. (1987) Cholinergic vasodilation of intracerebral arterioles in rats. *Am. J. Physiol.* **253**, H1253–H1260.
- Dauphin F., Lacombe P., Sercombe R., Hamel E. and Seylaz J. (1992) Hypercapnia and stimulation of the substantia innominata increase rat frontal cortical blood flow by different cholinergic mechanisms. *Brain Res.* **553**, 75–83.
- de Curtis M., Paré D. and Llinas R. R. (1991) The electrophysiology of the olfactory hippocampal system in the isolated and perfused adult mammalian brain *in vitro*. *Hippocampus* **1**, 341–354.
- de Curtis M., Arcelli P., De Biasi S., Spreafico R. and Avanzini G. (1994) Ultrastructural features of the isolated guinea-pig brain maintained *in vitro* by arterial perfusion. *Neuroscience* **59**, 775–788.
- de Curtis M., Biella G., Buccellati C. and Folco G. (1998) Simultaneous investigation of the neuronal and vascular compartments in the guinea pig brain isolated *in vitro*. *Brain Res. Protocols* **3**, 221–228.
- de Curtis M., Manfridi A. and Biella G. (1998) Activity-dependent pH changes and periodicity of spontaneous interictal spikes. *J. Neurosci.* **18**, 7543–7551.
- Edvinsson L., Owman C. and Sjöberg N.-O. (1976) Autonomic nerves, mast cells and amine receptors in human brain vessels: a histochemical and pharmacological study. *Brain Res.* **115**, 377–393.
- Edvinsson L., Falck B. and Owman C. (1977) Possibilities for cholinergic action on smooth musculature and on sympathetic axons in brain vessels mediated by muscarinic and nicotinic receptors. *J. Pharmac. exp. Ther.* **200**, 117–126.
- Faraci F. M., Brian J. E. and Heistad D. D. (1995) Response of cerebral blood vessels to an endogenous inhibitor of NOS. *Am. J. Physiol.* **269**, H1522–H1527.
- Feroli R., Folco G. C., Ferretti C., Gasco A. M., Medana C., Fruttero R. and Civelli G. A. (1995) A new class of furoxan derivatives as NO donors: mechanism of action and biological activity. *Br. J. Pharmac.* **114**, 816–820.
- Furchgott R. F. and Zawadzki J. V. (1980) The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholine. *Nature* **288**, 373–376.

23. Heistad D. D., Marcus M. L., Said S. I. and Gross P. M. (1980) Effect of acetylcholine and VIP on cerebral blood flow. *Am. J. Physiol.* **239**, H73–H80.
24. Hellmich B. and Gyermek L. (1997) *N*-Nitro-L-arginine methyl ester: a muscarinic receptor antagonist? *Fundam. clin. Pharmac.* **11**, 305–314.
25. Hylland P. and Nilsson G. E. (1995) Evidence that acetylcholine mediates increased cerebral blood flow velocity in crucian carp through an NO-dependent mechanism. *J. cerebr. Blood Flow Metab.* **15**, 512–524.
26. Iadecola C., Pelligrino D. A., Moskowitz M. A. and Lassen N. A. (1994) NO synthase inhibition and cerebrovascular regulation. *J. cerebr. Blood Flow Metab.* **14**, 175–192.
27. Iadecola C. and Zhang F. (1996) Permissive and obligatory role of NO in cerebrovascular response to hypercapnia and acetylcholine. *Am. J. Physiol.* **271**, R990–R1001.
28. Irikura K., Maynard K. I. and Moskowitz M. A. (1994) Importance of NO synthase inhibition to the attenuated vascular responses induced by topical L-NA during vibrissae stimulation. *J. cerebr. Blood Flow Metab.* **14**, 45–48.
29. Lacombe P., Sercombe R., Verrecchia C., Philipson V., MacKenzie E. T. and Seylaz J. (1989) Cortical blood flow increases induced by stimulation of the substantia innominata in the unanesthetized rat. *Brain Res.* **491**, 1–14.
30. Librizzi L., Folco G. and de Curtis M. (1999) Acetylcholine-mediated constriction of cerebral arteries is unmasked by NO synthase inhibitors in the *in vitro* isolated guinea-pig brain. *Neurosci. Lett., Suppl.* **52**, S24.
31. Llinas R., Yarom Y. and Sugimori M. (1981) Isolated mammalian brain *in vitro*: new technique for analysis of electrical activity of neuronal circuit function. *Fedn Proc.* **40**, 2240–2245.
32. Matsuda M., Meyer J. S., Deshmukh V. D. and Tagashira Y. (1976) Effect of acetylcholine on cerebral circulation. *J. Neurosurg.* **45**, 423–431.
33. McMahon T. J., Hoo J. S., Bellan J. A. and Kadowitz P. J. (1991) *N*-Nitro-arginine methyl ester selectively inhibits pulmonary vasodilator responses to acetylcholine and bradykinin. *J. appl. Physiol.* **71**, 2026–2031.
34. Muhlethaler M., de Curtis M., Walton K. and Llinas R. R. (1993) The isolated and perfused brain of the guinea-pig *in vitro*. *Eur. J. Neurosci.* **5**, 915–926.
35. Raszkievicz J. L., Linville D. G., Kerwin J. F., Wagenaar F. and Arneric S. P. (1992) NO synthase is critical in mediating basal forebrain regulation of cortical cerebral circulation. *J. Neurosci. Res.* **33**, 129–135.
36. Rosenblum W. I. (1997) Selective impairment of response to acetylcholine after ischemia/reperfusion in mice. *Stroke* **28**, 448–451.
37. Scobey C. G. and Faraci F. M. (1997) Effects of a novel inhibitor of guanylyl cyclase on dilator responses of mouse cerebral arterioles. *Stroke* **28**, 837–843.
38. Scremin O. U., Rovere A. A., Raynald A. C. and Giardini A. (1973) Cholinergic control of blood flow in the cerebral cortex of the rat. *Stroke* **4**, 233–239.
39. Shimizu T., Rosenblum W. I. and Nelson G. H. (1993) M3 and M1 receptors in cerebral arterioles *in vivo*: evidence for downregulated or ineffective M1 when endothelium is intact. *Am. J. Physiol.* **264**, 665–669.
40. Tanaka K., Gotoh F., Gomi S., Takashima S., Mihara B., Shirai T., Nogawa S. and Nagata E. (1991) Inhibition of NO synthesis induces a significant reduction in local cerebral blood flow in the rat. *Neurosci. Lett.* **127**, 129–132.
41. Tong X. K. and Hamel E. (1999) Regional cholinergic denervation of cortical microvessels and NOS-containing neurons in Alzheimer disease. *Neuroscience* **92**, 163–175.
42. Vaucher E., Borredon J., Seylaz J. and Lacombe P. (1995) Autoradiographic distribution of cerebral blood flow increase elicited by stimulation of the nucleus basalis magnocellularis in the unanesthetized rat. *Brain Res.* **691**, 68–75.
43. Vaucher E. and Hamel E. (1995) Cholinergic basal forebrain projections project to cortical microvessels in the rat: electron microscopic study with anterogradely transported *Phaseolus vulgaris* leucoagglutinine and choline acetyltransferase immunocytochemistry. *J. Neurosci.* **15**, 7427–7441.
44. Vaucher E., Linville D. and Hamel E. (1997) Cholinergic basal forebrain projections to nitric oxide synthase-containing neurons in the rat cerebral cortex. *Neuroscience* **79**, 827–836.
45. Wei E. P., Kukreja R. and Kontos H. A. (1992) Effects in cats of inhibition of NO synthesis on cerebral vasodilation and endothelium-derived relaxing factor from acetylcholine. *Stroke* **23**, 1623–1628.
46. White D. G., Drew G. M., Gurden J. M., Penny D. M., Roach A. G. and Watts I. S. (1993) The effect of *N*-nitro-L-arginine methyl ester upon basal blood flow and epithelium-dependent vasodilation in the dog hindlimb. *Br. J. Pharmac.* **108**, 763–768.
47. Zhang F., Xu S. and Iadecola C. (1995) Role of NO and acetylcholine in neocortical hyperemia elicited by basal forebrain stimulation: evidence for an involvement of endothelial NO. *Neuroscience* **69**, 1195–1204.