

Timing of dietary fat exposure and mammary tumorigenesis: Role of estrogen receptor and protein kinase C activity

Leena Hilakivi-Clarke^{1,2} and Robert Clarke^{1,3}

¹Lombardi Cancer Center; ²Department of Psychiatry, and ³Department of Physiology, Georgetown University, Washington, DC, USA

Abstract

The possible association between a high fat diet and increased breast cancer risk has remained controversial. This largely reflects the conflicting data obtained from migrant, case control and animal studies, which generally support this association, and cohort studies which often fail to show a link between fat and breast cancer. The mammary gland is particularly sensitive to estrogens during fetal development, leading us to hypothesize that dietary fat levels during this period may significantly influence breast cancer risk. Using chemically-induced mammary tumors in rats as our experimental model, we have demonstrated the ability of a maternal diet, high in the polyunsaturated fatty acid (PUFA) linoleic acid, to alter mammary gland differentiation, accelerate the onset of sexual maturation, and increase breast cancer risk. The mammary glands of female rats exposed to a high-fat diet *in utero* have more of the undifferentiated structures (terminal end buds) and fewer of the differentiated structures (alveolar buds) than the glands of rats exposed to a low-fat diet *in utero*. Furthermore, these mammary glands contain lower levels of total estrogen receptors and have reduced total protein kinase C activity. These effects appear to be mediated by an increase in the serum estradiol levels of pregnancy, which are elevated at least 30% in pregnant dams fed a high-fat diet. Furthermore, the administration of estradiol to pregnant dams produces effects on mammary gland development, onset of puberty and sensitivity to chemical carcinogenesis comparable to those seen in the offspring of rats fed a high fat diet during pregnancy. Our results, thus, support the hypothesis based on epidemiological data that high maternal estrogen levels increase daughters' breast cancer risk. The results also suggest that a high-fat diet may be an important factor in increasing pregnancy estrogenic activity. (Mol Cell Biochem **188**: 5–12, 1998)

Key words: breast cancer, estrogen, dietary fat, polyunsaturated fatty acid, estradiol, pregnancy

Dietary fat and breast cancer

The greatest potential for future success in reducing breast cancer mortality is to identify new prevention strategies for this disease. Diet is suggested to be involved in the multistep process that leads to clinical manifestation of breast cancer. It has been estimated that 35% of breast cancers could be prevented by appropriate dietary modifications [1]. However, despite extensive human and animal research, controversies exist concerning the possible link between nutritional components, including fat, fiber and vitamins, and breast cancer risk.

The initial observation linking dietary fat to breast cancer was the high correlation noted between national fat intake and breast cancer risk across different countries [2]. This observation led to a series of case control studies, the majority of which implicated that cases consumed more fat than controls [3, 4]. Numerous animal studies also showed that feeding a diet high in polyunsaturated fatty acids (PUFAs) promoted carcinogen-induced mammary tumorigenesis [5, 6]. Other types of fats, including saturated fats, were less effective in animals [5]. In humans, total or saturated fat intake appeared to be most closely linked to breast cancer [4]. Additional controversy arose when it became apparent that

the majority, but not all, cohort studies did not establish an increased breast cancer risk associated with a high-fat diet [7].

Sensitive periods and a high-fat exposure

We have proposed that a high-fat diet increases breast cancer risk if consumed during periods when the mammary gland is sensitive to endogenous estrogens, such as fetal life [8]. Evidence in support of the hypothesis includes the observation that Asian women, who consume a low-fat diet in their home country and have a low breast cancer incidence [9], when immigrating to the United States reach the high breast cancer incidence of Western countries within a few succeeding generations [10, 11]. Breast cancer risk among Asian immigrants is already 80% higher after a decade of living in the US, when compared with the risk of Asian women living in the East [11]. The most dramatic increase in risk occurs between Asian-Americans born in the West and those born in the East [11]. A higher dietary fat intake during pregnancy in the West than in the East may be responsible for the transition towards higher breast cancer risk between Asian generations living in the West.

Animal studies provide more direct evidence that a maternal intake of a high-fat diet during pregnancy increases spontaneous or carcinogen-induced mammary tumor incidence in the female offspring [12, 13]. The dietary fat source in these studies was corn oil that is high in PUFAs, particularly linoleic acid. The earlier studies have generally explored the effect of fat on promotion or progression of breast cancer. None of these studies were constructed/performed in a manner that would enable them to address our hypothesis, that dietary exposure during early life can significantly increase breast cancer risk in later life. Thus, an important period during which the mammary gland may be sensitive to dietary fat, might have been overlooked.

Mechanisms mediating the effects of dietary fat

Diet and estrogens

The mechanisms through which dietary fat alters mammary tumorigenesis, are unclear. Several possible mechanisms exist, including direct effects on lipid signaling pathways, and indirect effects mediated through perturbations in sex steroid levels (Table 1). Since estrogens have been extensively implicated in affecting breast cancer risk [14], one likely factor is the apparent ability of fat to alter serum estrogen

levels (Table 2). Several clinical studies show that both a high fat and/or total caloric intake increase the levels of circulating free estrogens, whereas a low-fat diet is associated with low plasma estrogen levels [15–18]. Results obtained in animal studies are less clear, but support the link between an isocaloric high-fat diet and high serum estrogen levels [19, 20]. Our findings clearly indicate that a diet high in corn oil increases serum estradiol (E2) levels in pregnant rats [13, 21]. This increase does not persist in the offspring after birth, which is consistent with the rapid clearance of maternal estrogens from neonates.

At least three possible mechanisms exist for a high-fat diet to increase circulating estrogens. Firstly, a high fat intake tends to lead to accumulation of adipose tissue, which is an important site for the conversion of androstenedione to estrone [25]. Secondly, arachidonic acid, a metabolite of PUFAs, activates P450 aromatase that then increases conversion of androstenedione to estrone [26]. Finally, PUFAs can reduce the binding of estrogens to serum binding proteins, including both sex-hormone binding globulin (SHBG) and albumin, thereby increasing the circulating levels of biologically potent estrogens [27].

Estrogen receptor and breast cancer

The fat-induced elevation in circulating estrogens is likely to indirectly affect mammary glands and tumors by influencing

Table 1. Putative mechanisms of action of dietary fat.

Generation of active lipid peroxides [22]
Indirect effects on signal transduction and gene expression through alterations in cellular membrane structure and function [23]
Direct effects on lipid-mediated signal transduction pathways, e.g., by altering the levels of arachidonic acid and eicosanoids [24]
Alterations in the levels/bioavailability of sex steroid hormones and their receptors [15–21]

Table 2. Observations demonstrating potential associations between dietary fat consumption, perturbations in serum estrogen levels and breast cancer risk.

Lifetime exposure to estrogens is lower in Asian women, who also consume a low fat diet and have a lower breast cancer risk [9]
A low fat diet can reduce serum estrogen levels [18]
Elevated serum estrogen levels are associated with increased breast cancer risk in some women [73]
Obese women have elevated serum estrogen levels, and postmenopausal obesity is associated with increased breast cancer risk [74]

estrogen's interactions with its nuclear estrogen receptors (ER). The ER is present *in utero* [28], and its concentrations in the mammary gland increase between birth and the first week of life [29]. The ER content remains at a relatively constant level after the prepubertal period, but appears to vary in concert with the estrus cycle [30]. A change in mammary ER levels also occurs during other periods when the levels of estrogens vary. For example, during pregnancy when circulating estrogen levels are high, the ER concentrations are low, but detectable [31]. A marked increase in mammary ER content and a decrease in serum estrogens occurs during lactation [32]. Thus, the changes in ER levels during pregnancy and lactation reflect the downregulation of ER by estrogens [33].

ER-positive breast tumors, which account for about 60% of all breast cancers [34] probably arise from within the ER positive epithelial cell populations of normal breast tissue [14]. Some 90% of these are associated with adjacent, ER-positive, non-neoplastic tissue [35]. ER-positive tumors tend to be more differentiated [36–38], exhibit a slower growth pattern [39, 40] and a better overall prognosis [41, 42]. A lower proportion of ER-positive tumors are found in premenopausal than postmenopausal breast cancer patients [43]. These findings could suggest that low mammary ER levels may be associated with an increased risk to develop premenopausal breast cancer. Since the data are based on ER α measurements, the role of ER β in affecting breast cancer risk remains to be determined.

Estrogen receptor and dietary fat intake

Since a high-fat intake elevates serum estrogens, a logical consequence of a high dietary fat consumption is an alteration in mammary ER status. Therefore, we have explored whether a diet high in n-6 PUFA affects ER protein levels in the mammary gland. Surprisingly, female mice consuming a high-fat diet exhibit a 6-fold increase in the mammary ER content [44]. A similar increase has been seen in DMBA-induced mammary tumors in female rats fed with a high corn oil diet [20]. These results would suggest that PUFAs may up-regulate ER, and this increase cannot be reversed with a simultaneous increase in circulating estrogens.

The effect of fat intake on ER in the breast has not been directly explored in human populations. Indirect evidence to suggest that a high-fat diet may induce ER is available from studies showing that either obese women, or women consuming a high-fat diet, are more likely to develop ER-positive mammary tumors than women consuming a low-fat diet [45]. Thus, a dietary fat appears to increase mammary ER content both in animals and humans. This increase is not likely to be caused by a fat-induced elevation in serum estrogens. However, it appears that the presence

ER may be required for fat to promote tumor growth and, perhaps, elevate the levels of this receptor. There is no correlation between the recurrence of ER negative breast tumors and dietary fat intake, while ER positive tumors are more likely to metastasize if a woman is consuming a high-fat diet [46].

Where associations between dietary fat and increased breast cancer risk have been reported, these are almost exclusively found in postmenopausal patients [47, 48]. A higher proportion of ER-positive tumors arise in postmenopausal women [43]. However, a significant proportion of human breast tumors are ER-negative [34], whereas the carcinogen-induced rodent mammary tumor models are ER-positive and strongly estrogen-dependent [49]. An exposure to a high-fat diet promotes the growth of mammary tumors in animal models [5]. These observations are consistent with the hypothesis that a high-fat diet preferentially affects breast cancer in ER-positive mammary glands. Thus, the effect of dietary fat on mammary gland ER content may explain some of the difference between animal studies and human cohort studies concerning a high-fat diet and breast cancer. In humans, a high-fat diet would preferentially increase breast cancer growth among the approximately 60% of the study population with ER-positive tumors. This could impact the power of some cohort studies to detect a significant overall trend for increased breast cancer risk in women consuming a high fat diet [7].

Estrogen receptor and dietary fat exposure during sensitive periods in mammary gland development

The results in humans and animals suggest that high ER levels in the mammary gland are associated with fat-induced promotion and progression of breast cancer. However, high ER levels prior to the carcinogen exposure may not affect the susceptibility to develop carcinogen-induced mammary tumors. An exposure to a high-fat diet prior to breast cancer initiation in adult animals does not appear to have significant effects on mammary tumorigenesis [5]. However, fat is likely to increase mammary ER content both before and after carcinogen exposure. Thus, high mammary ER levels in normal adult mammary glands do not increase the risk to develop carcinogen-induced tumors.

It was of interest to determine whether *in utero* exposure to a high-fat diet, that affects preinitiation events that increase susceptibility to develop mammary tumors, also affects mammary ER levels. Our data show that a maternal exposure to a high-fat diet induces a 4-fold reduction in the ER content in the offspring's mammary gland, when compared with ER content in the low-fat offspring [44]. This finding is similar to that showing that early postnatal treatment with DES causes a permanent reduction in the concentrations of ER

in the mouse mammary gland and DMBA induced tumors in rats [50, 51]. The decrease in the mammary ER content both in the high-fat and DES-exposed offspring suggest that the factor(s) responsible may be a high maternal estrogenic environment. There are several clinical observations that support the link between high maternal estrogen levels during pregnancy and increased breast cancer risk among daughters (Table 3). Thus, elevated circulating E2 levels during pregnancy in the mothers consuming a high-fat diet as well as maternal DES exposure could have resulted in a permanent down-regulation of ER in the offspring's mammary gland.

We do not have a clear explanation for why a high-fat diet increases ER levels when consumed by adult animals (and in breast tumors in humans), and why it reduces ER levels if the exposure occurs through a pregnant mother. In both cases, a high-fat diet increases serum E2 levels. However, the developmental state of the mammary gland, and the respective endocrinologic environments, are quite different in fetal and adult life. This suggests that the response of the gland to fat/E2 may vary during lifetime.

Another explanation for differences between adult and fetal exposure may be that *in utero* dietary fat may predominantly affect ER β receptors. These may be the predominant ER form during fetal life. We have used a ligand binding assay, which does not distinguish between ER α and ER β . The ER β cDNA was recently cloned from the rat prostate [52]. The DNA-binding domain of this ER β is highly homologous to that in the 'classical' ER α protein, while the C-terminal ligand binding domain bears only 55% homology. Most estrogenic substances or estrogenic antagonists compete with E2 for binding to both ER subtypes with identical preference and potency, but transcript tissue distribution is quite different for the ER α and ER β mRNA [53]. ER β mRNA expression is high in the ovary and prostate, moderate in uterus and testis, and low in the brain [53]. The distribution of ER subtypes in the breast is still unclear. However, human breast cancer cell lines appear to express less ER β mRNA than normal breast tissues [53, 54].

Taken together, the observations described above strongly suggest that both the timing and duration of exposure to a

high-fat diet may induce different responses in the mammary gland. Whether this is due to differential expression of ER β vs. ER α remains to be established. Nevertheless, the potential for the mammary gland to exhibit different responses to estrogens at different times could have considerable impact on how we think about estrogenic exposure and breast cancer risk, and how to modulate risk through chemoprevention or other dietary means.

Protein kinase C and breast cancer

Hormones can regulate cellular functions by activating some of the several isoforms of protein kinase C (PKC) [55, 56]. For example, E2 increases PKC δ expression in the uterus [57]. In human breast cancer cell lines PKC isoenzymes down-regulate ER mRNA expression [58–60]. Protein kinase C also is dependent on diacylglycerol (a fatty acid metabolite) and calcium for activation.

While PKC is linked to breast cancer [55, 61], its role is not fully understood. PKC activity is higher in the malignant than benign breast tissues [61], and higher in the more aggressive than in the less aggressive phenotype of human breast cancer cell lines [62]. However, reduced expression of some PKC isoforms, such as PKC η , is associated with increased neoplastic transformation in the mammary gland, while expression of other isoforms, such as PKC θ is linked to a more aggressive neoplastic process [62, 63].

Protein kinase C and dietary fat intake

Diet influences PKC activity. A high-fat diet increases PKC activity [64], whereas caloric restriction inhibits PKC activity [65] in epidermal cells in Sencar mice. A high corn oil diet also enhances PKC activity in the colon and carcinogen-induced colon tumors in male rats [66]. Further, a diet high in corn oil can block the inhibition of skin carcinogenesis, and reverse the reduction in PKC activity induced by moderate energy restriction [67]. Our data indicate that similarly to skin and colon, a diet high in n-6 PUFA increases PKC activity in the mouse mammary gland [44]. Since a high-fat intake increases mammary tumor incidence in animal models [5], the higher PKC activity in the mammary glands of the high-fat fed mice parallels the observed association between malignant progression and high PKC activity.

Protein kinase C and dietary fat exposure during sensitive periods

In contrast to the results obtained in adult mice fed with a high n-6 PUFA diet, PKC activity appears to be reduced in the

Table 3. Observations associating breast cancer risk with *in utero* estrogenic exposure.

Dizygotic twins have a high estrogenic pregnancy environment. The daughters of these pregnancies have an increased breast cancer risk [75–77].

High birth weight is associated with high levels of estrogens during pregnancy. The daughters of these pregnancies have an increased breast cancer risk [78, 79].

Preeclamptic and eclamptic pregnancies are accompanied by low serum estrogen levels. The daughters of these pregnancies have a lower breast cancer risk [80, 81].

mammary gland in the offspring of mothers that were kept on a high-fat diet during pregnancy [44]. It is possible that the reduced PKC activity reflects specific morphological changes in the mammary gland that increase susceptibility to neoplastic transformation. *In utero* exposure to a high-fat diet alters the normal development of a mammary gland [13, 44]. The number of terminal end buds, structures that are the targets of malignant transformation in the rodent mammary gland and possibly in the human breast [68, 69], is higher in the high-fat offspring than in a low-fat offspring. Thus, these structures that are sensitive to neoplastic changes, persist in the mammary glands of female mice exposed to a high-fat diet *in utero*. Persistent TEBs also have been reported in transgenic mice [70] and in rats exposed to either a high-fat diet or estradiol (E2) *in utero*, or during early postnatal period [13, 71]. All these groups exhibit an increased incidence of malignant growth in the mammary glands. These findings suggest that low PKC activity (as well as low ER content) in the mammary gland may increase the subsequent susceptibility to develop mammary tumors.

Conclusions

In conclusion, a diet high in PUFAs, when exposure occurs during adult life (mammary gland is well developed) increases the amounts of ER and PKC in the mammary gland. These events may be linked to the fat-induced increase in mammary tumorigenesis in animal models [5] and the effects (or lack of them) on breast cancer risk in humans [3, 7, 72]. In marked contrast, maternal intake of a high-fat diet during pregnancy reduces the ER content and PKC activity in the offspring's mammary gland and alters the gland's state of differentiation (Fig. 1). Since these offspring are at an increased risk to develop spontaneous and carcinogen-induced mammary tumors [12, 13], low amounts of ER and low PKC activity in the mammary gland may predict an increased breast cancer risk. In addition, the expression of ER β may be higher in the fetal than adult mammary gland [82], and therefore a maternal high-fat diet may affect offspring's breast cancer risk by affecting this novel ER subtype. Our future studies will determine whether a maternal high-fat intake specifically affects the ER α or ER β in the offspring. We also plan to investigate whether the low PKC activity reflects a reduction in activity of a specific isoform of PKC family of genes.

Acknowledgments

This work was supported by grants from the American Cancer Society (CN80420 to L.H.-C.), and the Public




Fig. 1. Plausible mechanisms mediating the effects of an exposure to a diet high in polyunsaturated fatty acids (a) after a carcinogen administration or (b) *in utero* on breast cancer risk. TEB – terminal end bud; AB – alveolar bud; ER – estrogen receptor (ER α and ER β); PKC – Protein kinase C.

Health Service (P50-CA58185 to R.C.) and the Lombardi Cancer Center Shared Animal Resource Facility (P30-CA51008).

References

- Boyd NF, Cousins M, Lockwood G, Tritchler D: The feasibility of testing experimentally the dietary fat-breast cancer hypothesis. *Br J Cancer* 62: 878–881, 1990
- Carroll KK, Khor HT: Dietary fat in relation to tumorigenesis: In: *Progress in Biochemistry and Pharmacology*. Basel, Karger, 1975, pp 308–353
- Howe GR, Hirohata T, Hislop TG, Iscovich JM, Yuan JM, Katsoyanni K, Lubin F, Marubini E, Modan B, Rohan T: Dietary factors and risk of breast cancer: combined analysis of 12 case-control studies. *J Nat Cancer Inst* 82: 561–569, 1990
- Boyd NF, Martin LJ, Noffel M, Lockwood GA, Tritchler DL: A meta-analysis of studies of dietary fat and breast cancer risk. *Br J Cancer* 68: 627–636, 1993
- Welsch CW: Relationship between dietary fat and experimental mammary tumorigenesis: A review and critique. *Cancer Res* 52: 2040–2048, 1992
- Fay MP, Freedman LS: Analysis of dietary fats and mammary neoplasms in rodent experiments. *Breast Cancer Res Treat* 46: 215–223, 1997
- Hunter DJ, Spiegelman D, Adami HO, Beeson L, van den Brandt PA, Folsom AR, Fraser GE, Goldbohm RA, Graham S, Howe GR, Kushi LH, Marshall JR, McDermott A, Miller AB, Speizer FE, Wolk A, Yuan S, Willett W: Cohort studies of fat intake and the risk of breast cancer - a pooled analysis. *N Engl J Med* 334: 356–361, 1996
- Hilakivi-Clarke LA: Mechanisms of high maternal fat intake during pregnancy on increased breast cancer risk in female rodent offspring. *Breast Cancer Res Treat* 46: 119–214, 1997
- Adlercreutz H, Gorbach SL, Goldin BR, Woods MN, Dwyer JT, Hamalainen E: Estrogen metabolism and excretion in oriental and caucasian women. *J Natl Cancer Inst* 86: 1076–1082, 1994
- Thomas DB, Karagas MR: Cancer in first and second generation Americans. *Cancer Res* 47: 5771–5776, 1987
- Ziegler RG, Hoover RN, Pike MC, Hildesheim A, Nomura AMY, West DW, Wu-Williams AH, Kolonel LN, Horn-Ross PL, Rosenthal JF, Hyer MB: Migration patterns and breast cancer risk in Asian-American women. *J Natl Cancer Inst* 85: 1819–1827, 1993
- Walker BE: Tumors in female offspring of control and diethylstilbestrol-exposed mice fed high-fat diets. *J Nat Cancer Inst* 82: 50–54, 1990
- Hilakivi-Clarke L, Clarke R, Onojafe I, Raygada M, Cho E, Lippman ME: A maternal diet high in n-6 polyunsaturated fats alters mammary gland development, puberty onset, and breast cancer risk among female rat offspring. *Proc Natl Acad Sci USA* 94: 9372–9377, 1997
- Clarke R, Dickson RB, Lippman ME: Hormonal aspects of breast cancer. Growth factors, drugs and stromal interactions. *Crit Rev Oncol Hematol* 12: 1–23, 1992
- Goldin BR, Adlercreutz H, Gorbach SL, Warram JH: Estrogen excretion patterns and plasma levels in vegetarian and omnivorous women. *N Engl J Med* 307: 1452–1457, 1982
- Bennett FC, Ingram DM: Diet and female sex hormone concentrations: an intervention study for the type of fat consumed. *Am J Clin Nutr* 52: 808–812, 1990
- Rose DP, Boyar AP, Cohen C, Strong LE: Effect of a low-fat diet on hormone levels in women with cystic breast disease. *J Natl Cancer Inst* 78: 623–626, 1987
- Rose DP, Connolly JM, Chlebowski RT, Buzzard IM, Wynder EL: The effects of a low-fat dietary intervention and tamoxifen adjuvant therapy on the serum estrogen and sex hormone-binding globulin concentrations of postmenopausal breast cancer patients. *Breast Cancer Res Treat* 27: 253–262, 1993
- Chan PC, Head JF, Cohen LA, Wynder EL: Influence of dietary fat on the induction of mammary tumors by nitrosomethylurea: Associated hormone changes and differences between Sprague-Dawley and F344 rats. *J Natl Cancer Inst* 59: 1279–1283, 1977
- Ip C, Ip MM: Serum estrogens and estrogen responsiveness in 7,12-dimethylbenz(a)anthracene-induced mammary tumors as influenced by dietary fat. *J Natl Cancer Inst* 66: 291–295, 1981
- Hilakivi-Clarke L, Onojafe I, Raygada M, Cho E, Clarke R, Lippman M: High-fat diet during pregnancy increases breast cancer risk in rats. *J Natl Cancer Inst* 88: 1821–1827, 1996
- Welsch CW: Review of the effects of dietary fat on experimental mammary gland tumorigenesis: Role of lipid peroxidation. *Free Radic Biol Med* 18: 757–773, 1995
- Stubbs CD, Smith AD: Modification of mammalian membrane polyunsaturated fatty acid composition in relation to membrane fluidity and function. *Biochim Biophys Acta* 779: 89–137, 1984
- Rose DP: Effects of dietary fatty acids on breast and prostate cancers: Evidence from *in vitro* experiments and animal studies. *Am J Clin Nutr* 66: 1513S–1522S, 1997
- Greenblatt RB, Colle ML, Mahesh VB: Ovarian and adrenal steroid production on the postmenopausal women. *Obstet Gynecol* 47: 383–387, 1976
- Noble LS, Simpson ER, Johns A, Bulun SE: Aromatase expression in endometriosis. *J Clin Endocrinol Metab* 81: 179, 1996
- Bruning PF, Bonfrer JMG: Possible relevance of steroid availability and breast cancer. *Ann NY Acad Sci* 257–264, 1990
- Holderegger C, Keefer D: The ontogeny of the mouse estrogen receptor: The pelvic region. *Am J Anatomy* 177: 285–297, 1986
- Muldoon TG: Steroid hormone receptor regulation by various hormonal factors during mammary development and growth in the normal mouse. *Ann NY Acad Sci* 464: 17–36, 1986
- Anderson E, Clarke RB, Howell A: Changes in the normal human breast throughout the menstrual cycle: Relevance to breast carcinogenesis. *Endocrine-related Cancer* 4: 23–33, 1997
- Kelly PA, Dijiane J, Malancon R: Characterization of estrogen, progesterone and glucocorticoid receptors in rabbit mammary glands and their measurement during pregnancy and lactation. *J Steroid Biochem* 18: 215–221, 1983
- Yu WC, Leung BS: Variation of cytoplasmic content of estrogen and progesterone receptors in the mammary gland and uterus of rats at time of parturition. *Biol Reprod* 27: 658–664, 1982
- Haslam SZ, Nummy KA: The ontogeny and cellular distribution of estrogen receptors in normal mouse mammary gland. *J Steroid Biochem Mol Biol* 42: 589–595, 1992
- McGuire WL, Carbone PP, Sears ME, Escher GC: Estrogen receptors in human breast cancer: An overview. In: W.L. McGuire, P.P. Carbone, M.E. Sears, G.C. Escher (eds). *Estrogen Receptors in Human Breast Cancer*. Raven Press, New York, 1975, pp 1–7
- Netto GJ, Cheek JH, Zachariah NY, Romer JC, Lee CF, Schoenfeld R, Chakmakjian ZH: Steroid receptors in benign mastectomy tissue. *Am J Clin Pathol* 94: 14–17, 1990
- Blanco G, Alavaikko M, Ojala A, Collan Y, Hietanen T, Aine R, Taskinen PJ: Estrogen and progesterone receptors in breast cancer: Relationships to tumor histopathology and survival of patients. *Anticancer Res* 4: 383–390, 1984
- Singh L, Wilson AJ, Baum M, Whimster WF, Birch IH, Jackson IM, Lowrey C, Palmer MK: The relationship between histological grade, oestrogen receptor status, events and survival at 8 years in the NATO (Nolvadex) trial. *Br J Cancer* 57: 612–614, 1988
- Henry JA, Nicholson JA, Farndon JR, Westley BR, May F: Measurement of oestrogen mRNA levels in human breast tumors. *Br J Cancer* 58: 600–605, 1988
- Anatnades K, Spector H: Quantitative estrogen receptor values and growth of carcinoma of the breast before surgical intervention. *Cancer* 50: 793–796, 1982

40. Meyers JS, Rao BR, Stevens SC, White WL: Low incidence of estrogen receptor in breast carcinomas with rapid rates of cellular proliferation. *Cancer* 40: 2290–2298, 1977
41. Aamdal S, Borner O, Jorgensen O, Host H, Eilassen G, Kaalhus O, Pihl A: Estrogen receptor and long term prognosis in breast cancer. *Cancer* 53: 2525–2529, 1984
42. Clark GM, McGuire WL: Steroid receptors and other prognostic factors in primary breast cancer. *Semin Oncol* 15: 20–25, 1988
43. Honig SF: Treatment of metastatic disease: hormonal therapy and chemotherapy. In: J.R. Harris, M.E. Lippman, M. Morrow, S. Hellman (eds). *Diseases of the Breast*. Lipponcott-Raven, Philadelphia, 6, pp 669–734
44. Hilakivi-Clarke LA, Raygada M, Stoica A, Martin M-B: Consumption of a high-fat diet during pregnancy alters estrogen receptor content, protein kinase C activity and morphology of mammary gland in the mother and her female offspring. *Cancer Res* 58: 654–660, 1998
45. Potter JD, Cerhan JR, Sellers TA, McGovern PG, Drinkard C, Kushi LR, Folsom AR: Progesterone and estrogen receptors and mammary neoplasia in the Iowa Women's Health Study: How many kinds of breast cancer are there. *J Natl Cancer Inst* 85: 32–36, 1995
46. Holm L-E, Nordevang E, Hjalmar ML, Lidbrink E, Callmer E, Nilsson B: Treatment failure and dietary habits in women with breast cancer. *J Natl Cancer Inst* 85: 32–36, 1993
47. Willett WC, Hunter DJ: Prospective studies of diet and breast cancer. *Cancer* 74: 1085–1089, 1994
48. Witte JS, Ursin G, Siemiatycki J, Thompson WD, Paganini-Hill A, Haile RW: Diet and premenopausal bilateral breast cancer: A case-control study. *Breast Cancer Res Treat* 42: 243–251, 1997
49. Russo IH, Medado J, Russo J: Endocrine influences on the mammary gland. In: T.C. Jones, U. Mohr, R.D. Hunt (eds). *Integument and Mammary Glands*. Springer-Verlag, Berlin, 1994, pp 252–266, 1994
50. Bern HA, Mills KT, Ederly M: Estrogen-associated defects in rodent mammary gland development. In: J.A. McLachlan (ed.). *Estrogens in the Environment*. Elsevier, The Netherlands, 1985, pp 319–326,
51. Verhoeven G, Vandoren G, Heyns W, Kuhn ER, Janssens JP, Teuwen D, Goddeeris E, Lesaffre E, DeMoor P: *J Endocrinol* 95: 357–368, 1982
52. Kuiper GGJM, Enmark E, Peltö-Huikko M, Nilsson S, Gustafsson JA: Cloning of a novel estrogen receptor expressed in rat prostate and ovary. *Proc Natl Acad Sci USA* 93: 5925–5930, 1996
53. Kuiper GGJM, Carlsson B, Grandien K, Enmark E, Haggblad J, Nilsson S, Gustafsson JA: Comparison of the ligand binding specificity and transcript tissue distribution of estrogen receptors alpha and beta. *Endocrinology* 138: 863–870, 1997
54. Vladusic EA, Hornby AK, Guerra FK, Lupu R: Expression of ERbeta in breast cancer cells: Does it play an important role? *Proc Am Assoc Cancer Res* 88, Abstract 1995, 1997
55. Nishizuka Y: Intracellular signalling by hydrolysis of phospholipids and activation of protein kinase C. *Science* 258: 607–614, 1992
56. Dekker LV, Parker PJ: Protein kinase C - a question of specificity. *Trends Biochem Sci* 19: 73–77, 1994
57. Cutler RE, Maizels ET, Hunzicker-Dunn M: delta protein kinase-C in the rat ovary: Estrogen regulation and localization. *Endocrinology* 135: 1669–1678, 1994
58. Tzukerman M, Zhang XK, Pfahl M: Inhibition of estrogen receptor activity by the tumor promoter 12-O-Tetradecanylphorbol-13-Acetate: A molecular analysis. *Mol Endocrinol* 5: 1893–1992, 1991
59. Saceda M, Knabbe C, Dickson RB, Lippman ME, Bronzert D, Lindsey RL, Gottardis M, Martin M-B: Post transcriptional destabilization of estrogen receptor mRNA in MCF-7 cells by 12-O-tetradecanolphorbol-13-acetate. *J Biol Chem* 266: 17809–17814, 1991
60. Martin M-B, Garcia-Morales P, Stoica A, Harrison BS, Pierce M, Katz D, Zhang S, Danielson M, Saceda M: Effects of 12-O-tetradecanoylphorbol-13-acetate on estrogen receptor activity in MCF-7 cells. *J Biol Chem* 270: 25244–25251, 1995
61. O'Brian CA, Ward N: Biology of the protein kinase C family. *Cancer Metast Rev* 8: 199–214, 1989
62. Ways DK, Kukoly CA, de Vente J, Hooker JL, Bryant WO, Posekany KJ, Fletcher DJ, Cook PP, Parker PP: MCF-7 breast cancer cells transfected with PKC-alpha exhibit altered expression of other protein kinase C isoforms and display a more aggressive neoplastic phenotype. *J Clin Invest* 95: 1906–1915, 1995
63. Manni A, Buckalter E, Etindi R, Kunselman S, Rossini A, Mauger D, Dabbs D, Demers L: Induction of less aggressive breast cancer phenotype by protein kinase-alpha and beta overexpression. *Cell Growth Diff* 7: 1187–1198, 1996
64. Choe M, Kris ES, Luthra R, Copenhaver J, Pelling JC, Donnelly TE, Birt DE: Protein kinase C is activated and diacylglycerol is elevated in epidermal cells from SENCAR mice fed high fat diets. *J Nutr* 122: 2322–2329, 1992
65. Kris E, Choe M, Luthra R, Conway H, Barnett T, Yakine A, Birt DE: Protein kinase C activity is reduced in epidermal cells from energy restricted SENCAR mice. *J Nutr* 124: 485–492, 1994
66. Reddy BS, Simi B, Patel N, Aliaga C, Rao CV: Effect of amount and types of dietary fat on intestinal bacterial 7 alpha-dehydroxylase and phosphatidylinositol-specific phospholipase C and colonic mucosal diacylglycerol kinase and PKC activities during different stages of colon tumor promotion. *Cancer Res* 56: 2314–2320, 1996
67. Birt DE, Barnett T, Pour PM, Copenhaver J: High-fat diet blocks the inhibition of skin carcinogenesis and reductions in protein kinase C by moderate energy restriction. *Mol Carcinogenesis* 16: 115–120, 1996
68. Russo J, Gusterson BA, Rogers AK, Russo IH, Wellings SR, van Zwieten MJ: Comparative study of human and rat mammary tumorigenesis. *Lab Invest* 62: 244–278, 1990
69. Russo J, Romero AL, Russo IH: Architectural pattern of the normal and cancerous breast under the influence of parity. *Cancer Epidemiol Biomarkers Prev* 3: 219–224, 1994
70. Krane IM, Leder P: NDF/herregulin induces persistence of terminal end buds and adenocarcinomas in the mammary glands of transgenic mice. *Oncogene* 12: 1781–1788, 1996
71. Hilakivi-Clarke L, Cho E, Raygada M, Kenney N: Alterations in mammary gland development following neonatal exposure to estradiol, transforming growth factor alpha, and estrogen receptor antagonist ICI 182,780. *J Cell Physiol* 170: 279–289, 1997
72. Wolk A, Bergström R, Hunter D, Willett W, Ljung H, Holmberg L, Bergkvist L, Bruce A, Adami HO: A prospective study of association of monounsaturated fat and other types of fat with risk of breast cancer. *Arch Intern Med* 158: 41–45, 1998
73. Toniolo PG, Levitz M, Zeleniuh-Jacquette A, Banerjee S, Koenig KL, Shore RE: A prospective study of endogenous estrogens and breast cancer in postmenopausal women. *J Natl Cancer Inst* 87: 190–197, 1995
74. Ingram D, Nottage E, Ng S, Sparrow L, Roberts A, Willcox D: Obesity and breast cancer. *Cancer* 64: 1049–1053, 1989
75. Weiss HA, Potischman N, Brinton LA, Brogan D, Coates RJ, Gammon MD, Malone KE, Schoenberg JB: Prenatal and perinatal risk factors for breast cancer in young women. *Epidemiology* 8: 181–187, 1997
76. Braun MM, Ahlbom A, Floderus B, Brinton LA, Hoover RN: Effect of twinning on incidence of cancer of the testis, breast, and other sites. *Cancer Causes Control* 6: 519–524, 1995
77. Swetdlow AJ, Stavola BLD, Swanwick MA, Maconochie NES: Risk of breast and testicular cancers in young adult twins in England and Wales: evidence on prenatal and genetic aetiology. *Lancet* 350: 1723–1728, 1997
78. Michels KB, Trichopoulos D, Robins JM, Rosner BA, Manson JE, Hunter D, Colditz GA, Hankinson SE, Speizer FE, Willett WC: Birth-weight as a risk factor for breast cancer. *Lancet* 348: 1542–1546, 1996

79. Sanderson M, Williams M, Malone KE, Stanford JL, Emanuel I, White E, Daling JR: Perinatal factors and risk of breast cancer. *Epidemiology* 7: 34–37, 1996
80. Ekbom A, Trichopoulos D, Adami HO, Hsieh CC, Lan SJ: Evidence of prenatal influences on breast cancer risk. *Lancet* 340: 1015–1018, 1992
81. Ekbom A, Hsieh C-c, Lipworth L, Adami H-O, Trichopoulos D: Intrauterine environment and breast cancer risk in women: A population-based study. *J Natl Cancer Inst* 89: 71–76, 1997
82. Brandenberger AW, Tee MK, Lee JY, Chao V, Jaffe RB: Tissue distribution of estrogen receptor α (ER α) and β (ER β) mRNA in the mid gestational human fetus. *J Clin Endocrinol Metab* 82: 3509–3512