


This article was downloaded by: [New York University]

On: 17 February 2015, At: 10:33

Publisher: Routledge

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of the American College of Nutrition

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uacn20>

Auto-immune complications of D-penicillamine—a possible result of zinc and magnesium depletion and of pyridoxine inactivation.

M S Seelig

Published online: 02 Sep 2013.

To cite this article: M S Seelig (1982) Auto-immune complications of D-penicillamine—a possible result of zinc and magnesium depletion and of pyridoxine inactivation., Journal of the American College of Nutrition, 1:2, 207-214, DOI: [10.1080/07315724.1982.10718989](https://doi.org/10.1080/07315724.1982.10718989)

To link to this article: <http://dx.doi.org/10.1080/07315724.1982.10718989>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Hypothesis

Auto-Immune Complications of D-Penicillamine – A Possible Result of Zinc and Magnesium Depletion and of Pyridoxine Inactivation

Mildred S. Seelig, MD, MPH, FACN

Department of Medicine, Goldwater Memorial Hospital, New York University Medical Center, Roosevelt Island

Long-term high-dosage penicillamine treatment of patients with advanced stages of diseases with autoimmune components has resulted in very few adverse reactions in a series of over 50 such patients also given selected nutrients: pyridoxine, zinc and magnesium (which penicillamine inactivates or chelates), and vitamins B₁, B₁₂, and E (which have sulfhydryl-protective activity). The patients on this regimen have been essentially free of the side effects that occur in about a third of patients treated with penicillamine without such supplements. Reports of myasthenia gravis—a disease with abnormalities of the thymus and of T-cells, as a side effect of penicillamine—suggest that zinc, magnesium, and pyridoxine might be the agents most likely to be protective. Pyridoxine is necessary for cellular accumulation of zinc and magnesium, deficiencies of which have caused thymic and other immunologic abnormalities. Whether the other vitamins administered contribute to the favorable results requires further study.

Key words: autoimmune reactions, D-penicillamine, immunology, magnesium, zinc, pyridoxine, thymus, T cells, rheumatoid arthritis, Laennec cirrhosis

INTRODUCTION

D-penicillamine's effect on immunoregulation [1,2] is indicated by its activity against autoimmune diseases [3-5] and its paradoxical production of immunologically mediated side effects [1,2,5]. Depending on circumstances, it can either inhibit or augment lymphocyte transformation in vitro [2]. The development of myasthenia gravis during the course of administration of D-penicillamine, rare though it is [6-10], provides a clue to its immunologic toxicity. Thymic hyperplasia [6] and abnormal

Received December, 1981; accepted February, 1982.

Address reprint requests to Dr. Mildred S. Seelig, Department of Medicine, Goldwater Memorial Hospital, New York University Medical Center, Roosevelt Island, New York NY 10044.

thymus-dependent lymphocytes (T cells) [2] have been reported in patients who developed this side effect. Similar changes have been described in spontaneous myasthenia gravis [11,12]. Thymic hyperplasia has also been seen in D-penicillamine toxicology studies [13]. The changes resemble those produced by experimental magnesium or zinc depletion [14–19] and those found in acrodermatitis enteropathica, a clinical zinc deficiency disorder [20]. Because penicillamine chelates zinc and magnesium [21] and because it inactivates pyridoxine [22], which is necessary for cellular uptake of zinc and magnesium [23–25], penicillamine's immunologic dysregulatory effects might be partially mediated by interference with utilization of these nutrients. In more than 50 patients with advanced chronic diseases (with autoimmune components) who were given selected nutrients before and during treatment with D-penicillamine in high dosages for prolonged periods, there was a very low incidence of adverse reactions [26–28]. We attribute this predominantly to protection against drug-induced deficiencies of magnesium, zinc, and pyridoxine.

METHODS

A combination of magnesium, zinc, and vitamins B₆, B₁, B₁₂, C, and E was given to patients with Laennec cirrhosis (because of underlying deficiencies and anticipated drug-induced losses) while awaiting approval for use of D-penicillamine [28]. This new application was to lower the elevated copper levels of cirrhosis [29], to apply its effect on collagen turnover [30] to hepatic fibrosis, and to treat the autoimmune component of cirrhosis [4]. The supplements were given orally three times a day with meals: vitamins B₆ (50 mg), B₁ (50 mg), B₁₂ (25 µg), C (500 mg), and vitamin E (400 IU), zinc (10–15 mg as the gluconate) and magnesium (100 mg as an amino acid chelate). Magnesium sulfate (100 mg of Mg) was given intramuscularly once a week to those with evidence of deficiency, as determined by percentage retention of a test dose. Vitamin B₁₂ was also given intramuscularly, (2,000 µg) once a week because of its demonstrated malabsorption in cirrhotic patients with pancreatic insufficiency [28,31]. Partial improvement of iron-refractory anemia, strength, and sense of well-being led us to continue the supplements when D-penicillamine treatment was started (given at least 1 hour apart from meals and supplements) 1–3 months after the nutrient regimen had been started [28]. There was early response to 250-mg and 500-mg daily doses of penicillamine for as long as 3 months at each dosage level, but cessation of improvement led to increase of dosage according to the accepted gradual incremental regimen [5]. A few patients required increases to as much as 2.75 g/day to sustain improvement. The same regimen was utilized in treatment of patients with multiple sclerosis in an attempt to obtain benefit from its activity in autoimmune disease and from its activity against neurotropic viruses [26], and with rheumatoid arthritis with and without vasculitis (unpublished observations) in order to obtain data with identical treatment programs. Hospital staff shortages necessitated reduction in the scope of our studies (in terms of number of patients and number of daily doses given) after 5 years. Among those remaining in the program, daily dosage was increased (in those whose dosage had been reduced to 250–500 mg/day) when prior gains were not sustained on lower doses. Patients given up to 500 mg of penicillamine daily are usually given one-third the total daily dosage of the supplements (at one meal). Several patients in another study of rheumatoid arthritis, who were not on the nutritional supplement program and who

were intolerant of penicillamine, had treatment cautiously restarted in combination with only tocopherol (I.A. Jaffe, personal communication). In addition to frequent evaluation for clinical changes, all patients have been monitored at least monthly throughout the treatment program for hematologic, urinary, and other side effects.

RESULTS

The patients with Laennec cirrhosis, who received the full therapeutic regimen, exhibited early correction of refractory anemia and diminution of hepatic pain and enlargement in those with concomitant hepatitis, as well as correction of inverted albumin/globulin ratio and needle liver biopsy evidence of less fibrosis [28]. Several of those with multiple sclerosis showed decreased spasticity and improved vision, mobility, and electroencephalograms during the full treatment phase of the study [26], which subsequently plateaued and showed no further improvement or some regression. A few with rheumatoid arthritis, who have been more recently entered into the study, have shown striking improvement. The improvement of the cirrhotic patients was sustained only on doses of 1–2.0 g/day (or higher in a few), with the exception of cirrhotic patients with encephalopathy or peripheral neuropathy, whose neurologic signs improved markedly on only 250 mg/day plus the nutrients. A patient with advanced active rheumatoid arthritis, who had improved while receiving 1.75 g/day, suffered an exacerbation when her penicillamine dosage was reduced to 250 mg/day. She showed gradual relief of pain, lowering of sedimentation rate, and improvement in anemia as the penicillamine dosage was gradually increased to (750 mg/day). Two, whose reactivation of advanced rheumatoid arthritis was not controlled by corticoids, have similarly improved on 500–750 mg of D-penicillamine plus supplements. Another, with severe rheumatoid arteritis and extensive necrosis of both legs, improved remarkably, with re-epithelialization and diminished pain, as dosage was gradually increased to 1,250 mg daily (M.S. Seelig and A.R. Berger, in preparation).

Among the more than 50 patients in this series, most of whom were treated for 6 months to 7 years, only one or possibly two serious side effects developed, and there were few transitory discomforts [27]. Hematuria developed in one cirrhotic patient; cystoscopy disclosed hemorrhagic cystitis. Since a renal biopsy to rule out immuno-complex glomerulitis as a contributory factor was refused, penicillamine treatment (but not the nutritional supplements) was discontinued. The patient's signs of hepatic dysfunction recurred in the hospital. He left, returned to drink, and died in hepatic failure the following year. A multiple sclerosis patient, with a history of anaphylaxis to penicillin, developed frank albuminuria (1 g/day) while receiving 750 mg penicillamine/day as an outpatient. Treatment was stopped, and he was lost to further study. Several patients developed slight to moderate proteinuria (1+ to 3+) at intervals, in association with urinary tract infections. Proteinuria subsided when the infection was controlled, without modification of penicillamine therapy. Hemorrhagic bullae over elbows and ecchymoses over legs and forearms developed in a cirrhotic patient while he was taking 2.75 g/day of penicillamine. The lesions cleared when the dosage was reduced to 500 mg/day and did not recur when clinical deterioration necessitated gradual increase of dosage to 1.75 g/day. Taste aberration or loss was encountered in two patients in their first month of treatment, before zinc supplementation had been

started. Transitory rashes were encountered in only two patients. No instance of thrombocytopenia or granulocytic leukopenia has been seen. Low pretreatment thrombocyte counts (in patients with Laennec cirrhosis or with multiple sclerosis), that necessitated very cautious institution of penicillamine therapy rose, during treatment. Low pretreatment hemoglobin and hematocrit levels also rose.

The rheumatoid arthritis patients who had been intolerant of D-penicillamine and had it restarted with tocopherol showed no improvement in tolerance of penicillamine (I.A. Jaffe, personal communication).

DISCUSSION

The unexpected occurrence of autoimmune disease complications in patients being treated with D-penicillamine [5-10], a drug with activity against autoimmune diseases [3-5,10,26,28], has led to many speculations and much study of its activity at the lymphocytic level [2,10] to elucidate the disparate findings. Thymic hyperplasia as a clinical [6] and experimental [13] toxicologic development, and the thymic and lymphocytic changes that are caused by depletion of two essential minerals namely, zinc and magnesium [14-20] that penicillamine chelates [21], suggest that the drug's first demonstrated clinical effect, as a chelator [32], might be responsible for the most serious problems associated with its use. Zinc is bound by penicillamine [21,33,34], which can cause retention [33,34] or loss of zinc [35,36], depending on whether it is given with or separately from the drug. Severe clinical manifestations of depletion developed in a child being treated for Wilson disease [35], and zinc levels were decreased in blood, brain, liver, and kidneys of mice given D-penicillamine [36]. Magnesium is low in the order of affinity of penicillamine for metals [21], but possibly when there is not an excess of trace or toxic minerals with greater affinity for the drug, even this macro-mineral can be removed. Preexistent magnesium inadequacy caused by cirrhosis or by drugs (such as diuretics or corticoids) might be intensified by penicillamine. The patients in this series have had marginally low serum magnesium levels while receiving penicillamine and have shown retention of most of parenterally administered test doses of magnesium (which is suggestive of deficiency), even though they all were being given magnesium supplements (100 mg three times daily), and several had received weekly intramuscular injections.

Inactivation of pyridoxine [22,36] has been demonstrated in rheumatoid arthritis patients being treated with penicillamine [22]; experimental vitamin B₆ deficiency causes cellular losses of both magnesium [23,24] and zinc [25], each of which is necessary for pyridoxine-dependent enzymes [37-39]. Cellular loss of magnesium and/or zinc might lead to aberrations of lymphoid tissue. Magnesium deficiency in rats has caused thymic hypertrophy [14-16] (or atrophy [16] under different conditions), lymphosarcoma, and leukemia [15,40] in certain strains. In mice, magnesium deficiency has reduced the number of antibody-synthesizing cells, thereby lowering immunoglobulin levels [39]. Optimal magnesium concentrations are required for normal *in vitro* lymphocyte transformation [42]. In rats, zinc deficiency causes inhibition of T-cell immunity [17], thymic atrophy and hypofunction [18], and impaired cell-mediated immunity [18]. Clinical zinc deficiency is associated with immunodeficiencies and T-cell dysfunction in infants [19,20] and thymic hypoplasia in calves [19,20] with acrodermatitis enteropathica, a disease that results from impaired intestinal zinc

absorption [20,43,44]. The effect of simultaneous deficiencies of magnesium, zinc, and pyridoxine, such as penicillamine can cause, has not been explored in animal models.

Few have investigated the effect of mineral supplements on the therapeutic or adverse effects of D-penicillamine. A mixture of minerals containing salts of magnesium, calcium, zinc, iron, and cobalt has been recommended to avoid depletion of essential minerals by the drug in Wilson disease patients [45]. The same supplement was given schizophrenic patients treated with penicillamine, with favorable results [46]. Improvement of schizophrenia with penicillamine was not confirmed by others who did not use the minerals [47]. It has been suggested that schizophrenics who improved on penicillamine plus minerals might have had pyroluria, a condition associated with zinc depletion, or kryptopyroluria (mauve factor), which is characterized by depletion of both zinc and pyridoxine and which improves somewhat on their administration [48]. Mineral supplements have been found to increase tolerance of high-dosage D-penicillamine in melanoma patients [49; and H.B. Demopoulos, personal communication].

The nutritional supplements given to the patients in this series [26–28] were selected to compensate for known penicillamine interference (with pyridoxine [22] and zinc [21,33–35]) and to substitute for the sulfhydryl-donor drug while we were awaiting clearance for administration of the drug, since increase in sulfhydryl levels might be a factor in therapeutic response to D-penicillamine [32,50,51]. Sulfhydryl-protective vitamins—B₁ [52], B₁₂ [53], E, and C [54,55]—were given at first with the intent of discontinuing them when penicillamine was started. A recent hypothesis, that auto-immune disorders might be mediated by free-radical-induced depression of T-cell suppressor function [56], suggests that the antioxidant activity of these vitamins, as well as that of D-penicillamine, might have played a role in the patients' improvement.

Subjective and hematologic improvement occurred while the patients were receiving the nutrients before penicillamine was started. Because vitamin assays disclosed persistently low levels, they were continued with penicillamine treatment. The same regimen was employed when new patients entered the study because of the absence of side effects, an incidence of 26–40% or more having been expected on the basis of reports of others [5,57,58]. Despite biochemical evidence of pyridoxine deficiency in rheumatoid arthritis patients on long-term penicillamine therapy [22], and of a disorder in pyridoxine metabolism in rheumatoid patients not receiving penicillamine [59], administration of only vitamin B₆ with penicillamine influenced neither the action of the drug nor its toxicity [5]. Zinc supplementation has corrected signs of severe zinc deficiency in a patient receiving penicillamine for Wilson disease [37], but its use has interfered with the therapeutic effects of low-dosage penicillamine in rheumatoid arthritis without preventing side effects [60]. In the present series, larger doses of zinc (30–45 mg/day) have not interfered with the therapeutic response to penicillamine in moderate to high doses (1–2.75 g/day) in patients with cirrhosis [28] or rheumatoid arthritis (unpublished observations).

Whether the vitamins that were provided in addition to pyridoxine, — which with magnesium and zinc are probably the most important components of the supplement mixture, contribute to the low incidence of side effects remains to be ascertained. Vitamin B₁ and magnesium activities are interdependent [61,62]. Magnesium deficiency intensifies some of the signs of vitamin E deficiency [63]. Vitamin E, magnesium, and zinc are each involved in maintaining membrane stability, and they and other vitamins and minerals play roles in immunosurveillance [64].

CONCLUSIONS

The most serious adverse reactions caused by D-penicillamine are those that develop late and that reflect its dysregulation of the immune system. The lack of such effects in a series of patients given high-dosage penicillamine might be the result of protecting against losses of cellular zinc and magnesium, depletions of which are associated with thymic and T-cell abnormalities.

ACKNOWLEDGMENTS

Appreciation is expressed to Herman Baker and Oscar Frank for vitamin assays and to Israeli A. Jaffe for unpublished data and for alerting me to the work demonstrating penicillamine's immunodysregulatory effects.

REFERENCES

1. Dawkins RL, Zilko PJ: Editorial. Penicillamine: Friend or foe? *Austral N Zeal J Med* 9:494-495, 1979.
2. Main RN, Roffe L: D-penicillamine and lymphocyte function in rheumatoid arthritis: The inhibitory and augmenting effects of D-penicillamine *in vitro* on phytohemagglutinin-induced lymphocyte transformation. In Munthe E (ed): "Penicillamine Research in Rheumatoid Disease." Oslo, Norway: Fabritius and Sonner, 1976, pp 172-184.
3. Jaffe IA: Rheumatoid arthritis with arteritis: Report of a case treated with D-penicillamine. *Ann Intern Med* 61:556-563, 1964.
4. Epstein O, DeVilliers D, Jains S, Potter BJ, Thomas HC, Sherlock C: Reduction of immune complexes and immunoglobins induced by D-penicillamine in primary biliary cirrhosis. *N Engl J Med* 300:274-278, 1979.
5. Jaffe A: Penicillamine treatment of rheumatoid arthritis. Rationale, pattern of clinical response and clinical pharmacology and toxicology. In Munthe E (ed): "Penicillamine Research in Rheumatoid Disease." Oslo, Norway: Fabritius and Sonner, 1976, pp 11-24.
6. Bucknall RC, Dixon AJ, Glick EN, Woodland J, Zutshi DW: Myasthenia gravis associated with penicillamine treatment of rheumatoid arthritis. *Br Med J* 1:600-602, 1975.
7. Masters CL, Dawkins RL, Zilko PJ, Simpson JA, Leedman RJ, Lindstrom J: Pencillamine associated myasthenia gravis, antiacetylcholine receptor and antistriational antibodies. *Am J Med* 63:689-694, 1977.
8. Gordon RA, Burnside JW: D-penicillamine-induced myasthenia gravis in rheumatoid arthritis. *Ann Intern Med* 87:578-579, 1977.
9. Atcheson SG, Ward JR: Ptosis and weakness after start of D-penicillamine therapy. *Ann Intern Med* 89:939-940, 1978.
10. Dawkins RL, Garlepp M, McDonald BL, Williamson J, Zilko PJ, Carrano J: Myasthenia gravis and penicillamine. *J Rheumatol* 8(Suppl 7):169-172, 1981.
11. Souadjian JV, Enriques P, Silverstein MN, et al: The spectrum of diseases associated with thymoma. Coincidence or syndrome? *Arch Intern Med* 134:374-379, 1974.
12. Richman DP, Patrick J, Arnason BGW: Cellular immunity in myasthenia gravis. Response to purified acetylcholine receptor and autologous thymocytes. *N Engl J Med* 294:694-698, 1976.
13. Jacobus DP, Bokelman DL, Majka JA: D-penicillamine in the monkey. In Munthe E Ed: "Penicillamine Research in Rheumatoid Disease." Oslo, Norway: Fabritius and Sonner, 1976, pp 25-40.
14. Hungerford GF, Karson EF: The eosinophilia of magnesium deficiency. *Blood* 16:1642-1650, 1960.
15. Jasmin G: Lymphoedeme, hyperplasie et tumefaction du tissu lymphatique chez le rat soumis a une diete deficiente en magnesium. *Rev Can Biol* 22:383-390, 1963.
16. Alcock NW, Shils ME, Lieberman PH, Erlandson RA: Thymic changes in the magnesium-depleted rat. *Cancer Res* 33:2196-2204, 1973.
17. Pekarek RS, Powanda MC, Hoagland AM: Effect of zinc deficiency on the immune response of the rat. *Fed Proc* 36:360, 1977.
18. Chandra RK, Au B: Single nutrient deficiency and cell-mediated immune responses. 1. Zinc. *Am J Clin Nutr* 33:736-738, 1980.
19. Andresen E, Basse A, Brummerstedt E, Flagstad T: Zinc and the immune system in

- cattle. *Lancet* 1:839-840, 1973.
20. Moynahan EJ: Zinc deficiency and cellular immune deficiency in acro-dermatitis enteropathica in man and zinc deficiency with thymic hypoplasia in Friesian calves: A possible genetic link. *Lancet* 2:710, 1975.
 21. Kuchinskas EJ, Rosen Y: Metal chelates of DL-penicillamine. *Arch Biochem Biophys* 97: 370-372, 1962.
 22. Jaffe IA, Altman K, Merryman P: The anti-pyridoxine effect of penicillamine in man. *J Clin Invest* 43:1869-1873, 1964.
 23. Aikawa JK: Effects of pyridoxine and desoxy-pyridoxine on magnesium metabolism in rabbits. *Proc Soc Exp Biol Med* 104:461-463, 1960.
 24. Durlach J: Donneés actuelles sur le mecanismes de synergie entre vitamine B₆ et magnesium. *J Med Besancon* 5:349-359, 1969.
 25. Hsu JM: Zinc content in tissues of pyridoxine deficient rats. *Proc Soc Exp Biol Med* 119: 177-180, 1965.
 26. Seelig MS, Alba A, Berger AR, Rudez A, Tarlau M: Pilot study of D-penicillamine, vitamins and minerals in multiple sclerosis. *J Clin Psychiatry* 39:170-174, 1978.
 27. Seelig MS, Berger AR: Do trace elements, magnesium and antioxidants protect against D-penicillamine toxicity? In Hemphill DD (ed): "Trace Substances in Environmental Health. A Symposium." Columbia, Mo: University of Missouri, 1977, Vol 11, pp 243-251.
 28. Seelig MS, Berger AR, Hazzi C, Milstoc M: Long term pilot study of D-penicillamine, minerals and vitamins in patients with advanced Laennec's cirrhosis. In Hemphill DD (ed): "Trace Substances in Environmental Health. A Symposium." Columbia, Mo: University of Missouri, 1979, Vol 13, pp 389-400.
 29. Butt EM, Nusbaum RE, Gilmour TC, DiDio SL: Trace metal patterns in disease states. *Am J Clin Pathol* 26:225-242, 1956.
 30. Harkness RD: In vitro and in vivo observations on the effect of penicillamine on collagen. *Postgrad Med J* 44:31-34, 1968.
 31. Matuchansky C, Rambaud R, Modigliani R, Bernier JJ: Vitamin B₁₂ malabsorption in chronic pancreatitis. *Gasteroenterology* 67: 406-407, 1974.
 32. Walshe JM: Penicillamine, a new oral therapy for Wilson's disease. *Am J Med* 21:487-495, 1956.
 33. McCall JT, Goldstein NP, Randall RV, Gross JB: Comparative metabolism of copper and zinc in patients with Wilson's disease (hepatolenticular degeneration). *Am J Med Sci* 254: 13-23, 1967.
 34. Aaseth J, Forre O, Munthe E, Froland SS, Abrahamsen T: Biological aspects of metal chelation during penicillamine treatment. In Munthe E (ed): "Penicillamine Research in Rheumatoid Disease." Oslo, Norway: Fabritius and Sonner, 1976, pp 80-87.
 35. Klingberg WG, Prasad AS, Oberleas D: Zinc deficiency following penicillamine therapy. In Prasad AS (ed): "Trace Elements in Human Health and Disease." New York: Academic Press, 1976, pp 51-65.
 36. Kuchinskas EJ, Horvath A, Duvigneaud V: On anti-vitamin B₆ effect of l-penicillamine. *Arch Biochem Biophys* 68:68-75, 1957.
 37. Lehninger AL: Role of metal ions in enzyme systems. *Physiol Rev* 30:393-429, 1950.
 38. McCormick DB, Gregory ME, Snell EE: Pyridoxal phosphokinases. I. Assay, distribution, purification and properties. *J Biol Chem* 236:2076-2084, 1961.
 39. Heaton FW: Magnesium in intermediary metabolism. In Cantin M, Seelig MS (eds): "Magnesium in Health and Disease." Jamaica, New York: SP Medical and Scientific Books, 1980, pp 43-55.
 40. Hass GM, McCreary PA, Laing GH: Lymphoproliferative and immunologic aspects of magnesium deficiency. In Cantin M, Seelig MS (eds): "Magnesium in Health and Disease." Jamaica, New York: SP Medical and Scientific Books, 1980, pp 185-200.
 41. Elin RJ: The effect of magnesium deficiency in mice on serum immunoglobulin concentrations and antibody plaque-forming cells. *Proc Exp Soc Biol Med* 148:620-624, 1975.
 42. Brennan JK, Lichtman MA: The effect of extracellular calcium and magnesium on the proliferation of murine lymphoblasts. *J Cell Physiol* 82:101-112, 1973.
 43. Moynahan EJ: Acrodermatitis enteropathica. A lethal inherited human zinc-deficiency disorder. *Lancet* 2:399, 1974.
 44. Hambidge KM, Neldner KH, Walravens PA. Zinc and acrodermatitis enteropathica. In Hambidge KM, Nichols BM (eds): "Zinc and Copper in Clinical Medicine." Jamaica, New York: SP Medical and Scientific Books, 1976, pp 81-98.
 45. Fister WP, Boulding JE, Baker RA: The treatment of hepatolenticular degeneration with penicillamine, with report of two cases. *Can Med Assoc J* 78:99-102, 1958.
 46. Nicolson GA, Greiner AC, McFarlane WJ, Barker A: Effect of penicillamine on schizophrenic patients. *Lancet* 1:344-347, 1966.

47. Brooker BK, Papeschi R, Murray L: Negative results of a trial of penicillamine in chronic schizophrenia. *Postgrad Med J* 44(Oct Suppl): 48-52, 1968.
48. Pfeiffer CC, Bacchi D: Copper, zinc, manganese, niacin and pyridoxine in the schizophrenias. *J Appl Nutr* 27:9-39, 1975.
49. Demopoulos HB: Effects of reducing the phenylalanine-tyrosine intake of patients with advanced malignant melanoma. *Cancer* 19: 657-664, 1966.
50. Lorber A, Chang C, Masuoka D, Meacham I: Effect of thiols in biological systems on protein sulfhydryl content. *Biochem Pharmacol* 19: 1551-1560, 1970.
51. Munthe E, Guldal G, Jellum E: Increased intracellular glutathione during penicillamine treatment for rheumatoid arthritis. *Lancet* 2: 1126-1127, 1979.
52. Kochetov GA, Lutovinova GF: Sulfhydryl groups and transketolase activity. *Biochem Biophys Res Commun* 22:129-134, 1966.
53. Register UD: Effect of vitamin B₁₂ on liver and blood non-protein sulphhydryl compounds. *J Biol Chem* 206:705-709, 1953.
54. Schwarz K: Vitamin E, trace elements and sulfhydryl groups in respiratory decline. *Vitamins Hormones* 20:463-484, 1962.
55. Tappel AL: Vitamin E as the biological lipid antioxidant. *Vitamins Hormones* 20:493-510, 1962.
56. Harman D: Free radical theory of aging: Beneficial effect of antioxidants on the life span of male NZB mice; role of free radical reactions in the deterioration of the immune system with age and in the pathogenesis of systemic lupus erythematosus. *Age* 3:64-73, 1980.
57. Stein HB, Patterson AC, Offer RA: Adverse effects of D-penicillamine in rheumatoid arthritis. *Ann Intern Med* 92:24-29, 1980.
58. Halverson PB, Kozin F, Bernhard GA, Goldman AL: Toxicity of penicillamine. A serious limitation to therapy in rheumatoid arthritis. *J Am Med Assoc* 240:1870-1871, 1978.
59. Bett IM: Effect of pyridoxine on tryptophan metabolism in rheumatoid arthritis. *Ann Rheum Dis* 21:388-391, 1962.
60. Mery C, Delrieu F, Ghozlan R, Saporta L, Simon F, Amor B, Menkes CJ, Delbarre F: Controlled trial of D-penicillamine in rheumatoid arthritis. Dose effect and the role of zinc. *Scand J Rheumatol* 5:241-247, 1976.
61. Itokawa Y, Inoue K, Natori Y, Okazaki K, Fujiwara M: Effect of thiamine on growth, tissue magnesium, thiamine levels and transketolase activity in magnesium deficient rats. *J Nutr Sci Vitaminol* 18:159-164, 1972.
62. Itokawa Y, Tseng LF, Fujiwara U: Thiamine metabolism in magnesium-deficient rats. *J Nutr Sci Vitaminol* 20:249-255, 1974.
63. Goldsmith LA: Relative magnesium deficiency in the rat. *J Nutr* 73:87-102, 1967.
64. Seelig MS: Magnesium (and trace substance) deficiencies in the pathogenesis of cancer. *Biol Trace Elem Res* 1:273-297, 1979.