

## Review

QJM

# Non-psychiatric comorbidity associated with Alzheimer's disease

A. DUTHIE<sup>1</sup>, D. CHEW<sup>2</sup> and R.L. SOIZA<sup>2</sup>

From the <sup>1</sup>Old Age Psychiatry Directorate, Royal Cornhill Hospital and <sup>2</sup>Department of Medicine for the Elderly, Woodend Hospital, Aberdeen, UK

Address correspondence to Dr R.L. Soiza, c/o Department of Medicine for the Elderly, Woodend Hospital, Aberdeen, AB15 6XS, UK. email: roy.soiza@nhs.net

## Summary

The burden of medical comorbidity in individuals with Alzheimer's disease is greater than that observed in matched individuals without dementia. This has important implications for all clinicians and health-care providers who deal with this common condition. The prevalence of vascular risk factors and vascular disease is particularly high. Additionally,

associations with a number of other chronic medical conditions have been described, including thyroid disorders, sleep apnoea, osteoporosis and glaucoma. This review gives an overview of evidenced medical (non-psychiatric) comorbidity associated with Alzheimer's disease and briefly explores the underlying mechanisms that may account for these associations.

## Introduction

In Alzheimer's disease, cognitive impairment influences the presentation and recognition of other medical conditions and the outcome of treatment. In turn, many physical health factors contribute to cognitive decline.

The burden of medical comorbidity in Alzheimer's disease is greater than matched individuals without Alzheimer's disease.<sup>1</sup> Higher medical comorbidity is significantly associated with lower cognition on mini-mental state examination taking into account age, gender, education and care setting.<sup>2</sup> Higher medical comorbidity is also significantly associated with poorer self-care, decreased mobility and greater incontinence, adjusted for the same demographic factors and cognition. Therefore, not only are medical comorbidities implicated in cognitive decline but also, independently of cognition, they may adversely affect function and independence. Moreover, an understanding of the non-psychiatric

associations of Alzheimer's disease may shed light on poorly understood pathophysiological mechanisms of this condition.

This article reviews the medical (non-psychiatric) comorbidity of Alzheimer's disease. Medical conditions with an evidenced association with Alzheimer's disease were identified through interrogating PubMed, Embase and Medline using a number of search terms including 'Alzheimer's', 'association' and 'comorbidity'. The review briefly explores the underlying mechanisms that may account for these associations and will be of practical use to clinicians dealing with patients with Alzheimer's disease.

## Vascular disease and Alzheimer's dementia

There is a particularly strong association between Alzheimer's disease and vascular disease as well as vascular risk factors. The pathological hallmark

of Alzheimer's disease is accumulation of  $\beta$ -amyloid plaques and the formation of neurofibrillary tangles,<sup>3</sup> while vascular dementia is characterized by ischaemic damage and multiple infarctions due to occlusion of cerebral blood vessels. Although these two diseases have long been considered separate entities, increasing evidence has shown that the ischaemic damage known to cause vascular dementia is also responsible for the development of Alzheimer's dementia, which could explain the overlap of the presence of both these diseases, known as mixed dementia.

In animal models, ischaemia results in increased amyloid deposition in the brain cortex,<sup>4,5</sup> and it is likely similar to processes that occur in humans.<sup>6</sup> Ischaemic damage disrupts the blood-brain barrier which is protective against the deposition of  $\beta$ -amyloid in the neurones and finally cause an up-regulation of apolipoprotein E that binds to  $\beta$ -amyloid causing conformational change in the peptide, which renders it neurotoxic. There is up-regulation of the expression of amyloid precursor protein and amyloid beta in response to ischaemic damage to the brain.

Elevated blood pressure is established as a strong risk factor for vascular dementia. The relationship between blood pressure and Alzheimer's disease is more complicated. Evidence shows that elevated blood pressure in midlife is a strong risk factor for Alzheimer's disease.<sup>7</sup> However, hypotension in later life, defined as a diastolic blood pressure <70 mmHg, has a similar effect. This relationship was described in the Chicago Health and Ageing Project,<sup>8</sup> where low later life blood pressure was related to higher incidence of Alzheimer's disease. This can be explained by the fact that hypotension in later life leads to ischaemic conditions that promote neuronal death.<sup>9</sup> The decreasing auto-regulatory capacity of the brain in later life causes the brain to be more vulnerable to ischaemic insults when systemic blood pressure falls beyond a critical level to maintain adequate perfusion.<sup>10</sup>

In a prospective study of 3734 Japanese-Asian men followed up for 25 years, there was a strong association between midlife hypertension and Alzheimer's and vascular dementia.<sup>11</sup> This effect was most pronounced in men not treated with anti-hypertensives. Data from autopsy and neuroimaging findings from the same study demonstrated that elevated systolic blood pressure in midlife caused a vasculopathic change along with reduced brain weight and a significant amount of neuritic ( $\beta$ -amyloid) plaques in the neocortex and hippocampus.<sup>7</sup> Similar findings have been observed in a Finnish study.<sup>12</sup>

Elevated blood pressure causes the blood-brain barrier disruption around the capillaries in the brain which in turn cause the dysfunction of subcortical vessels.<sup>13</sup> This subsequently impairs critical nutrient delivery to the brain, especially glucose which in turn impairs its performance. Hypertension also causes endothelial damage which has a 2-fold effect on the brain. First, it leads to impaired nitric oxide production resulting in the generation of free radicals.<sup>14</sup> Second, it triggers an inflammatory response that promotes neuritic plaque formation.<sup>15</sup> Extreme hypoxia has also been shown to cause amyloid precursor protein accumulation and its subsequent cleavage to amyloid.<sup>16</sup>

Diabetes mellitus may also be a contributing risk factor for Alzheimer's disease. Several large studies have shown that diabetes is associated with cognitive impairment.<sup>17–21</sup> A positive association has also been found between impaired glucose tolerance and Alzheimer's disease.<sup>22–24</sup> Leibson *et al.*<sup>21</sup> concluded that adult onset diabetes mellitus increased the risk of Alzheimer's disease by 37% in women and 127% in men. These findings are largely supported by results of other studies—the Rotterdam study showed a link between diabetes and Alzheimer's disease with a relative risk of 1.3.<sup>22,25</sup> Patients with diabetes have also been shown to have greater degrees of whole-brain atrophy and hippocampal atrophy.<sup>26,27</sup>

The association between diabetes mellitus and Alzheimer's disease probably involves several mechanisms. Peripheral hyperinsulinemia seen in type 2 diabetes results in blood-brain barrier dysfunction and decreased insulin transport to the brain.<sup>28,29</sup> Reduced brain insulin signalling causes increased tau phosphorylation and  $\beta$ -amyloid levels in a streptozotocin mouse model of diabetes.<sup>30</sup> Insulin receptors are found in high concentration in areas associated with memory functions such as the hypothalamus and limbic system, perhaps explaining the strong association between peripheral hyperinsulinemia and impaired cognition.<sup>31</sup> Hyperinsulinemia also provokes increase in central inflammation.<sup>32</sup> Animal studies have shown that this leads to neurodegeneration and cognitive deficits. In a study of diabetic rats, hyperglycaemia caused an upregulation of amyloid precursor protein, neuronal loss and neuritic degeneration.<sup>33</sup>

Obesity is the primary cause of insulin resistance. Lobo *et al.*<sup>34</sup> showed that 80% of obese individuals are insulin resistant. The public health implications of the growing number of obese individuals in the population may extend to exacerbate the predicted future rise in prevalence of dementia. Both a recent meta-analysis and a newly published twin study confirmed that obesity in midlife is a significant

risk factor for both Alzheimer's and vascular dementia,<sup>35,36</sup> though this association may disappear or even become reversed in later life.<sup>35</sup> Obesity results in free fatty acid elevation and inhibition of enzyme metalloprotease, which facilitates clearance of  $\beta$ -amyloid and normal insulin signalling.<sup>37</sup> Free fatty acid further stimulates the aggregation of amyloid and tau proteins as shown in the Honolulu Asia Ageing study.<sup>38</sup> Elevated free fatty acid produces a pro-inflammatory state with a rise in TNF- $\alpha$ ,<sup>39</sup> which may contribute to Alzheimer's dementia.<sup>40</sup> Elevated midlife total cholesterol level is associated with a 2- to 3-fold increased risk of Alzheimer's disease.<sup>41</sup> Cholesterol plays a large role in  $\beta$ -amyloid aggregation. Cholesterol-derived aldehydes promote Schiff's base formation, accelerating the early stages of amyloidogenesis.<sup>42,43</sup> Cerebral atherosclerosis is strongly linked to neuritic plaque formation<sup>44,45</sup> and cholesterol has also been implicated in their pathogenesis.<sup>46,47</sup>

## Thyroid

Both hypothyroidism and hyperthyroidism are well-recognized as potentially reversible causes of cognitive impairment. There is also a link between thyroid function at the higher and lower ends of the normal clinical range and risk of dementia.<sup>48</sup>

Kalmijn *et al.*<sup>49</sup> demonstrated a 3-fold increased risk of Alzheimer's disease with sub-clinical hyperthyroidism >2 years. In the 2009 Honolulu-Asia ageing study, thyroxine (but not thyrotropin) concentration was associated with increased risk of Alzheimer's disease in their male sample over a mean follow-up period of almost 5 years.<sup>50</sup> The authors found a higher Alzheimer's pathology load in those with higher free thyroxine in a sub-sample at post-mortem. A large sample of original Framingham cohort participants demonstrated an association between both high and low thyroid-stimulating hormone levels and dementia in females only.<sup>51</sup> Tan and Vasan<sup>48</sup> suggest that thyroid abnormalities are more likely causative of Alzheimer's pathology than consequence. Suggested mechanisms include effects on amyloid and tau deposition,<sup>50</sup> cholinergic function, or neuronal survival, or alternatively the indirect mediation of risk via vascular factors.<sup>50,51</sup>

## Sleep apnoea

Sleep apnoea is more common in the elderly than in younger individuals,<sup>52</sup> and various small studies demonstrate increased prevalence in Alzheimer's disease.<sup>53,54</sup> Various neuropsychological deficits have been described in sleep apnoea patients

without dementia. However, the deficits are mild and dissimilar in pattern to Alzheimer's disease and to vascular dementia.<sup>55</sup> In sleep apnoea, continuous positive airway pressure (CPAP) has been shown to improve cognitive functions in several domains.<sup>56</sup> A group investigating the effect of CPAP for sleep apnoea in Alzheimer's disease patients found improved sleep patterns<sup>56</sup> and cognition.<sup>57,58</sup> Deficits in cholinergic neurotransmission have been suggested as a cause for sleep apnoea in Alzheimer's disease. Moraes *et al.*<sup>59</sup> found that oxygen saturation, REM sleep duration and cognition improved with donepezil in a small group of Alzheimer's patients compared to placebo controls.

However, across studies the magnitude of difference of rates of sleep apnoea in Alzheimer's disease compared to controls has been generally small,<sup>60,61</sup> and other studies have failed to confirm an association.<sup>62</sup> Sleep apnoea overlaps with multiple medical conditions linked to impaired cognition, including hypothyroidism and hypertension. An association with apoE4 has been demonstrated.<sup>63</sup> Overall, the relationship between cognition and sleep apnoea remains unclear. To clarify the mechanisms involved, Alzheimer's disease and other types of cognitive impairment in sleep apnoea should be examined separately.

## Osteoporosis

A number of studies have demonstrated an association between low bone mineral density and both cognitive decline<sup>64</sup> and Alzheimer's disease.<sup>65,66</sup> Alzheimer's disease patients are more prone to osteoporosis, falls and fractures than non-demented subjects, with poorer outcomes following fracture. The health of bones is therefore important for disability and mortality in Alzheimer's disease patients since bone mineral density strongly predicts fracture. Weller and Schnatzer<sup>67</sup> found a prevalence of osteoporosis of 27% in nursing home residents with Alzheimer's disease cases compared to 16% in those without dementia. The Alzheimer's disease group also suffered higher rates of falls and an excess of hip fractures, even after accounting for the higher number of falls. The authors suggest that a further independent factor must account for the higher rates of fracture in Alzheimer's disease. However, their study design involved use of a questionnaire to identify the presence of osteoporosis and would have failed to identify undiagnosed bone loss.

More recently, an association has been demonstrated between low bone mineral density and brain atrophy in early Alzheimer's disease.<sup>68</sup> A

further paper by the same authors demonstrated reduced volume in the hypothalamus specifically in early Alzheimer's disease, again associated with low bone mineral density.<sup>69</sup> The authors propose that neurodegeneration leads to bone loss through neural or hormonal mechanisms. Alternatively, factors including weight loss, reduced soft tissue and nutritional deficiencies may underlie the relationships between cognition, bone mass and fractures.

## Glaucoma

Increased prevalence with age and neurodegenerative pathology are common factors in both Alzheimer's disease and glaucoma. Studies have demonstrated higher rates of glaucoma in Alzheimer's disease,<sup>70</sup> which are not explained by a shared factor of apoE4.<sup>71</sup> A study of 112 nursing home residents with Alzheimer's disease revealed an occurrence rate of 25.9% compared to 5.2% in matched non-demented controls.<sup>70</sup> Glaucoma is characterized by retinal ganglion cell death. Raised intraocular pressure is typically seen in glaucoma, but studies demonstrate that the disease can occur in the absence of raised pressure indicating the role of other pathological mechanisms. Considerable evidence supports the presence of the characteristic pathological mechanisms of Alzheimer's disease, of  $\beta$ -amyloid accumulation, neuronal apoptosis and cell loss, in glaucoma.<sup>72–75</sup> Optic nerve degeneration and loss of retinal ganglion cells have also been found in Alzheimer's disease. However, this has been found to occur in the absence of retinal or optic nerve amyloid or tau pathology.<sup>76,77</sup> Wostyn *et al.*<sup>78</sup> proposed low CSF pressure occurring in Alzheimer's disease as a factor leading to increased risk of glaucoma. While further research is required to clarify overlapping disease processes in glaucoma and Alzheimer's disease, existing work raises some interesting possibilities for therapy. Guo *et al.*<sup>73</sup> investigated treatments targeting  $\beta$ -amyloid for glaucoma, with some success.

## Cancer

Not all conditions are more prevalent in Alzheimer's disease. There is mounting evidence of a protective effect of Alzheimer's disease against developing cancer. One population-based prospective study found a slower rate of developing cancer in Alzheimer's disease, and also reduced risk of Alzheimer's disease in cancer.<sup>79</sup> The same investigating group failed to demonstrate an association between cancer and vascular dementia, indicating a link with neurodegenerative disease specifically.<sup>79</sup>

Disruption to biological mechanisms regulating cell apoptosis may be crucial to both cancer and Alzheimer's disease. One enzyme involved in cellular signalling, Pin1, appears to be over-expressed in many cancers and downregulated in Alzheimer's disease.<sup>80</sup>

## Rheumatoid arthritis and NSAIDs

Epidemiological studies have also demonstrated lower rates of Alzheimer's disease in rheumatoid arthritis.<sup>81</sup> Non-steroidal anti-inflammatory drugs (NSAIDs) are a mainstay of treatment for arthritis, and theoretical models implicate inflammation as key to Alzheimer's disease pathogenesis. Consequently, there has been considerable research interest into the role of NSAIDs in Alzheimer's disease. Substantial evidence from longitudinal observation studies supports a protective role for NSAIDs against Alzheimer's disease.<sup>81,82</sup> NSAIDs appear to delay Alzheimer's disease onset<sup>83</sup> and supporting evidence has demonstrated diminished risk reduction with increasing age.<sup>84</sup> However, clinical trials have failed to demonstrate benefit.<sup>85</sup> A plausible explanation is that studies did not involve intervention at a sufficiently early stage. There is general agreement that sustained treatment of >2 years duration is required<sup>86</sup> and that NSAID ingestion must occur at least several years before the onset of Alzheimer's disease for risk reduction.<sup>82,87</sup>

Most traditional NSAIDs inhibit both COX1 and COX2 enzymes. COX1 is implicated in the activation of microglia in Alzheimer's disease, and the theoretical rationale for the use of NSAIDs targeting this enzyme is supported by epidemiological work.<sup>85,88</sup> Some clinical trials have investigated the newer selective COX 2 inhibitors such as celecoxib, finding no benefit. Results from animal models and epidemiological studies also support a lack of efficacy for selective COX2 inhibitors in reducing Alzheimer's disease risk.<sup>85</sup> However, NSAIDs and associated compounds have a wide range of effects beyond inflammation. Some traditional NSAIDs are also anti- $\beta$ -amyloidogenic, including ibuprofen and indomethacin, most likely acting via Y-secretase.<sup>89</sup> As yet, it is unclear whether NSAIDs exert their perceived preventative effects by modulating inflammatory response, influencing  $\beta$ -amyloid deposition, or through interaction between the two.

There is a need for further prospective prevention trials which could usefully focus on mild cognitive impairment.<sup>90</sup> Unfortunately, multiple challenges face such trials in Alzheimer's disease, as demonstrated in the ADAPT study,<sup>91</sup> which terminated early due to adverse effects. Concerns exist



regarding the safety and tolerability of NSAIDs used at a population level for prevention.<sup>90</sup>

## End of life in Alzheimer's

There are surprisingly few autopsy studies that are focused on physical comorbidity in Alzheimer's disease.<sup>92</sup> Bronchopneumonia is the most frequently cited cause of death, typically accounting for almost half of all deaths.<sup>92–95</sup> This is frequently related to eating problems and probably reflects general frailty.<sup>95</sup> Compared to a matched, non-demented population, vascular-related deaths are more common in Alzheimer's disease, as is co-existent vascular disease,<sup>92,96</sup> while cancer is a much less common cause of death.<sup>92,94</sup> The high incidence of comorbidity at the end of life in Alzheimer's disease poses special medical and ethical dilemmas for clinicians.<sup>97</sup> The medical instinct to treat normally reversible pathology, such as bronchopneumonia, can arguably delay death and potentially prolong suffering in what most clinicians view as a terminal illness.<sup>95</sup> Furthermore, at this stage of their illness, affected individuals are unable to make informed decisions about their preferences. A better understanding of associated comorbidity and underlying pathophysiological mechanisms could help clinicians and patient welfare guardians make the most appropriate decisions in advanced Alzheimer's disease.

## Conclusion

Physical comorbidities make important contributions to cognitive and functional decline in Alzheimer's disease. This has important implications for healthcare providers, as patients will often require access to services and expertise beyond that dealing with the behavioural and psychological manifestations of the condition. Patients are likely to benefit from improved team working between general practitioners, old-age psychiatrists, geriatricians and other specialists. Moreover, non-psychiatric associations of Alzheimer's disease may provide clues to pathological mechanisms of this important condition. In particular, the importance of vascular risk factors and arterial disease is increasingly recognized as important in the pathogenesis of the condition.

## References

1. Zhao Y, Kuo TC, Weir S, Kramer MS, Ash AS. Healthcare costs and utilization for Medicare beneficiaries with Alzheimer's. *BMC Public Health* 2008; **8**:108.
2. Doraiswamy PM, Leon J, Cummings JL, Marin D, Neumann PJ. Prevalence and impact of medical comorbidity in Alzheimer's disease. *J Gerontol Biol A Sci Med Sci* 2002; **57**:M173–7.
3. Masters CL, Simms G, Weinmann NA, Multhaup G, McDonald BL, Beyreuther K. Amyloid plaque core protein in Alzheimer disease and Down syndrome. *Proc Natl Sci USA* 1985; **82**:4245–9.
4. Abe K, Tanzi RE, Kogure K. Selective induction of Kunitz-type protease inhibitor domain containing amyloid precursor protein mRNA after persistent focal ischemia in the rat cerebral cortex. *Neurosci Lett* 1991; **125**:172–4.
5. Tomimoto H, Wakita H, Aikiguchi I, Nakamura S, Kimura J. Temporal profiles of accumulation of amyloid  $\beta$ /A4 protein precursor in the gerbil after graded ischemic stress. *Cereb Blood Flow Metab* 1994; **14**:565–73.
6. Jendroska K, Poewe W, Daniel SE, Pluess J, Iwerssen-Schmidt H, Paulsen J, et al. Ischemic stress induces deposition of amyloid beta immunoreactivity in human brain. *Acta Neuropathol* 1995; **90**:461–6.
7. Launer LJ, Ross GW, Petrovich H, Masaki K, Foley D, White LR, et al. Midlife blood pressure and dementia: the Honolulu Asia Ageing study. *Neurobiol Ageing* 2000; **21**:49–55.
8. Bienias JL, Beckett LA, Bennet DA, Wilson RS, Evans DA. Design of the Chicago Health and ageing Project. *J Alzheimer's Dis* 2003; **5**:349–55.
9. Korf ES, White LR, Scheltens P, Launer LJ. Midlife blood pressure and the risk of hippocampal atrophy: the Honolulu Asia Ageing Study. *Hypertension* 2004; **44**:29–34.
10. Guo LJ, Viitanen M, Fratiglioni L. Low blood pressure and dementia in elderly people. *BMJ* 1996; **312**:805–8.
11. Freitag MH, Peila R, Masaki K, Petrovich H, Ross GW, White LR, et al. Midlife pulse pressure and the incidence of dementia: the Honolulu Asia Ageing Study. *Stroke* 2006; **37**:33–7.
12. Kivipelto M, Helkala EL, Laasko MP, Hanninen T, Hallikainen M, Alhainen K, et al. Midlife vascular factors and Alzheimer's disease in later life: longitudinal population based study. *BMJ* 2001; **322**:1447–51.
13. Skoog I, Lernfelt B, Landahl S, Palmertz B, Andreasson LA, Nilsson L, et al. 15-year longitudinal study of blood pressure and dementia. *Lancet* 1996; **347**:1141–5.
14. Girodard H, Iadecola C. Neurovascular coupling in the normal brain and in hypertension, stroke, and Alzheimer's disease. *J Appl Physiol* 2006; **100**:328–35.
15. Petrovich H, White LR, Izmirilian G, Ross GW, Havlik RJ, Markesbery W, et al. Midlife blood pressure and neuritic plaques, neurofibrillary tangles and brain weight at death: the Honolulu Asia Ageing study. *Neurobiol Ageing* 2000; **21**:57–62.
16. Webster NJ, Green KN, Peers C, Vaughan PF. Altered processing of human amyloid precursor protein in human neuroblastoma SH-SY5Y. *J Neurochem* 2002; **83**:1262–71.
17. Perlmutter LC, Hakami MK, Hodson-Harrington C, Ginsberg J, Katz J, Singer DE, et al. Decreased cognitive function in ageing non-insulin-dependant diabetic patients. *Am J Med* 1984; **77**:1043–8.
18. Mooradian AD, Perryman K, Fitten J, Kavonian GD, Morley JE. Cortical function in elderly non-insulin dependant

- diabetic patients: behavioural and electrophysiological studies. *Arch Intern Med* 1988; **148**:2369–72.
19. U'Ren RC, Riddle MC, Lezak MD, Bennington-Davis MI. The mental efficiency of the elderly person with type II diabetes mellitus. *J Am Geriatric Soc* 1990; **38**:505–10.
  20. Reaven GM, Thompson LW, Nahum D, Haskins E. Relationship between hyperglycaemia and cognitive function in older NIDDM patients. *Diabetes Care* 1990; **13**:16–21.
  21. Leibson CL, Rocca WA, Hanson VA, Cha R, Kokmen E, O'Brien PC, et al. Risk of dementia among persons with diabetes mellitus: a population based cohort study. *Am J Epidemiol* 2007; **145**:301–8.
  22. Ott A, Stolk RP, Hofman A, van Harskamp F, Grobbee DE, Breteler MM. Association of diabetes mellitus and dementia: the Rotterdam Study. *Diabetologia* 1996; **39**:1392–7.
  23. Craft S, Dagogo-Jack SE, Wiethop BV, Murphy C, Nevins RT, Fleischman S, et al. Effects of hyperglycaemia on memory and hormone levels in dementia of the Alzheimer's type: a longitudinal study. *Behav Neurosci* 1993; **107**:926–40.
  24. Menielly GS, Hill A. Alterations in glucose metabolism in patients with Alzheimer's disease. *J Am Geriatric Soc* 1993; **41**:710–4.
  25. Ott A, Stolk RP, Van Harskamp F, Pols HA, Hofman A, Breteler MM. Diabetes mellitus and the risk of dementia: the Rotterdam Study. *Neurology* 1999; **53**:1937–42.
  26. Araki Y, Noumura M, Tanaka M, Yamamoto H, Yamamoto T, Tsukaguchi I, et al. MRI of the brain in diabetes mellitus. *Neuroradiology* 1994; **36**:101–3.
  27. den Heijer T, Veemer SE, van Dijk EJ, Prins ND, Koudstaal PJ, Hofman A, et al. Type 2 diabetes and atrophy of medial temporal lobe structures on brain MRI. *Diabetologia* 2003; **46**:1604–10.
  28. Cavalieri M, Ropele S, Petrovic K, Pluta-Fuerst A, Homayoon N, Enzinger C, et al. Metabolic syndrome, brain magnetic resonance imaging, and cognition. *Diabetes Care* 2010; **33**:2489–95.
  29. Craft S. The role of metabolic disorders in Alzheimer's disease and vascular dementia: two roads converged? *Arch Neurol* 2009; **66**:300–5.
  30. Jolivalck CG, Townsend M. Insulin resistance and amyloidogenesis as common molecular foundation for type 2 diabetes and Alzheimer's disease. *Biochem Biophys Acta* 2009; **1792**:482–96.
  31. Young SE, Mainous AG, Carnemolla M. Hyperinsulinemia and cognitive decline in middle aged cohort. *Diabetes Care* 2006; **29**:2688–93.
  32. Fishel MA, Watson GS, Montine SJ, Wang Q, Green PS, Kulstad JJ, et al. Hyperinsulinemia provokes synchronous increase in central inflammation and beta-amyloid in normal adults. *Arch Neurol* 2005; **62**:1539–44.
  33. Li ZG, Zhang WX, Sima AAF. Alzheimer-like change in rat models of spontaneous diabetes. *Diabetes* 2007; **56**:1817–24.
  34. Lobo A, Launer LJ, Fratiglioni L, Andersen K, Di Carlo A, Breteler MM, et al. Prevalence of dementia and major subtypes in Europe: a collaborative study of population-based cohorts. Neurologic Diseases in the Elderly Research Group. *Neurology* 2000; **54**:S4–9.
  35. Anstey KJ, Cherbuin N, Budge M, Young J. Body mass index in midlife and late-life as a risk factor for dementia: a meta-analysis of prospective studies. *Obes Rev* 2011; **12**:e426–37.
  36. Xu WL, Atti AR, Gatz M, Pedersen NL, Johansson B, Fratiglioni L. Midlife overweight and obesity increase late-life dementia risk: a population-based twin study. *Neurology* 2011; **76**:1568–74.
  37. Kovari E, Gold G, Herrmann FR, Canuto A, Hof PR, Michel JP, et al. Cortical microinfarcts and demyelination significantly affect cognition in brain ageing. *Stroke* 2004; **35**:410–4.
  38. Kalmijn S, Foley D, White L, Burchfiel CM, Curb JD, Petrovich H, et al. Metabolic cardiovascular syndrome and risk of dementia in Japanese American elderly men. The Honolulu-Asia ageing study. *Arterioscler Thromb Vasc Biol* 2000; **20**:2255–60.
  39. Grundy SM. Metabolic syndrome: a multiplex cardiovascular risk factor. *J Clin Endocrinol Metab* 2007; **92**:399–404.
  40. Guerreiro RJ, Santana I, Bras JM. Peripheral inflammatory cytokines as biomarkers in Alzheimer's disease and mild cognitive impairment. *Neurodegener Dis* 2007; **4**:406–12.
  41. Bonarek M, Barberger-Gateau P, Letenneur L, Deschamps V, Iron A, Dubroca B, et al. Relationship between cholesterol, apolipoprotein E polymorphism and dementia: a cross-sectional analysis from the PAQUID study. *Neuroepidemiology* 2000; **19**:141–8.
  42. Bjorntorp P, Rosmond R. Neuroendocrine abnormalities in visceral obesity. *Int J Obes Relat Metab Disord* 2000; **24**:S80–5.
  43. Small SA. The longitudinal axis of hippocampal formation: its anatomy, circuitry, and role in cognitive function. *Rev Neurosci* 2002; **13**:183–94.
  44. Ohm TG, Kirca M, Bohl J, Scharnagl H, Gross W, Marz W. Apolipoprotein E polymorphism influences not only cerebral senile plaque load but also Alzheimer like neurofibrillary tangle formation. *Neuroscience* 1995; **66**:583–7.
  45. Beach TG, Wilson JR, Sue LI, Newell A, Poston M, Cisneros R, et al. Circle of Willis atherosclerosis: association with Alzheimer's disease, neuritic plaques and neurofibrillary tangles. *Acta Neuropathol* 2007; **113**:13–21.
  46. Burns MP, Noble WJ, Olm V, Gaynor K, Casey E, LeFrancois J, Wang L, Duff K. Co-localization of cholesterol, apolipoprotein E and fibrillar Abeta in amyloid plaques. *Brain Res Mol Brain Res* 2003; **110**:119–25.
  47. Stefani M, Liguri G. Cholesterol in Alzheimer's disease: unresolved questions. *Curr Alzheimer's Res* 2009; **6**:15–629.
  48. Tan ZS, Vasan RS. Thyroid function and Alzheimer's disease. *J Alzheimer's Dis* 2009; **16**:503–7.
  49. Kalmijn S, Mehta KM, Pols HAP, Hofman A, Drexhage HA, Breteler MMB. Subclinical hyperthyroidism and the risk of dementia. The Rotterdam study. *Clin Endocrinol* 2000; **53**:733–7.
  50. de Jong FJ, Masaki K, Chen H, Remaley AT, Breteler MM, Petrovitch H, et al. Thyroid function, the risk of dementia and neuropathologic changes: the Honolulu-Asia aging study. *Neurobiol Aging* 2009; **30**:600–6.
  51. Tan ZS, Beiser A, Vasan RS, Au R, Auerbach S, Kiel DP, et al. Thyroid function and the risk of Alzheimer's disease: the Framingham Study. *Arch Intern Med* 2008; **168**:1514–20.

52. Ancoli-Israel S, Kripke DF, Klauber MR, Mason WJ, Fell R, Kaplan O. Sleep-disordered breathing in community-dwelling elderly. *Sleep* 1991; **14**:486–95.
53. Hoch CC, Reynolds CF III, Kupfer DJ, Houck PR, Berman SR, Stack JA. Sleep-disordered breathing in normal and pathologic aging. *J Clin Psychiatry* 1986; **47**:499–503.
54. Gehrman PR, Martin JL, Shochat T, Nolan S, Corey-Bloom J, Ancoli-Israel S. Sleep-disordered breathing and agitation in institutionalized adults with Alzheimer disease. *Am J Geriatr Psychiatry* 2003; **11**:426–33.
55. Incalzi RA, Marra C, Salvigni BL, Petrone A, Gemma A, Selvaggio D, et al. Does cognitive dysfunction conform to a distinctive pattern in obstructive sleep apnea syndrome? *J Sleep Res* 2004; **13**:79–86.
56. Ferini-Strambi L, Baietto C, Di Gioia MR, Castaldi P, Castronovo C, Zucconi M, et al. Cognitive dysfunction in patients with obstructive sleep apnea (OSA): partial reversibility after continuous positive airway pressure (CPAP). *Brain Res Bull* 2003; **61**:87–92.
57. Cooke JR, Ancoli-Israel S, Liu L, Loreda JS, Natarajan L, Palmer BS, et al. Continuous positive airway pressure deepens sleep in patients with Alzheimer's disease and obstructive sleep apnea. *Sleep Med* 2009; **10**:1101–6.
58. Ancoli-Israel S, Palmer BW, Cooke JR, Corey-Bloom J, Fiorentino L, Natarajan L, et al. Cognitive effects of treating obstructive sleep apnea in Alzheimer's disease: a randomized controlled study. *J Am Geriatr Soc* 2008; **56**:2076–81.
59. Cooke JR, Ayalon L, Palmer BW, Loreda JS, Corey-Bloom J, Natarajan L, et al. Sustained use of CPAP slows deterioration of cognition, sleep, and mood in patients with Alzheimer's disease and obstructive sleep apnea: a preliminary study. *J Clin Sleep Med* 2009; **5**:305–9.
60. Moraes W, Poyares D, Sukys-Claudino L, Guilleminault C, Tufik S. Donepezil improves obstructive sleep apnea in Alzheimer disease. *Chest* 2008; **133**:677–83.
61. Bliwise DL. Sleep apnea, APOE4 and Alzheimer's disease 20 years and counting? *J Psychosom Res* 2002; **53**:539–46.
62. Bliwise DL, Yesavage JA, Tinklenberg JR, Dement WC. Sleep apnea in Alzheimer's disease. *Neurobiol Aging* 1989; **10**:343–6.
63. Kadotani H, Kadotani T, Young T, Peppard PE, Finn L, Colrain M, et al. Association between apolipoprotein E epsilon4 and sleep-disordered breathing in adults. *JAMA* 2001; **285**:2888–90.
64. Lui L, Stone K, Cauley JA, Hillier T, Yaffe K. Bone loss predicts subsequent cognitive decline in older women: the study of osteoporotic fractures. *J Am Geriatr Soc* 2003; **51**:38–43.
65. Tan ZS, Seshadri S, Beiser A, Zhang Y, Felson D, Hannan MT, et al. Bone mineral density and the risk of Alzheimer disease. *Arch Neurol* 2005; **62**:107–11.
66. Yaffe K, Browner W, Cauley J, Launer L, Harris T. Association between bone mineral density and cognitive decline in older women. *J Am Geriatr Soc* 1999; **47**:1176–82.
67. Weller I, Schatzker J. Hip fractures and Alzheimer's disease in elderly institutionalized Canadians. *Ann Epidemiol* 2004; **14**:319–24.
68. Loskutova N, Honea RA, Vidoni ED, Brooks WM, Burns JM. Bone density and brain atrophy in early Alzheimer's disease. *J Alzheimers Dis* 2009; **18**:777–85.
69. Loskutova N, Honea RA, Brooks WM, Burns JM. Reduced limbic and hypothalamic volumes correlate with bone density in early Alzheimer's disease. *J Alzheimers Dis* 2010; **20**:313–22.
70. Bayer AU, Ferrari F, Erb C. High occurrence rate of glaucoma among patients with Alzheimer's disease. *Eur Neurol* 2002; **47**:165–8.
71. Tamura H, Kawakami H, Kanamoto T, Kato T, Yokoyama T, Sasaki K, et al. High frequency of open-angle glaucoma in Japanese patients with Alzheimer's disease. *J Neurol Sci* 2006; **246**:79–83.
72. McKinnon SJ. Glaucoma: ocular Alzheimer's disease? *Front Biosci* 2003; **8**:s1140–56.
73. Guo L, Salt TE, Luong V, Wood N, Cheung W, Maass A, et al. Targeting amyloid-beta in glaucoma treatment. *Proc Natl Acad Sci USA* 2007; **104**:13444–9.
74. Normando EM, Coxon KM, Guo L, Cordeiro MF. Focus on: amyloid beta. *Exp Eye Res* 2009; **89**:446–7.
75. Wostyn P, Audenaert K, De Deyn PP. Alzheimer's disease and glaucoma: is there a causal relationship? *Br J Ophthalmol* 2009; **93**:1557–9.
76. Hinton DR, Sadun AA, Blanks JC, Miller CA. Optic-nerve degeneration in Alzheimer's disease. *N Engl J Med* 1986; **315**:485–7.
77. Blanks JC, Hinton DR, Sadun AA, Miller CA. Retinal ganglion cell degeneration in Alzheimer's disease. *Brain Res* 1989; **501**:364–72.
78. Wostyn P, Audenaert K, De Deyn PP. Alzheimer's disease-related changes in diseases characterized by elevation of intracranial or intraocular pressure. *Clin Neurol Neurosurg* 2008; **110**:101–9.
79. Roe CM, Fitzpatrick AL, Xiong C, Sieh W, Kuller L, Miller JP, et al. Cancer linked to Alzheimer disease but not vascular dementia. *Neurology* 2010; **74**:106–12.
80. Lu KP. Pinning down cell signaling, cancer and Alzheimer's disease. *Trends Biochem Sci* 2004; **29**:200–9.
81. McGeer PL, Schulzer FRCP, McGeer M, Edith G. Arthritis and anti-inflammatory agents as possible protective factors for Alzheimer's disease: a review of 17 epidemiologic studies. *Neurology* 1996; **47**:425–32.
82. Zandi PP, Breitner JC. Do NSAIDs prevent Alzheimer's disease? And, if so, why? The epidemiological evidence. *Neurobiol Aging* 2001; **22**:811–7.
83. Szekely CA, Thorne JE, Zandi PP, Ek M, Messias E, Breitner JC, et al. Nonsteroidal anti-inflammatory drugs for the prevention of Alzheimer's disease: a systematic review. *Neuroepidemiology* 2004; **23**:159–69.
84. Breitner JCS, Haneuse SJA, Walker R, Dublin S, Crane PK, Gray SL, et al. Risk of dementia and AD with prior exposure to NSAIDs in an elderly community-based cohort. *Neurology* 2009; **72**:1899–905.
85. McGeer PL, McGeer EG. NSAIDs and Alzheimer disease: epidemiological, animal model and clinical studies. *Neurobiol Aging* 2007; **28**:639–47.
86. Etminan M, Gill S, Samii A. Effect of non-steroidal anti-inflammatory drugs on risk of Alzheimer's disease: systematic review and meta-analysis of observational studies. *BMJ* 2003; **327**:128.
87. Hayden KM, Zandi PP, Khachaturian AS, Szekely CA, Fotuhi M, Norton MC, et al. Does NSAID use modify

- cognitive trajectories in the elderly? *Neurology* 2007; **69**:275–82.
88. Vlad SC, Miller DR, Kowall NW, Felson DT. Protective effects of NSAIDs on the development of Alzheimer disease. *Neurology* 2008; **70**:1672–7.
  89. Sastre M, Gentleman SM. NSAIDs: how they work and their prospects as therapeutics in Alzheimer's disease. *Front Aging Neurosci* 2010; **2**:20.
  90. Pasinetti GM. From epidemiology to therapeutic trials with anti-inflammatory drugs in Alzheimer's disease: the role of NSAIDs and cyclooxygenase in beta-amyloidosis and clinical dementia. *J Alzheimers Dis* 2002; **4**:435–45.
  91. ADAPT research group. Martin BK, Szekely C, Brandt J, Piantadosi S, Breitner JC, et al. Cognitive function over time in the Alzheimer's disease anti-inflammatory prevention trial (ADAPT)- results of a randomized, controlled trial of naproxen and celecoxib. *Arch Neurol* 2008; **65**:896–905.
  92. Fu C, Chute DJ, Farag ES, Garakian J, Cummings JL, Vinters HV. Comorbidity in dementia: an autopsy study. *Arch Pathol Lab Med* 2004; **128**:32–8.
  93. Atterns J, Konig C, Huber M, Lintner F, Jellinger KA. Cause of death in demented and non-demented elderly inpatients: an autopsy study of 308 cases. *J Alzheimers Dis* 2005; **8**:57–62.
  94. Brunnstrom HR, Englund EM. Cause of death in patients with dementia disorders. *Eur J Neurol* 2009; **16**:488–92.
  95. Mitchell SL, Teno JM, Kiely DK, Shaffer ML, Jones RN, Prigerson HG, et al. The clinical course of advanced dementia. *N Engl J Med* 2009; **361**:1529–38.
  96. Brosselin P, Dupont N, Bloch J. Mortality with Alzheimer's disease and dementia in France, 2006. *Rev Epidemiol Sante Publique* 2010; **58**:269–76.
  97. Petrini C. The clinical course of advanced dementia. *N Engl J Med* 2010; **362**:364–5;author reply 364–5.