

Increased CO₂ stimulates K/Rb reabsorption mediated by H-K-ATPase in CCD of potassium-restricted rabbit

XIAOMING ZHOU, SUGURU NAKAMURA, SHEN-LING XIA, AND CHARLES S. WINGO

Division of Nephrology, Hypertension and Transplantation, Department of Medicine,
College of Medicine, University of Florida, and Nephrology Section,
Veterans Affairs Medical Center, Gainesville, Florida 32608-1197

Received 25 October 1999; accepted in final form 15 March 2001

Zhou, Xiaoming, Suguru Nakamura, Shen-Ling Xia, and Charles S. Wingo. Increased CO₂ stimulates K/Rb reabsorption mediated by H-K-ATPase in CCD of potassium-restricted rabbit. *Am J Physiol Renal Physiol* 281: F366–F373, 2001.—Apical H-K-ATPase in the cortical collecting duct (CCD) plays an important role in urinary acidification and K reabsorption. Our previous studies demonstrated that an H-K-ATPase mediates, in part, Rb reabsorption in rabbit CCD (Zhou X and Wingo CS. *Am J Physiol Renal Fluid Electrolyte Physiol* 263: F1134–F1141, 1992). The purpose of these experiments was to examine using in vitro microperfused CCD from K-restricted rabbits 1) whether an acute increase in PCO₂ and, presumably, intracellular acidosis stimulate K absorptive flux; and 2) whether this stimulation was dependent on the presence of a functional H-K-ATPase. Rb reabsorption was significantly increased after exposure to 10% CO₂ in CCD, and this effect was persistent for the entire 10% CO₂ period, whereas 10 μM SCH-28080 in the perfusate totally abolished the stimulation of Rb reabsorption by 10% CO₂. After stimulation of Rb reabsorption by 10% CO₂, subsequent addition of 0.1 mM methazolamide, an inhibitor of carbonic anhydrase, failed to affect Rb reabsorption. However, simultaneous exposure to 10% CO₂ and methazolamide prevented the stimulation of Rb reabsorption. Treatment with the intracellular calcium chelator MAPTAM (0.5 μM) inhibited the stimulation of Rb reabsorption by 10% CO₂. Similar inhibition was also observed in the presence of either a calmodulin inhibitor, W-7 (0.5 μM), or colchicine (0.5 mM), an inhibitor of tubulin polymerization. In time control studies, the perfusion time did not significantly affect Rb reabsorption. We conclude the following: 1) stimulation of Rb reabsorption on exposure to 10% CO₂ is dependent on the presence of a functional H-K-ATPase and appears to be regulated in part by the insertion of this enzyme into the apical plasma membrane by exocytosis; 2) insertion of H-K-ATPase requires changes in intracellular pH and needs a basal level of intracellular calcium concentration; and 3) H-K-ATPase insertion occurs by a microtubule-dependent process.

exocytosis; intracellular pH; calcium; calmodulin; microtubules; cortical collecting duct

IN THE RENAL COLLECTING DUCT, three distinct pathways of K/Rb reabsorption have been identified: a Ba-sensitive K-conductive pathway due to Ba-sensitive K channels, an electroneutral pathway due to an H-K-ATPase, and

the paracellular pathway (45). Previous studies have shown that 10% CO₂ stimulates luminal acidification via an apical SCH-28080-sensitive H-K-ATPase under K-replete conditions (44). K restriction increased an apical H-K-ATPase activity of the microperfused collecting duct in rabbit (15, 38). Functional studies have demonstrated that 10% CO₂ increases Rb absorption in the colon, which is mediated by a related K-absorbing pump (27), and an acute increase in ambient PCO₂ in vitro enhances the rate of luminal acidification in the collecting duct (22). The role of ambient PCO₂ in the control of exocytosis has been studied extensively in other acid-transporting epithelia. In particular, the turtle urinary bladder has many transport characteristics similar to those of the collecting duct (4). Acetazolamide has been shown to alkalinize intracellular pH (pH_i) in the turtle urinary bladder (4), and exposure to CO₂ also increases H secretion in this epithelia. This effect of CO₂ on luminal acidification appears dependent on the exocytosis of vesicles containing proton pumps at the apical membrane. Exocytosis may be an important mechanism for regulation of both proton pumps and has been proposed to occur by the following sequence: 1) exposure to CO₂ decreases pH_i, and the rate of pH_i reduction is affected by the presence of carbonic anhydrase; 2) rapid alterations in pH_i increase intracellular free Ca²⁺ concentration ([Ca²⁺]_i), which accelerates the organization of the cytoskeleton; and 3) cytoskeleton rearrangement is dependent on calmodulin and results in the translocation of tubulovesicles from the cytoplasm to the apical membrane (4, 30, 32, 33, 35). These observations suggest that an intracellular acid-base balance affects the activity of this enzyme and prompted us to speculate that increased CO₂ might also stimulate K absorption by the cortical collecting duct (CCD) during K-restricted conditions, which could be attributable to enhanced H-K-ATPase activity. The purpose of this study was to examine whether increased CO₂ induced by exposure to 10% CO₂ affects H-K-ATPase-dependent K absorption of the CCD in animals adapted to a low-K diet (K-restricted animals). In addition, we examined pos-

Address for reprint requests and other correspondence: C. S. Wingo, Nephrology Section, Dept. of Veterans Affairs Medical Ctr., Gainesville, FL 32608-1197 (E-mail: wingocs@ufl.edu).

The costs of publication of this article were defrayed in part by the payment of page charges. The article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

sible intracellular mechanisms that could explain this effect. Rb reabsorption was used as a qualitative measure of K reabsorption, and rabbits were adapted to K-restricted conditions to enhance the component of Rb/K reabsorption mediated by H-K-ATPase.

MATERIALS AND METHODS

In vivo conditions. Female New Zealand White rabbits were maintained on a K-restricted diet (0.25% K, TD 87433, Teklad, Madison, WI) for at least 4 days before experimentation and allowed free access to tap water.

In vitro methods. Standard in vitro microperfusion methods (2) as modified in this laboratory were used (38, 43). Briefly, rabbits were decapitated, the left kidney was quickly removed, and 1- to 2-mm slices were placed in a chilled petri dish containing Ringer bicarbonate solution [(in mM) 145 Na, 5 K, 112 Cl, 25 HCO₃, 1.8 Ca, 2.3 PO₄, 1.0 Mg, 1.0 SO₄, 10 acetate, 8 glucose, and 5 alanine] gassed with 5% CO₂-95% O₂ with 5% vol/vol fetal calf serum.

Dissection proceeded superficially from the corticomedullary junction. Tubules were transferred to a thermostatically controlled chamber (37°C), and the two ends of the tubule were aspirated into the holding pipettes. The perfusing pipette was advanced 100 μm beyond the holding pipette. The composition of the bath solution was identical to that of dissection solution and was gassed with 5% CO₂-95% O₂ (pH = 7.4 ± 0.04) or 10% CO₂-90% O₂ (pH = 7.1 ± 0.03) as appropriate. The bath solution was continuously exchanged at a rate of 0.64 ml/min. The equilibration time between two periods was 30 min unless otherwise indicated. The electrolyte composition of the perfusate was identical to that of the bath and was gassed with 5% CO₂-95% O₂. The perfusate contained 50 μCi of [methoxy-³H]inulin exhaustively dialyzed according to the method of Schafer et al. (28). The flow rate of the perfusate was maintained between 4 and 6 nl/min. Effluent fluid was collected into a constant-volume pipette for measurement of volume and isotopic flux.

Volume flux was determined from timed collections of the effluent fluid by using the equation $J_v = (\text{cpm}_o/\text{cpm}_i - 1) \cdot V_o/L$, where J_v is the net volume absorption in nanoliters per millimeter per minute; cpm_o and cpm_i are the [³H]inulin counts per min per nanoliter in the collected and the perfused fluid, respectively; V_o is the rate of fluid collection in nanoliters per minute; and L is the tubule length in millimeters. In all experiments, the percent recovery of [³H]inulin was >95% or the experimental result was discarded.

K absorptive flux was assessed as the ⁸⁶Rb lumen-to-bath efflux coefficient (K_{Rb} or Rb reabsorption). Isotopic ⁸⁶Rb was used as a tracer marker for measurement of K reabsorption because stimulation of K-ATPase activity with Rb is similar to that observed with K (5), and there is no evidence that the two ions are transported by different pathways (37). ⁸⁶Rb reabsorption was determined by the disappearance of ⁸⁶Rb from the perfusate according to the following equation

$$K_{\text{Rb}} = V_o/L \cdot 2 \cdot (\text{Rb}_i^* - \text{Rb}_o^*)/(\text{Rb}_i^* + \text{Rb}_o^*)$$

where Rb_i^* and Rb_o^* are the ⁸⁶Rb counts per minute per nanoliter in the perfused and collected fluid, respectively. At least three and generally four collections were obtained for measurement of Rb reabsorption. Counts for ³H and ⁸⁶Rb were measured by an LS-7800 liquid scintillation counter (Beckman Instruments, Irvine, CA). The overlap of ⁸⁶Rb counts in the ³H channel was corrected as previously described (43).

The [Ca²⁺]_i of CCD cells was measured with fluo-4 and Fura Red using a confocal fluorescence system (Laser Scan-

ning Module; LSM 510, Zeiss, Thornwood, NY). Tubules were loaded with Ringer solution containing 10 μM fluo-4 acetoxymethyl ester (Molecular Probes, Eugene, OR) by 60-min incubation (either loaded from lumen or from bath), then washed at least 15 min by the Ringer solution. To prevent possible artifacts by the use of single-wavelength fluorescent studies, we used the fluorescent indicator Fura Red (15 μM) to coload cells at the same time, so that a ratio signal could be measured (16). The dyes were dissolved in DMSO, and the final concentration of DMSO in the loading solution was always <0.25% (vol/vol).

Fluorescence measurements were carried out with the fluar ×20 (numerical aperture = 0.75) and plan-neo ×40 (numerical aperture = 0.75) objectives with an inverted microscope (Axiovert 100M, Zeiss). Images were obtained at a rate of 1 image/10 s and stored onto a hard disk as 515 × 512 pixels for further analysis. Windows NT-based Zeiss LSM image-analysis software was used to calculate the fluorescent intensities and plot the time courses.

All chemicals were analytical grade or of the highest available purity. [³H]inulin and ⁸⁶Rb were obtained from New England Nuclear (Boston, MA). SCH-28080 (graciously supplied by Dr. James J. Kaminski, Schering, Bloomfield, NJ) was dissolved in DMSO and applied to the perfusate with a final concentration of 10 μM. The final concentration of DMSO was 0.1% (vol/vol), and this concentration has been previously demonstrated not to affect Rb efflux (44). Methazolamide (American Cyanamide, Pearl River, NY), bis-(2-amino-5-methyl-phenoxy)-ethane-*N,N,N',N'*-tetraacetic acid tetraacetoxymethyl ester (MAPTAM; Sigma, St. Louis, MO), and *N*-(6-aminoethyl)-5-chloro-1-naphthalene-sulfonamide (W-7, Sigma) were also dissolved in DMSO and applied to the bath with the final concentration of 0.1 mM, 0.5 μM, and 0.5 μM, respectively. Colchicine (Sigma) was dissolved in a 154 mM NaCl solution and applied to the bath with a final concentration of 0.5 mM.

Data are expressed as means ± SE. Statistical analyses were performed by ANOVA for repeated measures or paired *t*-test as appropriate. Post hoc comparisons were made by the Ryan-Einot-Gabriel-Welch *F*-test. The null hypothesis was rejected at the 0.05 level of significance.

RESULTS

To examine whether an acute increase in PCO₂ stimulates K absorptive flux, we perfused the CCD with symmetrical Ringer bicarbonate solution gassed with 5% CO₂ (5% CO₂ period) or 10% CO₂ (10% CO₂ period). After measurement of the basal rate of Rb reabsorption (5% CO₂ period), effluent fluid was collected from 30 to 120 min after exposure to 10% CO₂. Exposure to 10% CO₂ decreases pH_i (29, 35), which could provide a greater concentration of protons (or, more precisely, hydronium ions) as a substrate for the H-K-ATPase. The rate of reduction of pH_i is dependent on the activity of carbonic anhydrase that catalyzes the hydration of CO₂ to carbonic acid (8, 10, 20). To evaluate the role of carbonic anhydrase on the stimulation of Rb reabsorption by 10% CO₂, 0.1 mM methazolamide, an inhibitor of this enzyme, was added to the bath from 120 to 180 min after exposure to 10% CO₂. As shown in Fig. 1, exposure to 10% CO₂ significantly stimulated Rb efflux, and this effect was persistent for the entire 10% CO₂ period. The subsequent addition of methazolamide did not significantly inhibit K_{Rb} (72.4 ± 11.8, 5% CO₂




Fig. 1. Effect of 10% CO₂ and 0.1 mM methazolamide in the absence of SCH-28080 on the ⁸⁶Rb lumen-to-bath efflux coefficient (K_{Rb} ; $n = 6$). * $P < 0.05$ compared with the 5% CO₂ period (ANOVA).

period; 121 ± 29.9 , 135 ± 34.3 , and 135 ± 41.3 nm/s at 30–60, 60–90, and 90–120 min after exposure to 10% CO₂, respectively; 133 ± 29.0 nm/s, methazolamide period, $n = 6$). To examine whether the H-K-ATPase contributes to the effect of 10% CO₂ on Rb reabsorption, we repeated the experiments under conditions identical to those described above (Fig. 1) except for the presence of luminal 10 μ M SCH-28080, a specific inhibitor of H-K-ATPase. SCH-28080 totally blocked the stimulation of Rb reabsorption in response to 10% CO₂, and methazolamide had no significant effect on Rb reabsorption under these conditions (90.8 ± 16.5 , 5% CO₂ period; 80.5 ± 12.3 , 83.9 ± 14.3 , and 78.7 ± 10.9 nm/s at 30–60, 60–90, and 90–120 min after exposure to 10% CO₂, respectively; 81.9 ± 13.5 nm/s, methazolamide period, $n = 7$, Fig. 2). Thus H-K-ATPase appears to mediate the stimulatory effect of 10% CO₂ on K_{Rb} . The lack of inhibitory effect of methazolamide on K_{Rb} after exposure to 10% CO₂ implies that carbonic anhydrase is not necessary to maintain the activation of H-K-ATPase by 10% CO₂. To examine whether carbonic anhydrase was required for initiating the stimulation of K_{Rb} by 10% CO₂, we simultaneously exposed the CCD to methazolamide and 10% CO₂ after mea-

surement of the basal rate of K_{Rb} . Under these conditions, 10% CO₂ did not stimulate Rb reabsorption (98.6 ± 14.1 vs. 86.2 ± 16.5 nm/s, $n = 6$, Fig. 3). Time control experiments demonstrated that perfusion time did not significantly affect K_{Rb} (Table 1). These data suggest that carbonic anhydrase is necessary for initiating the activation of H-K-ATPase on exposure to 10% CO₂. To examine whether methazolamide inhibited the basal rate of Rb reabsorption, we perfused the CCD in the presence of 5% CO₂ throughout the experiment and added methazolamide during the second period. As shown in Fig. 4, methazolamide did not significantly inhibit K_{Rb} (60.4 ± 10.1 vs. 59.1 ± 10.6 nm/s, $n = 6$). Because our previous studies demonstrated that H-K-ATPase contributed to the Rb reabsorption when the CCD was perfused in the presence of 5% CO₂ (43), it appears that methazolamide has no significant effect on basal H-K-ATPase activity.

The time frame for the stimulation of Rb efflux on exposure to 10% CO₂ suggests that existing H-K-ATPase pump units contribute to this effect. This could involve activation of previously inactive H-K-ATPase located at the apical plasma membrane or translocation of H-K-ATPase by the insertion of the enzyme to




Fig. 2. Effect of 10% CO₂ and 0.1 mM methazolamide in the presence of SCH-28080 on K_{Rb} ($n = 7$; ANOVA). None of the means of the 10% CO₂ periods were significantly different from those of the 5% CO₂ period.




Fig. 3. Effect of simultaneous exposure to 10% CO₂ and 0.1 mM methazolamide on K_{Rb} [$n = 6$; not significant (NS) by paired t -test].




Fig. 4. Effect of 0.1 mM methazolamide on K_{Rb} . The entire experiment was conducted in the presence of 5% CO₂ ($n = 6$; NS by paired t -test).

the apical membrane of the CCD via exocytosis, or a combination of both mechanisms. To further test the insertion hypothesis, we examined the effects of several agents known to block the insertion cascade on Rb reabsorption after exposure to 10% CO₂. First, the CCD was perfused in the presence of 0.5 μ M MAPTAM, a cell-permeable intracellular Ca²⁺ chelator, throughout the experiment. In this case, 10% CO₂ failed to stimulate Rb reabsorption (89.5 ± 19.5 vs. 85.6 ± 23.6 nm/s, $n = 6$, Fig. 5). To assess whether 10% CO₂ increased [Ca²⁺]_i, we perfused the CCD and measured [Ca²⁺]_i by using a confocal fluorescence system with fluo-4 and Fura Red. [Ca²⁺]_i failed to increase during the exposure to 10% CO₂ (5–15 min; $n = 44$ cells/3 tubules). However, a decrease in [Ca²⁺]_i was observed by adding 5 mM EGTA to the bath solution containing 1.5 mM Ca²⁺. These observations suggest that basal-level [Ca²⁺]_i is critical for 10% CO₂ to stimulate K_{Rb} . Second, to evaluate the role of calmodulin in the stimulation of K_{Rb} by 10% CO₂, the CCD was perfused in the presence of 0.5 μ M W-7, an inhibitor of calmodulin, throughout the experiment. W-7 completely abolished the stimulation of 10% CO₂ on Rb reabsorption. In fact, Rb reabsorption was significantly reduced during the 10% CO₂ period (89.3 ± 14.0 vs. 65.1 ± 13.0 nm/s, $P < 0.05$, $n = 6$, Fig. 6). These observations indicate a critical role for calmodulin in the activation of K_{Rb} by 10% CO₂. Third, to examine whether microtubules were involved in the effect of 10% CO₂, the CCD was simultaneously exposed to 10% CO₂ and 0.5 mM colchicine, an inhibitor of tubulin

polymerization. Colchicine totally blocked the stimulatory effect of 10% CO₂ on Rb reabsorption (85.0 ± 15.0 vs. 81.3 ± 13.6 nm/s, $n = 6$, Fig. 7). These data indicate a requirement for an intact cytoskeletal system in the activation of K_{Rb} by 10% CO₂. In contrast, after stimulation of K_{Rb} by 10% CO₂, subsequent addition of colchicine had no inhibitory effect on Rb reabsorption (72.5 ± 7.9 nm/s, 5% CO₂ period; 97.3 ± 10.1 nm/s, 10% CO₂ period; and 110 ± 13.2 nm/s, 10% CO₂+colchicine period, $n = 6$, Fig. 8).

DISCUSSION

In renal collecting ducts, K reabsorption occurs in the intercalated cells of CCD and outer medullary collecting duct (OMCD) and is mediated by H-K-ATPase pumps in the luminal membrane that reabsorb K and secrete H. The activity of these pumps is increased by hypokalemia (15, 25, 38). The renal response to K depletion (KD) is sufficiently effective that

Table 1. Effect of perfusion time

	Time After Decapitation, min			
	Basal, 100–150	150–200	100–250	150–300
K_{Rb} , nm/s	82.1 ± 13.0	83.6 ± 11.7	106 ± 8.8	100 ± 15.1

Values are means \pm SE. K_{Rb} , ⁸⁶Rb lumen-to-bath efflux coefficient ($n = 5$). None of the means are significantly different from the basal values.




Fig. 5. Effect of 10% CO₂ in the presence of 0.5 μ M bis-(2-amino-5-methyl-phenoxy)-ethane- N,N,N',N' -tetraacetic acid tetraacetoxymethyl ester (MAPTAM) throughout the experiment on K_{Rb} (NS by paired t -test).




Fig. 6. Effect of 10% CO₂ on K_{Rb} in the presence of 0.5 μ M *N*-(6-aminoethyl)-5-chloro-L-naphthalene-sulfonamide (W-7) throughout the experiment (paired *t*-test).

a low-K diet will not lead to significant K losses. Microperfusion studies have demonstrated a K-dependent, SCH-28080-sensitive, acid-secreting pathway in the collecting ducts of normal and KD animals (15, 23–25, 36). Previous functional studies have demonstrated H-K-ATPase sensitivity to SCH-28080 in KD rats and rabbits (1, 15, 23, 25). On the basis of the inhibitory profile of H-K-ATPase activity in KD animals by SCH-28080, it has been presumed that the gastric H-K-ATPase is the isoform responsible for increased activity (15, 41). However, colonic H-K-ATPase (HK α_2) is induced and mediates increased bicarbonate reabsorption in OMCD and inner medullary collecting duct (IMCD) of KD rats (23, 25). Recent studies show that, in KD, an H-K-ATPase activity (as assayed by ATP hydrolysis) is induced in OMCD and CCD and is sensitive to both ouabain and SCH-28080 (1). HK α_2 is sensitive to both ouabain and SCH-28080 in KD (1, 25) and is likely responsible for increased bicarbonate and K reabsorption in collecting ducts. As such, this iso-




Fig. 7. Effect of simultaneous exposure to 10% CO₂ and 0.5 mM colchicine on K_{Rb} (NS by paired *t*-test).




Fig. 8. Effect of 0.5 mM colchicine on K_{Rb} after prior stimulation of K_{Rb} by 10% CO₂. **P* < 0.05 compared with the 5% CO₂ period (ANOVA).

form appears to play an important role in the maintenance of metabolic alkalosis in KD. Our results provide direct evidence that increased CO₂ stimulates K-absorptive flux, assessed as the K_{Rb} or Rb efflux, in the CCD of rabbits under K-restricted conditions. Moreover, SCH-28080 inhibited the stimulation of Rb efflux by 10% CO₂, suggesting that this enhanced K_{Rb} is mediated by an H-K-ATPase. The rapid response to 10% CO₂ and the effects of various maneuvers designed to inhibit exocytosis suggest that the stimulus increases the number of functional H-K-ATPase units in the apical membrane. In the gastric gland, exocytosis of H-K-ATPase to the canalicular (apical) membrane occurs in response to several stimuli (11, 32). We propose that the stimulation of Rb reabsorption on exposure to 10% CO₂ is dependent on the exocytotic insertion of H-K-ATPase into the apical membrane of the collecting duct. Three different lines of evidence are consistent with this hypothesis. First, the present studies show that maneuvers designed to prevent exocytosis also block the stimulatory effect of 10% CO₂ on Rb reabsorption (see Figs. 5–7). Second, H-K-ATPase has been localized by immunocytochemistry to the intercalated cell in the CCD and OMCD (40), and inhibition of this enzyme by SCH-28080 blocks the effect of 10% CO₂ on K_{Rb} (42). Third, alterations in ambient PCO₂ stimulate the translocation of tubulovesicles to the luminal membrane in the CCD (22, 29). Respiratory acidosis increases the surface density of the apical membrane of intercalated cells concomitantly with a decrease in the number of tubulovesicular profiles in the apical region, but no such finding occurred in the principal cell (18). These findings have been taken as evidence that increased CO₂ stimulates the fusion of tubulovesicular profiles to the apical membrane (18). In addition, experiments using fluorescent dextran demonstrate that CO₂ stimulated the exocytotic fusion of vesicles with the apical membrane in the intercalated cell of rabbit collecting duct (29). Taken together,

these studies indicate that 10% CO₂ stimulates exocytosis both in vivo and in vitro.

Functional studies have demonstrated that an increase in CO₂ causes a decrease in pH_i in ventricular myocytes (14) and in the perfused CCD (I. D. Weiner, personal communication). The decrease in pH_i stimulates the exocytotic insertion of H pumps into the apical membrane of the turtle urinary bladder tissue (4, 30, 35). Consistent with the present observations, simultaneous exposure to 10% CO₂ and methazolamide inhibited the stimulatory effect of 10% CO₂ on K_{Rb} (Fig. 3), whereas after 10% CO₂ had increased Rb reabsorption, the subsequent addition of methazolamide had no detectable effect on K_{Rb} (Fig. 1). Previous studies have demonstrated that methazolamide dramatically reduced the initial rate of acidification produced by the carbonic anhydrase-catalyzed hydration of CO₂ (26). This may explain the lack of the late effect on Rb efflux by the addition of methazolamide after the CCD was acidified. In addition, our previous studies suggest that the H-K-ATPase contributes to Rb reabsorption in the basal (5% CO₂) state (43), and our present studies demonstrate that methazolamide does not significantly affect K_{Rb} when the CCD is perfused in the presence of 5% CO₂ throughout the experiment (Fig. 4). These data support the hypothesis that, after H-K-ATPase is inserted into the apical membrane, methazolamide does not significantly inhibit the activity of this enzyme. Therefore, methazolamide inhibition of H-K-ATPase appears to be indirect, and this effect is mediated by inhibition of exocytosis, which is pH dependent. The proton or hydronium ion source for the H-K-ATPase after the administration of methazolamide is an additional issue that is not addressed by these studies. The hydrolysis of ATP may supply protons for the pump, as suggested for the gastric H-K-ATPase (17). Alternatively, the noncatalyzed hydration of CO₂ (20) may in part provide protons for apical secretion.

[Ca²⁺]_i concentration is responsible for a variety of cellular events. Many Ca²⁺ responses depend on protein phosphorylation or dephosphorylation through the reactions with calmodulin. For example, calmodulin regulates Ca²⁺-dependent microtubule assembly and disassembly (19). Exposure to CO₂ in the turtle urinary bladder appears to stimulate the insertion of proton pumps into the apical membrane (4). When alterations in [Ca²⁺]_i are minimized (4, 22, 35) or the function of calmodulin is antagonized (22), exocytosis and the enhancement of proton secretion are mitigated. Compatible with these observations, two maneuvers that should affect [Ca²⁺]_i or calmodulin activity also inhibited the enhancement of H-K-ATPase-dependent Rb reabsorption on exposure to 10% CO₂ (Figs. 5 and 6). In the present studies, we observed that MAPTAM prevented an adaptive increase in Rb efflux in response to 10% CO₂. Microperfusion experiments in the OMCD have demonstrated that 1,2-bis(2-aminophenoxy) ethane-*N,N,N',N'*-tetraacetic acid (BAPTA; analogous to MAPTAM, an intracellular Ca²⁺ chelator) substantially attenuated an adaptive increase in H⁺ secretion in response to in

vitro acidosis (34). On the basis of our results that [Ca²⁺]_i failed to increase under 10% CO₂, we speculate that the inhibitory effect of MAPTAM is related to a decrease in basal intracellular calcium. However, the failure to detect the increase of [Ca²⁺]_i under the 10% CO₂ condition is consistent with the functional studies in rat ventricular myocytes (14). Studies in vascular smooth muscle cells have shown that intracellular acidification is associated with changes in free cytosolic Ca²⁺ via a Ca²⁺-ATPase (7). These observations suggest that increased Ca²⁺ extrusion from the cell may be another reason for the lack of change in [Ca²⁺]_i when Pco₂ is increased. A significant decrease in Rb reabsorption was observed during the 10% CO₂ period when the CCD was perfused in the presence of W-7. We do not exclude the possibility that the effect of W-7 may be due to an action on the basolateral K exit pathway or due to inhibition of the apical H-K-ATPase. The mechanism(s) for the decrease in K_{Rb} remains to be elucidated.

Exocytosis is, in large part, dependent on the organization of the cytoskeleton, which includes microtubules (3, 11, 22). Therefore, we examined conditions that were expected to inhibit microtubule function at two different stages. First, we simultaneously exposed the CCD to 10% CO₂ and colchicine. Under these conditions, we observed no stimulation of Rb reabsorption (Fig. 7). Second, we further pursued this question by stimulating Rb reabsorption with 10% CO₂ and subsequently examining the effect of colchicine. In this case, colchicine did not reduce Rb reabsorption (Fig. 8). These results suggest that exocytosis of the H-K-ATPase is dependent on the intact function of microtubules. After the H-K-ATPase inserts into the membrane, the present data suggest that colchicine does not inhibit the enzyme. Alternatively, exocytosis may be necessary for the insertion into the apical membrane of a protein that is essential for the functional activity of the H-K-ATPase.

Exposure to 10% CO₂, which stimulates exocytosis of H-K-ATPase, may not be a phenomenon unique to animals adapted to a K-restricted diet. McKinney and Davidson (22) have previously shown that the enhancement of acidification by the collecting duct on exposure to 15% CO₂ involves the intact function of the cell cytoskeleton in normal (K-replete) animals. In CCD of K-replete animals, the principal cells possess pH-sensitive K conductance at both the apical and basolateral membrane, whereas intercalated cells possess H-K-ATPase but small K conductance. CO₂ (10%) could be predicted to decrease Rb reabsorption via K conductance but increase Rb reabsorption that is mediated by an H-K-ATPase. Our previous findings that 10% CO₂ failed to increase Rb reabsorption during K repletion suggest that an apical K exit mechanism is the predominant pathway for K flux under these conditions and that 10% CO₂ stimulates luminal acidification via an H-K-ATPase-dependent process in K-replete animals (44). Clearly, the response to 10% CO₂ during K repletion differs from that during K restriction.

In summary, these studies demonstrate that 10% CO₂ stimulates Rb reabsorption, and this stimulation was totally abolished by SCH-28080. The rapid response to 10% CO₂ implies that this maneuver increases the functional units of H-K-ATPase at the apical membrane or the activity of existing pump units. The subsequent addition of methazolamide after exposure to 10% CO₂ had no significant effect on K_{Rb} , whereas simultaneous exposure of the tubules to 10% CO₂ and methazolamide prevented the enhancement of Rb reabsorption by 10% CO₂, suggesting that carbonic anhydrase is not necessary for maintaining activation of H-K-ATPase by 10% CO₂ but is necessary for initiating activation of this enzyme. MAPTAM, W-7, and colchicine inhibited the stimulation of Rb reabsorption after exposure to 10% CO₂, suggesting that the effect of 10% CO₂ on Rb reabsorption requires at least a basal level of $[Ca^{2+}]_i$ as well as the intact function of calmodulin and microtubules. These studies suggest that increased CO₂ activates an SCH-28080-sensitive H-K-ATPase in the CCD of K-restricted rabbits.

These studies were supported by the Medical Research Service of the Department of Veterans Affairs and National Institute of Diabetes and Digestive and Kidney Diseases Grant R01-DK-49750.

Present address of X. Zhou: Dept. of Medicine, Uniformed Services Univ. of the Health Sciences, 4301 Jones Bridge Rd., Bethesda, MD 20814-4799.

REFERENCES

1. Buffin-Meyer B, Younes-Ibrahim M, Barlet-Bas C, Cheval L, and Doucet A. K depletion modifies the properties of SCH-28080-sensitive K-ATPase in rat collecting duct. *Am J Physiol Renal Physiol* 272: F124–F131, 1997.
2. Burg M, Grantham J, Abramson M, and Orloff J. Preparation and study of fragments of single rabbit nephrons. *Am J Physiol* 210: 1293–1298, 1966.
3. Burgess TL and Kelly RB. Constitutive and regulated secretion of proteins. *Annu Rev Cell Biol* 3: 243–293, 1987.
4. Cannon C, van Adelsberg J, Kelly S, and Al-Awqati Q. Carbon-dioxide-induced exocytotic insertion of H⁺ pumps in turtle-bladder luminal membrane: role of cell pH and calcium. *Nature* 314: 443–446, 1985.
5. Cheval L, Barlet-Bas C, Khadouri C, Feraille E, Marsy S, and Doucet A. K⁺-ATPase-mediated Rb⁺ transport in rat collecting tubule: modulation during K⁺ deprivation. *Am J Physiol Renal Fluid Electrolyte Physiol* 260: F800–F805, 1991.
6. Curran KA, Hebert MJ, Cain BD, and Wingo CS. Evidence for the presence of a K-dependent acidifying adenosine triphosphatase in the rabbit renal medulla. *Kidney Int* 42: 1093–1098, 1992.
7. Daugirdas JT, Arrieta J, Ye M, Flores G, Battle DC. Intracellular acidification associated with changes in free cytosolic calcium. Evidence for Ca²⁺/H⁺ exchange via a plasma membrane Ca²⁺-ATPase in vascular smooth muscle cells. *J Clin Invest* 95: 1480–1489, 1995.
8. Dobyan DC and Bulger RE. Renal carbonic anhydrase. *Am J Physiol Renal Fluid Electrolyte Physiol* 243: F311–F324, 1982.
9. Doucet A and Marsy S. Characterization of K-ATPase activity in distal nephron: stimulation by potassium depletion. *Am J Physiol Renal Fluid Electrolyte Physiol* 253: F418–F423, 1987.
10. DuBose TD Jr and Lucci MS. Effect of carbonic anhydrase inhibition on superficial and deep nephron bicarbonate reabsorption in the rat. *J Clin Invest* 71: 55–65, 1983.
11. Forte JG, Black JA, Fortes TM, Machen TE, and Wolosin JM. Ultrastructural changes related to functional activity in gastric oxyntic cells. *Am J Physiol Gastrointest Liver Physiol* 241: G349–G358, 1981.
12. Garg LC and Narang N. Ouabain-insensitive K-adenosine triphosphatase in distal nephron segments of the rabbits. *J Clin Invest* 81: 1204–1208, 1988.
13. Gifford JD, Rome L, and Galla JH. H⁺-K⁺-ATPase activity in rat collecting duct segments. *Am J Physiol Renal Fluid Electrolyte Physiol* 262: F692–F695, 1992.
14. Hayashi H, Miyata H, Noda N, Kobayashi A, Hirano M, Kawai T, Yamazaki N. Intracellular Ca²⁺ concentration and pH_i during metabolic inhibition. *Am J Physiol Cell Physiol* 262: C628–C634, 1992.
15. Kuwahara M, Fu WJ, and Marumo F. Functional activity of H-K-ATPase in individual cells of OMCD: localization and effect of K⁺ depletion. *Am J Physiol Renal Fluid Electrolyte Physiol* 270: F116–F122, 1996.
16. Lipp P and Niggli E. Ratiometric confocal Ca²⁺-measurements with visible wavelength indicators in isolated cardiac myocytes. *Cell Calcium* 14: 359–372, 1993.
17. Ljungstrom M, Vega FV, and Mardh S. Effects of pH on the interaction of ligands with the (H⁺+K⁺)-ATPase purified from pig gastric mucosa. *Biochim Biophys Acta* 769: 220–230, 1984.
18. Madsen KM and Tisher CC. Cellular response to acute respiratory acidosis in rat medullary collecting duct. *Am J Physiol Renal Fluid Electrolyte Physiol* 245: F670–F679, 1983.
19. Marcum JM, Dedman JR, Brinkley BR, and Means AR. Control of microtubule assembly-disassembly by calcium dependent regulator protein. *Proc Natl Acad Sci USA* 75: 3771–3775, 1978.
20. Maren TH. Chemistry of the renal reabsorption of bicarbonate. *Can J Physiol Pharmacol* 52: 1041–1050, 1974.
21. Mattsson JP, Vaananen K, Wallmark B, and Lorentzon P. Omeprazole and bafilomycin, two proton pump inhibitors: differentiation of their effects on gastric, kidney and bone H⁺-translocating ATPase. *Biochim Biophys Acta* 1065: 261–268, 1991.
22. McKinney TD and Davidson KK. Effects of respiratory acidosis on HCO₃⁻ transport by rabbit collecting duct. *Am J Physiol Renal Fluid Electrolyte Physiol* 255: F656–F665, 1988.
23. Nakamura S, Amlal H, Galla J, and Soleimani M. Colonic H⁺-K⁺-ATPase is induced and mediates increased HCO₃⁻ reabsorption in inner medullary collecting duct in potassium depletion. *Kidney Int* 54: 1233–1239, 1998.
24. Nakamura S, Amlal H, Schultheis PJ, Galla J, Shull GE, and Soleimani M. HCO₃⁻ reabsorption in renal collecting duct of NHE-3-deficient mouse: a compensatory response. *Am J Physiol Renal Physiol* 276: F914–F921, 1999.
25. Nakamura S, Wang Z, Galla J, and Soleimani M. K⁺ depletion increases HCO₃⁻ reabsorption in OMCD by activation of colonic H⁺-K⁺-ATPase. *Am J Physiol Renal Physiol* 274: F687–F692, 1998.
26. Newman EA. A physiological measure of carbonic anhydrase in Muller cells. *Glia* 11: 291–299, 1994.
27. Perrone RD and McBride DE. Aldosterone and PCO₂ enhance rubidium absorption in rat distal colon. *Am J Physiol Gastrointest Liver Physiol* 254: G898–G906, 1988.
28. Schafer JA, Troutman SL, and Andreoli TE. Volume reabsorption, transepithelial potential differences, and ionic permeability properties in mammalian superficial proximal straight tubules. *J Gen Physiol* 64: 582–607, 1974.
29. Schwartz GJ and Al-Awqati Q. Carbon dioxide causes exocytosis of vesicles containing H⁺ pumps in isolated perfused proximal and collecting tubules. *J Clin Invest* 75: 1638–1644, 1985.
30. Schwartz GJ and Al-Awqati Q. Regulation of transepithelial H⁺ transport by exocytosis and endocytosis. *Annu Rev Physiol* 48: 153–161, 1986.
31. Smolka A, Helander HF, and Sachs G. Monoclonal antibodies against gastric H⁺ K⁺ ATPase. *Am J Physiol Gastrointest Liver Physiol* 245: G589–G596, 1983.
32. Stetson DL and Steinmetz PR. Role of membrane fusion in CO₂ stimulation of proton secretion by turtle bladder. *Am J Physiol Cell Physiol* 245: C113–C120, 1983.
33. Stetson DL and Steinmetz PR. Correlation between apical intramembrane particles and H⁺ secretion rates during CO₂ stimulation in turtle bladder. *Pflügers Arch* 407: 580–584, 1986.

34. **Tsuruoka S and Schwartz GJ.** Adaptation of the outer medullary collecting duct to metabolic acidosis in vitro. *Am J Physiol Renal Physiol* 275: F982–F990, 1998.
35. **Van Adelsberg J and Al-Awqati Q.** Regulation of cell pH by Ca²⁺-mediated exocytotic insertion of H⁺-ATPase. *J Cell Biol* 102: 1638–1645, 1986.
36. **Wall SM, Truong AV, and DuBose TD Jr.** H⁺-K⁺-ATPase mediates net acid secretion in rat terminal inner medullary collecting duct. *Am J Physiol Renal Fluid Electrolyte Physiol* 271: F1037–F1044, 1996.
37. **Warden DH, Hayashi M, Schuster VL, and Stokes JB.** K⁺ and Rb⁺ transport by rabbit CCD: Rb⁺ reduces K⁺ conductance and Na transport. *Am J Physiol Renal Fluid Electrolyte Physiol* 257: F43–F52, 1989.
38. **Wingo CS.** Active proton secretion and potassium absorption in the rabbit outer medullary collecting duct: functional evidence for proton-potassium-activated adenosine triphosphatase. *J Clin Invest* 84: 361–365, 1989.
39. **Wingo CS and Armitage FE.** Rubidium absorption and proton secretion by rabbit outer medullary collecting duct via H-K-ATPase. *Am J Physiol Renal Fluid Electrolyte Physiol* 263: F849–F857, 1992.
40. **Wingo CS, Madsen KM, Smolka A, and Tisher CC.** H-K-ATPase immunoreactivity in cortical and outer medullary collecting duct. *Kidney Int* 38: 985–990, 1990.
41. **Wingo CS and Smolka A.** Function and structure of H-K-ATPase in the kidney. *Am J Physiol Renal Fluid Electrolyte Physiol* 269: F1–F16, 1995.
42. **Zhou X, Lynch IJ, Xia S-L, and Wingo CS.** Activation of H⁺-K⁺-ATPase by CO₂ requires a basolateral Ba²⁺-sensitive pathway during K restriction. *Am J Physiol Renal Physiol* 279: F153–F160, 2000.
43. **Zhou X and Wingo CS.** H-K-ATPase enhancement of Rb efflux by the cortical collecting duct. *Am J Physiol Renal Fluid Electrolyte Physiol* 263: F43–F48, 1992.
44. **Zhou X and Wingo CS.** Stimulation of total CO₂ flux by 10% CO₂ flux in rabbit CCD: role of an apical Sch-28080- and Ba-sensitive mechanism. *Am J Physiol Renal Fluid Electrolyte Physiol* 267: F114–F120, 1994.
45. **Zhou X and Wingo CS.** Mechanisms of rubidium permeation by rabbit cortical collecting duct during potassium restriction. *Am J Physiol Renal Fluid Electrolyte Physiol* 263: F1134–F1141, 1992.

