

Effects of genistein, resveratrol, and quercetin on steroidogenesis and proliferation of MA-10 mouse Leydig tumor cells

Yu-Chyu Chen^{1,2}, Madan L Nagpal^{1,2}, Douglas M Stocco³ and Tu Lin^{1,2}

¹Medical and Research Services, WJB Dorn Veterans Medical Center, Columbia, South Carolina 29201, USA

²Department of Medicine, Medical Library Building, Suite 316, School of Medicine, University of South Carolina School of Medicine, Columbia, South Carolina 29208, USA

³Department of Cell Biology and Biochemistry, Texas Tech University Health Sciences Center, Lubbock, Texas 79430, USA

(Requests for offprints should be addressed to T Lin at Department of Medicine, University of South Carolina; Email: lin@med.sc.edu)

Abstract

This study was performed to compare the effects of three well-known phytoestrogens such as genistein, resveratrol, and quercetin on steroidogenesis in MA-10 mouse tumor Leydig cells. Addition of genistein or resveratrol to MA-10 cells resulted in decreases in the cAMP-stimulated progesterone secretion, but quercetin had an opposite response. Steroidogenic acute regulatory (StAR) mRNA expression and StAR promoter activity in transiently transfected MA-10 cells were significantly reduced by genistein or resveratrol, but increased by quercetin. Genistein was found to inhibit MA-10 cell proliferation, while resveratrol and quercetin had no effect. Quercetin-induced increase in cAMP-stimulated progesterone secretion was reversed by ICI 182,780, an estrogen receptor (ER) antagonist. However, ICI 182,780 had no effect on cAMP plus quercetin-stimulated StAR promoter activity. To examine whether non-ER factors are associated with quercetin-stimulated progesterone production, we

treated MA-10 cells with EGTA to deprive them of extracellular Ca^{2+} . We found that EGTA inhibited quercetin- plus cAMP-stimulated progesterone secretion and StAR promoter activity. Blocking of Ca^{2+} influx through L- or T-type voltage-gated Ca^{2+} channels with verapamil or nifedipine respectively, attenuated quercetin-stimulated progesterone secretion, while they had no effect on quercetin- plus cAMP-stimulated StAR promoter activity. Blocking of intracellular Ca^{2+} efflux by sodium orthovanadate, a Ca^{2+} - pump inhibitor, blocked quercetin- plus cAMP-stimulated progesterone secretion and StAR promoter activity in MA-10 cells. Finally, EGTA or vanadate reduced quercetin and cAMP-increased in StAR mRNA expression in MA-10 cells, while ICI 182,780 had no effect. Taken together, these results indicate that phytoestrogens have differential effects on steroidogenesis in MA-10 cells.

Journal of Endocrinology (2007) **192**, 527–537

Introduction

Converging lines of evidence suggested that estrogen acts as an important paracrine factor to modulate testicular functions (O'Donnell *et al.* 2001). Estrogen receptor (ER) α is present in the Leydig cells and ER β is detected exclusively in the Sertoli cells (Saunders *et al.* 1998, Pelletier *et al.* 2000). Estradiol treatment suppressed progesterone synthesis (Freeman 1985), but stimulated proliferation of mouse Leydig tumor cells (Sato *et al.* 1987, DuMond *et al.* 2001). Furthermore, mice with a genetic disruption of ER α (α ERKO) appeared infertile, had lower sperm counts and elevated serum testosterone levels (Eddy *et al.* 1996), and were not responsive to the estrogen benzoate-mediated decreases in both serum luteinizing hormone (LH) and androgen levels (Akingbemi *et al.* 2003). In addition, cytochrome P450 17 α -hydroxylase (C17 α), 17 β -hydroxysteroid dehydrogenase (17 β -HSD), and StAR mRNAs were higher in α ERKO mice than in wild-type mice (Akingbemi *et al.* 2003). This

finding suggested that estrogen and ER-mediated signals regulate Leydig cell steroidogenesis.

Phytoestrogens such as genistein, resveratrol, and quercetin are widely distributed in human and animal diet and have chemopreventive properties against estrogen-responsive diseases, including inhibition of tumor cell growth (Setchell & Cassidy 1999, Miodini *et al.* 1999, Mitchell *et al.* 1999, Xing *et al.* 2001), lowering serum cholesterol, and prevention of bone loss in rodents (Mizutani *et al.* 2000, Nakajima *et al.* 2001, Wattel *et al.* 2003). Genistein is the most potent estrogenic compound in soy and soy products. Resveratrol is a potent antioxidant in grapes and red wines. Quercetin is found in apples, onions, and vegetables. Importantly, the exposure of young animals to phytoestrogens has potential impact on male and female gonadal functions. For example, a dietary (Weber *et al.* 2001, Wisniewski *et al.* 2003, Svechnikov *et al.* 2005) or s.c. (Ohno *et al.* 2003) administration of male rats with genistein resulted in lower serum testosterone levels and demasculinization. Exposure of resveratrol to female rats led to

the development of ovarian hypertrophy and to shorter estrous cycle (Henry & Witt 2002). All of these phenomena are believed to be due to the perturbation on gonadal steroidogenesis by the phytoestrogens. In fact, steroidogenic enzymes such as cytochrome P450 side-chain cleavage complex (P450_{sc}) and 17 β -HSD in the gonad and adrenal (Le Bail 2002, Ohno *et al.* 2002, Supornsilchai *et al.* 2005, Svechnikov *et al.* 2005) were altered by phytoestrogen treatments. As yet, reports for phytoestrogen-mediated effects on steroidogenic acute regulatory (StAR) protein expression are limited. Therefore, we focused on phytoestrogen-mediated effects on *StAR* gene expression and steroidogenesis.

In the gonad, the initial step of steroid synthesis depends on the delivery of cholesterol from cytosol into the mitochondria by the StAR protein (Clark *et al.* 1994, 1997), which is the rate-limiting step in steroidogenesis. The expression of StAR is especially abundant in testis, ovary, and adrenal glands (Sugawara *et al.* 1995, Clark *et al.* 1997), and its importance in steroidogenesis was clearly shown by the virtual absence of steroid synthesis in StAR knockout mice (Caron *et al.* 1997b) and in patients with congenital lipoid adrenal hyperplasia caused by mutations in the *StAR* gene (Lin *et al.* 1995, Bose *et al.* 1996). The expression of the *StAR* gene is regulated by numerous factors (Manna & Stocco 2005) and have been shown to be suppressed by 17 β -estradiol in MA-10 cells (Houk *et al.* 2004). Apart from their estrogenic effects, genistein (Sargeant *et al.* 1993) and quercetin (Lloyd *et al.* 1995) are potent tyrosine kinase inhibitors and have effects in the alteration of cytoplasm Ca^{2+} level. Resveratrol has been characterized as a nonspecific cyclooxygenase (COX) inhibitor (Murias *et al.* 2004). The aim of the present study was to compare the effects of quercetin, resveratrol, and genistein on *StAR* gene expression and steroid hormone synthesis in MA-10 cells, in which progesterone is the main steroid product due to the lack of P450_{c17} expression.

Materials and Methods

Materials and reagents

ICI 182,780 was purchased from Tocris (Ellisville, MO, USA). EGTA, mibefradil, verapamil, BK4866, sodium orthovanadate, and 8-bromo-adenosine-cAMP (8-Br-cAMP) were obtained from Sigma. DNase I, Murine Moloney Leukemia Virus (M-MLV) reverse transcriptase, Dual Luciferase Assay System, pRL-SV40 *Renilla* luciferase vector, and CellTiter 96 Aqueous Non-Radioactive Cell Proliferation Assay kits were purchased from Promega. Progesterone Enzyme Immunoassay kits were obtained from Cayman (Ann Arbor, MI, USA). Bicinchoninic Acid Protein Assay kits were purchased from Pierce Biochemicals (Rockford, IL, USA). The Genepot transfection reagent was obtained from Gene Therapy System, Inc. (San Diego, CA, USA). Trizol reagent, horse serum, and Waymouth's MB572/1 culture medium were obtained from Invitrogen Life Technologies, Inc.

Cell line and culture condition

The MA-10 mouse Leydig tumor cell line (Ascoli 1981) was a generous gift from Dr Mario Ascoli (University of Iowa, Iowa City, IA). MA-10 cells were grown in complete Waymouth's MB572/1 medium containing 2.2 g sodium bicarbonate, 100 U/ml penicillin/streptomycin, and 15% horse serum in a 37 °C and 5% CO₂ incubator.

Progesterone assay

MA-10 cells (5×10^4 /ml) were seeded into each well of 96-well plates. On the next day, the cells were incubated with 1, 10, 25, or 50 μM quercetin, resveratrol, or genistein. After 24 h, the cells were washed with Ca^{2+} -free phosphate buffer saline twice and replaced with assay medium (Waymouth MB572/1 containing 0.1% BSA (pH 7.4)) containing 0.2 mM 8-Br-cAMP for 4 h. The media were collected for the measurement of progesterone. Steroids were extracted by mixing 1 vol medium with 10 vol diethyl ether in 12 \times 75 mm glass tubes. The steroid extracts were evaporated in a 34 °C water bath in a ventilated chemical hood. The amount of progesterone in the extracts was measured according to the protocol accompanying the Progesterone Enzyme Immunoassay Kit. Ten microliters of radio immunoprecipitation assay (RIPA) lysis buffer (50 mM Tris-HCl (pH 7.5), 150 mM NaCl, 1 mM EGTA, 1% NP-40, 0.25% sodium deoxycholate, 2 $\mu\text{g}/\text{ml}$ aprotinin, 1 $\mu\text{g}/\text{ml}$ leupeptin, and 100 $\mu\text{g}/\text{ml}$ phenyl methyl sulfonyl fluoride (PMSF)) were added to each well containing cell monolayer and the protein concentration was determined using a Bicinchoninic Acid Protein Assay Kit.

StAR promoter-luciferase assay

MA-10 cells (1×10^5 /ml) were seeded into each well of six-well culture plates for 2 days until they were 50–60% confluent. The cells in each well were transfected with 2.0 μg mouse StAR promoter (P-966)–luciferase reporter gene (Caron *et al.* 1997a), 0.2 μg pRL-SV40 *Renilla* luciferase vector, and 15 μl GenePORTER transfection reagent in serum-free Waymouth's MB752/1 medium for 5 h. Cells were replaced with complete Waymouth's MB752/1 medium and incubated at 37 °C, 5% CO₂. On the next day, the cells were incubated with fresh Waymouth's MB752/1 medium containing 0.5, 1, 10, 25, or 50 μM quercetin, resveratrol, or genistein for another 24 h. The cells were then treated with or without 0.2 mM 8-Br-cAMP for 6 h and the lysate was harvested using Passive Lysis Buffer (Promega). The luciferase activity of the cell lysate was determined according to the manufacturer's protocol of Dual-Luciferase Reporter Assay System. The light intensity of each sample was measured in an EG&G Berthold LB 9507 Luminometer (Berthold, Bad Wildband, German). The StAR P-996 luciferase activity was normalized to that of the *Renilla* luciferase vector in order to compensate for the variation in transfection efficiency. Each treatment group consisted of triplicate cultures, and each experiment was repeated at least thrice.

Cell proliferation assay

MA-10 cells (2.5×10^4 /ml) were seeded into each well of 96-well plates. After 24 h, the cells were treated with 1, 10, 25, or 50 μ M quercetin, resveratrol, or genistein. Cell growth was measured from days 0 to 5 with the MTS (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium) assay according to the manufacturer's protocol. Absorbance was recorded at 490 nm using a Bio-Rad Benchmark microplate reader (Bio-Rad). Each treatment contained four replicates, and this experiment was performed at least thrice.

Real-time reverse transcription-PCR

MA-10 cells were treated with 50 μ M quercetin, resveratrol, or genistein and 0.2 mM 8-Br-cAMP for 24 h. Total RNA was prepared using Trizol reagent according to the manufacturer's protocol. RNA samples were incubated with RQ1 DNase at 37 °C for 30 min and then inactivated at 65 °C for 10 min. First strand cDNA was generated using 5 μ g DNase-treated RNAs and M-MLV reverse transcriptase at 42 °C for 1 h. The first strand cDNA was used for real-time PCR amplification reactions that were carried out using the MyIQ system (Bio-Rad). Reactions were initiated with hot start at 94 °C for 5 min, followed by 48 cycles at 94 °C for 30 s, 60 °C for 30 s, and 72 °C for 1 min with an extension of 5 min. The *StAR* gene was amplified using the primer pairs: forward: 5'-TTCA-AGCTGTGTGCTGG GAGCTCCTA-3' and reverse: 5'-TTAACACTGGGCC TCAGAGGCAGGGCTGG-3' which gave an amplification product of 247 bp. Primer S14 ribosomal protein was amplified using the forward: 5'-TTTGGTGTCTGCCACATCTTTG-3' and reverse: 5'-ATGGGGGTGACATCCTCAAT-3' sequences which yielded an amplification product of 323 bp (Mamchaoui *et al.* 2002). Reactions were performed in triplicate, and the average threshold cycle (C_T) was used in subsequent calculations to determine the *StAR* mRNA levels using the $\Delta\Delta C_T$ method of MyIQ system (Bio-Rad). The mean relative values \pm S.E.M. from three independent experiments were determined.

Statistical analysis

All data were analyzed by one-way ANOVA followed by Fisher's least significant difference test using the StatView 4.5 program (Abacus Concepts, Inc., Berkeley, CA, USA) fitted for the Macintosh computer. The results were expressed as the mean \pm S.E.M., and $P \leq 0.05$ was considered statistically significant.

Results

Effects of phytoestrogens on progesterone productions in MA-10 cells

In the present study, progesterone secretion of MA-10 cells in response to cAMP stimulation was decreased by genistein or

resveratrol at concentrations higher than 25 μ M (Fig. 1). On the other hand, quercetin increased progesterone synthesis (Fig. 1). These results suggested that there are differential effects of phytoestrogens on MA-10 cell steroidogenesis.

Effects of phytoestrogens on *StAR* mRNA levels and *StAR* promoter activity in transiently transfected MA-10 cells

We next evaluated the effects of genistein, resveratrol, and quercetin on *StAR* promoter activities in transiently transfected MA-10 cells. *StAR* promoter activity was inhibited by treatment with either genistein or resveratrol at concentrations higher than 10 or 25 μ M respectively (Fig. 2), when compared with cells treated with cAMP only. On the other hand, *StAR* promoter activity was enhanced by quercetin (Fig. 2). Cyclic AMP-stimulated *StAR* mRNA levels in MA-10 cells were decreased by treatment with 50 μ M genistein or resveratrol, but were increased by quercetin (Fig. 3). These results indicated that genistein and resveratrol have inhibitory effects, while quercetin has a stimulatory effect on MA-10 cell steroidogenesis.




Figure 1 Effects of phytoestrogens on progesterone synthesis. MA-10 cells were incubated with indicated amounts of genistein (G), resveratrol (R), or quercetin (Q) for 24 h. The cells were then replaced with assay medium containing 0.2 mM 8-Br-cAMP. After 4 h, the medium was collected and subjected to the extraction with diethyl ether, and the cell lysate was prepared by adding RIPA solution to the monolayer in each well. The progesterone levels in the ether extracts were measured using progesterone EIA kit as described under 'Materials and Methods'. The progesterone concentration in each well was normalized to the amount of total cellular protein. Each group contained triplicate samples. This experiment was repeated at least thrice. * Or † denotes significant increase or decrease when compared with the cells treated with cAMP only ($P < 0.05$ or 0.01 respectively). § Denotes significant increase when compared with the cells without cAMP treatment ($P < 0.001$). § Denotes significant difference when compared with the cells treated with cAMP plus genistein or resveratrol ($P < 0.01$).




Figure 2 Effects of phytoestrogens on StAR transcription. MA-10 cells were transiently transfected with a StAR P-966 luciferase reporter construct and pRL-Renilla luciferase plasmid. After 16 h, the transfected cells were incubated with indicated amounts of genistein (G), resveratrol (R), and quercetin (Q) for another 24 h. The cells were then treated with 0.2 mM 8-Br-cAMP for 6 h. Both StAR-reporter luciferase and Renilla luciferase activities were measured in the cell lysates as described under ‘Materials and Methods’. The activity of StAR P-966 luciferase was normalized to that of Renilla luciferase. For each experiment, the data were expressed as the -fold increase or decrease in treated cells relative to untreated cells that was set to a value of 1.0. The graph displays the mean values \pm s.e.m. for normalized data from three independent transfection experiments for StAR luc/Renilla luc. Each treatment contained triplicate samples. * Or † denotes a significant increase or decrease when compared with the cells treated with cAMP only ($P<0.05$ or $P<0.1$ respectively). ‡ Denotes a significant increase compared with the cells without cAMP treatment ($P<0.001$). § Denotes a significant difference when compared with the cells treated with cAMP plus genistein or resveratrol ($P<0.01$).

Effects of phytoestrogens on the proliferation of MA-10 cells

It has not been reported whether genistein, resveratrol, or quercetin affects MA-10 cell proliferation. In the present study, genistein inhibited MA-10 cell proliferation in a dose-dependent manner (Fig. 4A), while resveratrol or quercetin had no effect. The proliferation rate between days 1 and 5 in cells treated with genistein exhibited a gradual decline when compared with those treated with 0.1% dimethyl sulfoxide (DMSO). However, this phenomenon was not observed following treatment with either resveratrol or quercetin (Fig. 4B). During the first 48 h, incubation with 50 μ M genistein did not show any obvious inhibition on MA-10 cell growth and its cytotoxicity became discernable after 72 h in the cultures.

The role of estrogen receptor in quercetin-stimulated steroidogenesis

To investigate the role of ER in quercetin-stimulated steroid hormone synthesis, ICI 182,780, a potent ER antagonist




Figure 3 Effects of phytoestrogens on StAR mRNA expression. MA-10 cells were treated with 0.2 mM 8-Br-cAMP and 50 μ M genistein, resveratrol, or quercetin. After 24 h, cell monolayer was used for the preparation of total cellular RNA. The first strand of cDNA was synthesized from DNase-treated RNA samples by M-MLV reverse transcriptase. The first strand of cDNA was used as the template for real-time RT-PCR analysis as described under ‘Materials and Methods’. S14 rRNA was used as an internal standard to normalize StAR mRNA expression. The mRNA levels were expressed relative to control using $\Delta\Delta C_T$ method and represented as means \pm s.e.m. This experiment was repeated thrice using three independent sets of RNA samples. * Denotes a significant decrease when compared with the cells treated with cAMP ($P<0.01$). † Denotes a significant increase when compared with the cells treated with cAMP ($P<0.05$).

(Wakeling *et al.* 1991), was used to block endogenous ER activity in MA-10 cells. ICI 182,780 had no effect on cAMP-induced progesterone secretion, but reduced quercetin-plus cAMP-stimulated progesterone secretion by 31% (Fig. 5A). The effect of ICI 182,780 on quercetin-stimulated StAR reporter activity was next investigated. ICI 182,780 did not alter StAR promoter activity when cells were treated with cAMP alone or with cAMP plus quercetin (Fig. 5B). If activated ER has an inhibitory effect on MA-10 cell steroidogenesis, ER antagonists should increase MA-10 cell steroidogenesis. This result in the present study indicated that the effects of quercetin are not mediated by ER and are at a step beyond StAR.

The role of extracellular Ca^{2+} in quercetin-stimulated steroidogenesis

Steroid hormone synthesis in the gonads and adrenal glands is dependent on the trophic hormones and modulation by Ca^{2+} and various cytokines (Huhtaniemi & Toppari 1995, Rossier 1997). Whether extracellular Ca^{2+} plays a role in




Figure 4 Effects of phytoestrogens on MA-10 cell proliferation. MA-10 cells (5×10^3 cells/well) were seeded onto the 96-well plates and allowed to grow for 5 days in the presence of (A) indicated amounts of genistein, resveratrol, or quercetin. Cell proliferation was measured using MTS assay as described under 'Materials and Methods'. * Denotes a significant difference when compared with the cells treated with 50 μ M resveratrol or quercetin ($P < 0.05$). Data show the mean \pm S.E.M. of the values combined from three independent experiments. Four samples were used for each group. (B) The cells were incubated with 0.1% DMSO (control) or 50 μ M of each phytoestrogens. Cell growth was measured from days 0 to 5. * Denotes a significant decrease when compared with the control cells ($P < 0.05$). Data show the mean values \pm S.E.M. combined from three independent experiments. Four samples were used for each group.

quercetin-stimulated steroid hormone synthesis remains to be determined. In the presence of cAMP, EGTA decreased progesterone production and attenuated quercetin-mediated increase in progesterone secretion by 28% (Fig. 6A). EGTA also reduced cAMP- and cAMP- plus quercetin-stimulated StAR




Figure 5 Effects of ICI 182,780 on quercetin-stimulated steroidogenesis and StAR transcription. MA-10 cells were treated with 0.1% DMSO (control) or 50 μ M quercetin. After 24 h, cells were replaced with an assay medium containing 1 μ M ICI 182,780 and 0.2 mM 8-Br-cAMP for another 4 h. The medium in each well was collected and progesterone level was measured as described in Fig. 1. (A) ICI 182,780 decreased cAMP- and quercetin-stimulated progesterone secretion. * Denotes a significant increase when compared with the cells treated with cAMP ($P < 0.05$). † Denotes a significant decrease when compared with the cells treated with cAMP and quercetin ($P < 0.05$). (B) MA-10 cells were subjected to transient transfection as mentioned in Fig. 2, except treatment with 1 μ M ICI 182,780 and 0.2 mM 8-Br-cAMP for another 6 h. The relative luciferase activities were measured. Data show the mean values \pm S.E.M. combined from three independent sets of RNA samples. Each treatment contained four replicates. * Denotes a significant increase when compared with the cells treated with cAMP only ($P < 0.01$).

promoter activities (Fig. 6B). These results indicated that extracellular Ca^{2+} is required for quercetin-stimulated steroidogenesis in MA-10 cells. Voltage-gated Ca^{2+} channels and calcium-release-activated Ca^{2+} (CRAC) channels provide the major routes for Ca^{2+} to enter the cells in response to a




Figure 6 Effects of deprivation of extracellular calcium by EGTA on quercetin-stimulated progesterone secretion and StAR transcription. MA-10 cells were treated with 0.1% DMSO or 50 μ M quercetin for 24 h. The cells were replaced with an assay medium containing 4 mM EGTA and 0.2 mM 8-Br-cAMP for 4 h. The medium in each well was collected and progesterone levels were measured as described in Fig. 1. (A) EGTA reduced cAMP- and quercetin-stimulated progesterone secretion. (B) MA-10 cells were subjected to transient transfection as mentioned in Fig. 2, except treatment with 4 mM EGTA and 0.2 mM 8-Br-cAMP for 6 h. The relative luciferase activities were measured as described in Fig. 2. * Denotes a significant decrease when compared with the cells treated with cAMP only ($P < 0.05$). [†] Denotes a significant decrease when compared with the cells treated with cAMP and quercetin ($P < 0.01$). [‡] Denotes a significant increase when compared with the cells treated with cAMP only ($P < 0.01$). [§] Denotes a significant increase when compared with the cells treated with cAMP and EGTA ($P < 0.05$). Data show the mean values \pm S.E.M.1 combined from three independent experiments. Four samples were used for each group.

variety of stimulations (Rossier 1997). We sought to determine if blockade of Ca^{2+} channels affects quercetin-mediated responses in MA-10 cells. We utilized mibefradil (Massie 1998) and verapamil (Atlas & Adler 1981) to block T- and L-type voltage-gated Ca^{2+} channel. Treatment of MA-10 cells with 10 μ M of either mibefradil or verapamil had no effect on cAMP-stimulated progesterone synthesis, but significantly reduced quercetin-plus cAMP-stimulated progesterone secretion (Fig. 7A). However, neither mibefradil nor verapamil had an inhibitory effect on quercetin-plus cAMP-stimulated StAR promoter activity (Fig. 7B). Together, these results supported the view that extracellular Ca^{2+} influx is required for quercetin-plus cAMP-stimulated steroidogenesis in MA-10 cells and that Ca^{2+} acts at a step beyond StAR.

Effects of vanadate on quercetin-stimulated steroidogenesis

Calcium leakage through the plasma membrane (PM) Ca^{2+} pump or uptake by intracellular Ca^{2+} stores is essential for maintaining the homeostatic environment of the cytosol after elevation of cytosolic Ca^{2+} levels (Strehler & Treiman 2004). Vanadate and quercetin have been reported to inhibit the parathyroid hormone-stimulated PM Ca^{2+} pump activity in osteoblasts (Lloyd *et al.* 1995). We investigated the effect of sodium orthovanadate on MA-10 cell steroidogenesis. Treatment with 10 μ M vanadate inhibited progesterone secretion by 28% and quercetin-stimulated progesterone secretion by 36% (Fig. 8A). In the parallel experiments, vanadate had no effect on cAMP-stimulated StAR promoter, but significantly inhibited quercetin-plus cAMP-stimulated StAR promoter (Fig. 8B). These results indicate that Ca^{2+} pump activity is required for quercetin-stimulated progesterone synthesis.

Effects of ICI 182,780, EGTA, and vanadate on quercetin-increased StAR mRNA expression in MA-10 cells

We next examined the effect of ICI 182,780, EGTA, or vanadate on quercetin-induced StAR mRNA expression in the MA-10 cells. We observed that vanadate alone significantly reduced cAMP-stimulated StAR mRNA expression, while EGTA or ICI 182,780 had no effect. Vanadate and EGTA reduced quercetin-plus cAMP-increased StAR mRNA level by 50 or 36% respectively, while ICI 182,780 had no effect (Fig. 9).

Discussion

In the present study, we found that the phytoestrogens, quercetin, resveratrol, and genistein have differential effects on steroid hormone synthesis in MA-10 cells. Resveratrol and genistein inhibited progesterone secretion through down-regulation of StAR gene expression at the transcriptional and mRNA levels. MA-10 cell proliferation was decreased by treatment with genistein, but was unaffected by resveratrol or




Figure 7 Effects of blockade of T- or L-type voltage-gated Ca^{2+} channel on quercetin-stimulated steroidogenesis and StAR transcription. MA-10 cells were treated with either 0.1% DMSO or 50 μM quercetin for 24 h. Cells were incubated with an assay medium containing 10 μM mibefradil or verapamil and 0.2 mM 8-Br-cAMP for another 4 h. The medium in each well was collected and progesterone levels were measured as described in Fig. 1. (A) Mibefradil or verapamil reduced cAMP- and quercetin-stimulated progesterone synthesis. * Denotes a significant increase when compared with the cells treated with cAMP only ($P < 0.05$). † Denotes a significant decrease when compared with the cells treated with cAMP and quercetin ($P < 0.05$). (B) MA-10 cells were subjected to transient transfection as mentioned in Fig. 2, except treatment with 10 μM mibefradil or verapamil for 6 h. * Or † denotes a significant increase when compared with the cells treated with cAMP only ($P < 0.05$ or $P < 0.01$ respectively). Four samples were used for each treatment. Data show the mean values \pm S.E.M. combined from three independent experiments.




Figure 8 Effects of vanadate on quercetin-stimulated progesterone synthesis and StAR promoter activities in MA-10 cells. MA-10 cells were treated with either 0.1% DMSO or 50 μM quercetin for 24 h. Cells were incubated with an assay medium containing 10 μM sodium orthovanadate (Na_3VO_4) and 0.2 mM 8-Br-cAMP for another 4 h. The medium in each well was collected and progesterone levels were measured as described in Fig. 1. (A) Vanadate reduced cAMP- and quercetin-stimulated progesterone synthesis. * Denotes a significant decrease when compared with the cells treated with cAMP only ($P < 0.05$). † Denotes a significant increase when compared with the cells treated with cAMP and quercetin ($P < 0.05$). (B) MA-10 cells were subjected to transient transfection as mentioned in Fig. 2, except treatment with 10 μM Na_3VO_4 for 6 h. * Denotes a significant increase when compared with the cells treated with cAMP only ($P < 0.01$). † Denotes a significant decrease when compared with the cells treated with cAMP and quercetin ($P < 0.01$). Four samples were used for each treatment. Data show the mean values \pm S.E.M. combined from three independent experiments.




Figure 9 Effects of ICI 182,780, EGTA, and vanadate on quercetin-plus (cAMP+Q)-stimulated StAR mRNA expression. Cells were treated with 0.2 mM 8-Br-cAMP and 50 μ M quercetin for 24 h, followed by treatment with 1 mM ICI 182,780, 4 mM EGTA, or 10 μ M Na₃VO₄ for 4 h. The cell monolayer was used for the preparation of total cellular RNA. The first strand of cDNA was synthesized and subjected to real-time PCR analysis as mentioned in Fig. 3. The mRNA levels were expressed relative to cAMP treatment alone using $\Delta\Delta C_T$ method and represent the mean \pm S.E.M.. This experiment was repeated thrice using three independent sets of RNA samples. * Denotes a significant increase when compared with the cells treated with cAMP ($P < 0.01$). † Denotes a significant decrease when compared with the cells treated with cAMP and quercetin ($P < 0.05$). ‡ Denotes a significant decrease when compared with the cells treated with cAMP ($P < 0.01$).

quercetin. On the contrary, quercetin stimulated steroid hormone synthesis by up-regulation of StAR promoter activity and mRNA expression. ICI 182,780 decreased quercetin-stimulated progesterone secretion, but had no effect on quercetin-induced StAR promoter activity and StAR mRNA expression in MA-10 cells, suggesting that its effect is at a step beyond StAR. The role of cytosolic Ca²⁺ in the quercetin-stimulated steroidogenesis was next investigated. We found that Ca²⁺ is involved in quercetin-stimulated steroid hormone synthesis. We observed that the quercetin-induced increase in progesterone secretion is reduced by (1) deprivation of extracellular Ca²⁺ by EGTA treatment and (2) blocking extracellular Ca²⁺ influx with either mibefradil or verapamil in MA-10 cells. Quercetin-plus cAMP increased StAR promoter activity and StAR mRNA expression were reduced by EGTA, while mibefradil or verapamil had no effect. Blocking intracellular Ca²⁺ efflux by vanadate suppressed the quercetin-induced progesterone secretion, StAR promoter activity, and StAR mRNA expression, suggesting that protein phosphatases may be involved in quercetin-mediated response in MA-10 cells. These findings suggest that phytoestrogens have differential effects on steroid hormone synthesis in MA-10 cells.

Feeding rats with a phytoestrogen- or genistein-rich diet resulted in lower serum testosterone levels, but had no obvious effect on testicular StAR protein levels when compared with those fed with a regular diet (Weber *et al.* 2001, Svechnikov *et al.* 2005). Serum genistein levels were approximately 1.5 μ M in rats (Weber *et al.* 2001) and 10 μ M in men (Busby *et al.* 2002) after consumption with a phytoestrogen-rich diet. However, it was not clear whether such concentrations of genistein are sufficient to suppress StAR gene expression. In the present study, we treated MA-10 cells with 0.5–50 μ M genistein, resveratrol, and quercetin. We found that genistein and resveratrol in concentrations higher than 25 μ M decreased cAMP-stimulated progesterone secretion through down-regulation of StAR expression, while quercetin had an opposite effect. Our results are comparable with the previous report that genistein had an inhibitory effect on steroid hormone synthesis in adrenal glomerulosa (Bodart *et al.* 1995, Mesiano *et al.* 1999, Aptel *et al.* 1999, Ohno *et al.* 2002), Leydig (Svechnikov *et al.* 2005), and ovarian cells (Lacey *et al.* 2005). Moreover, genistein, resveratrol, and quercetin have been shown to suppress the proliferation of several tumor cells (Miodini *et al.* 1999, Mitchell *et al.* 1999, Setchell & Cassidy 1999, Xing *et al.* 2001). Since our assays were performed by the incubation with genistein for less than 30 h and 50 μ M genistein had no obvious inhibitory effect on MA-10 cell proliferation during the first 48 h in the cultures (Fig. 4B), therefore it is unlikely that genistein-mediated decreases on steroidogenesis and StAR gene expression were due to its cytotoxicity. On the other hand, we cannot rule out the possibility that genistein-mediated cytotoxicity has an impact on steroidogenesis in the animals receiving a long-term exposure to the higher doses of genistein. Feeding rats with resveratrol has been shown to decrease serum corticosterone levels (Supornsilchai *et al.* 2005) and our results showed that resveratrol decreased progesterone production in MA-10 cells. Resveratrol-mediated decrease in StAR gene expression is probably due to its estrogenic effect rather than its inhibitory effect on COX activities (Murias *et al.* 2004), because StAR expression in the MA-10 cells was increased by COX-2 inhibitors (Wang *et al.* 2003). Together, we conclude that genistein and resveratrol play a negative role in steroidogenesis. In contrast, we report for the first time that quercetin has a stimulatory effect on steroid production in MA-10 cells through up-regulation of StAR expression. In fact, quercetin alone was sufficient to induce a 2.5-fold increase in basal progesterone secretion in MA-10 cells (data not shown). Previously, quercetin was reported to have no suppressive effect on dibutyryl cAMP-stimulated cortisol secretion in H295R cells (Ohno *et al.* 2002), supporting our present finding.

Genistein, resveratrol, and quercetin are classified as phytoestrogens because of their capability to bind to both ER α and ER β and to activate the transcription of several estrogen-responsive genes *in vitro* (Kuiper *et al.* 1998, Bowers *et al.* 2000). The hierarchy of the binding affinity for both ER subtypes is genistein > quercetin \geq resveratrol when compared with 17 β -estradiol. The differences in the binding

affinity for ER α and ER β among these three phytoestrogens are unlikely to explain the discrepancy in quercetin-induced progesterone production in MA-10 cells. In the present study, ICI 182,780 did not affect cAMP-induced progesterone production but inhibited cAMP-plus quercetin-induced progesterone production. Our result is in agreement with the previous report that ICI 182,780 did not increase LH-stimulated testosterone production in isolated primary rat Leydig cells (Akingbemi *et al.* 2003). ICI 182,780 suppressed cAMP- plus quercetin-increased progesterone production without affecting StAR promoter activity and StAR mRNA expression, suggesting that the effect of ICI 182,780 is at a step beyond StAR.

Many factors such as trophic hormones, cytokines, Ca²⁺, kinases, and phosphatases have been reported to affect StAR gene expression (Manna & Stocco 2005). Previously, genistein was reported to decrease AngII-stimulated aldosterone synthesis by diminishing Ca²⁺ influx in adrenal glomerulosa cells (Aptel *et al.* 1999). In the present study, EGTA significantly decreased quercetin-plus cAMP-stimulated StAR transcriptional activity and StAR mRNA expression that attenuated progesterone production in MA-10 cells, suggesting that Ca²⁺ signal is involved in quercetin-stimulated steroidogenesis. Verapamil significantly reduced human chorionic gonadotropin (hCG)-plus potassium (K⁺)-stimulated progesterone synthesis and StAR mRNA levels in mLTC-1 mouse Leydig tumor cells (Manna *et al.* 1999). In the presence of quercetin plus cAMP stimulation, verapamil as well as mibefradil significantly reduced progesterone secretion, but fail to decrease StAR promoter activity and StAR mRNA expression in MA-10 cells. These results indicate that the major effect of Ca²⁺ influx through voltage-gated Ca²⁺ channels may contribute to quercetin-stimulated increases in steroidogenesis and its effect is at a step beyond StAR.

In the present report, vanadate suppressed cAMP- and cAMP-plus quercetin-stimulated progesterone secretion in MA-10 cells, implicating that cellular signals associated with vanadate may play a role in regulating quercetin-stimulated steroidogenesis. Vanadate did not alter StAR promoter activity, but reduced StAR mRNA expression when MA-10 cells were treated with cAMP, suggesting that there is a post-transcriptional regulation on StAR gene expression by vanadate. On the other hand, vanadate inhibited StAR promoter activity and StAR mRNA expression when the cells were treated with cAMP plus quercetin, suggesting that vanadate-sensitive cellular factors are responsible for quercetin effects on steroidogenesis. Vanadate is a well-known inhibitor for phosphoprotein phosphatase (PP) and alkaline phosphatase (ALP) (Gordon 1991) and is able to enhance the sensitivity of mouse Leydig cell proliferation in response to 17 β -estradiol stimulation (Sato *et al.* 1987). Recently, PP was reported to play a role in regulating cAMP-mediated increase in StAR protein expression and steroidogenesis in mouse Y1 adrenocortical cells (Jones *et al.* 2000), while PPs are also involved in the elevation of mitochondria Ca²⁺ level that

may be associated with K⁺-stimulated aldosterone synthesis in H295R cells (Lalevee *et al.* 2003).

In conclusion, phytoestrogens have differential effects on MA-10 cell steroidogenesis. Genistein and resveratrol have a negative role, while quercetin has a stimulatory effect on steroidogenesis. The effects of quercetin are complex and may depend on the non-ER-mediated events and Ca²⁺ in MA-10 cells.

Acknowledgements

This work was supported by the Veterans Affairs Medical Research Fund (to T L), a South Carolina Cancer Research Award (to Y C C), and by NIH Grant HD17481 and funds from the Robert A Welch Foundation Grant B1-0028 (to D M S). The authors declare that there is no conflict of interest that would prejudice the impartiality of this scientific work.

References

- Akingbemi BT, Ge R, Rosenfeld CS, Newton LG, Hardy DO, Catterall JF, Lubahn DB, Korach KS & Hardy MP 2003 Estrogen receptor- α gene deficiency enhances androgen biosynthesis in the mouse Leydig cell. *Endocrinology* **144** 84–93.
- Aptel HB, Burnay MM, Rossier MF & Capponi AM 1999 The role of tyrosine kinases in capacitative calcium influx-mediated aldosterone production in bovine adrenal zona glomerulosa cells. *Journal of Endocrinology* **163** 131–138.
- Ascoli M 1981 Characterization of several clonal lines of cultured Leydig tumor cells: gonadotropin receptors and steroidogenic responses. *Endocrinology* **108** 88–95.
- Atlas D & Adler M 1981 Alpha-adrenergic antagonists as possible calcium channel inhibitors. *PNAS* **78** 1237–1241.
- Bodart V, Ong H & De Lean A 1995 A role for protein tyrosine kinase in the steroidogenic pathway of angiotensin II in bovine zona glomerulosa cells. *Journal of Steroid Biochemistry and Molecular Biology* **54** 55–62.
- Bose HS, Sugawara T, Strauss JF III & Miller WL 1996 The pathophysiology and genetics of congenital lipid adrenal hyperplasia. International Congenital Lipoid Adrenal Hyperplasia Consortium. *New England Journal of Medicine* **335** 1870–1878.
- Bowers JL, Tyulmenkov VV, Jernigan SC & Klinge CM 2000 Resveratrol acts as a mixed agonist/antagonist for estrogen receptors α and β . *Endocrinology* **141** 3657–3667.
- Busby MG, Jeffcoat AR, Bloedon LT, Koch MA, Black T, Dix KJ, Heizer WD, Thomas BF, Hill JM, Crowell JA *et al.* 2002 Clinical characteristics and pharmacokinetics of purified soy isoflavones: single-dose administration to healthy men. *American Journal of Clinical Nutrition* **75** 126–136.
- Caron KM, Ikeda Y, Soo SC, Stocco DM, Parker KL & Clark BJ 1997a Characterization of the promoter region of the mouse gene encoding the steroidogenic acute regulatory protein. *Molecular Endocrinology* **11** 138–147.
- Caron KM, Soo SC, Wetsel WC, Stocco DM, Clark BJ & Parker KL 1997b Targeted disruption of the mouse gene encoding steroidogenic acute regulatory protein provides insights into congenital lipid adrenal hyperplasia. *PNAS* **94** 11540–11545.
- Clark BJ, Wells J, King SR & Stocco DM 1994 The purification, cloning, and expression of a novel luteinizing hormone-induced mitochondrial protein in MA-10 mouse Leydig tumor cells. Characterization of the steroidogenic acute regulatory protein (StAR). *Journal of Biological Chemistry* **269** 28314–28322.
- Clark BJ, Combs R, Hales KH, Hales DB & Stocco DM 1997 Inhibition of transcription affects synthesis of steroidogenic acute regulatory protein and steroidogenesis in MA-10 mouse Leydig tumor cells. *Endocrinology* **138** 4893–4901.

- DuMond JW Jr, Singh KP & Roy D 2001 Regulation of the growth of mouse Leydig cells by the inactive stereoisomer, 17 α -estradiol: lack of correlation between the elevated expression of ER α and difference in sensitivity to estradiol isomers. *Oncology Reports* **8** 899–902.
- Eddy EM, Washburn TF, Bunch DO, Goulding EH, Gladen BC, Lubahn DB & Korach KS 1996 Targeted disruption of the estrogen receptor gene in male mice causes alteration of spermatogenesis and infertility. *Endocrinology* **137** 4796–4805.
- Freeman DA 1985 Estradiol acts as a competitive inhibitor of the 3 β -hydroxysteroid dehydrogenase/delta 5-delta 4 isomerase enzyme of cultured Leydig tumor cells. *Endocrinology* **117** 2127–2133.
- Gordon JA 1991 Use of vanadate as protein-phosphotyrosine phosphatase inhibitor. *Methods in Enzymology* **201** 477–482.
- Henry LA & Witt DM 2002 Resveratrol: phytoestrogen effects on reproductive physiology and behavior in female rats. *Hormones and Behavior* **41** 220–228.
- Houk CP, Pearson EJ, Martinelle N, Donahoe PK & Teixeira J 2004 Feedback inhibition of steroidogenic acute regulatory protein expression *in vitro* and *in vivo* by androgens. *Endocrinology* **145** 1269–1275.
- Huhtaniemi I & Toppari J 1995 Endocrine, paracrine and autocrine regulation of testicular steroidogenesis. *Advances in Experimental Medicine and Biology* **377** 33–54.
- Jones PM, Sayed SB, Persaud SJ, Burns CJ, Gyles S & Whitehouse BJ 2000 Cyclic AMP-induced expression of steroidogenic acute regulatory protein is dependent upon phosphoprotein phosphatase activities. *Journal of Molecular Endocrinology* **24** 233–239.
- Kuiper GG, Lemmen JG, Carlsson B, Corton JC, Safe SH, van der Saag PT, van der Burg B & Gustafsson JA 1998 Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor beta. *Endocrinology* **139** 4252–4263.
- Lacey M, Bohday J, Fonseka SM, Ullah AI & Whitehead SA 2005 Dose-response effects of phytoestrogens on the activity and expression of 3 β -hydroxysteroid dehydrogenase and aromatase in human granulosa-luteal cells. *Journal of Steroid Biochemistry and Molecular Biology* **96** 279–286.
- Lalevee N, Resin V, Arnaudeau S, Demaux N & Rossier MF 2003 Intracellular transport of calcium from plasma membrane to mitochondria in adrenal H295R cells: implication for steroidogenesis. *Endocrinology* **144** 4575–4585.
- Le Bail JC, Champavier Y, Chulia AJ & Habrioux G 2000 Effects of phytoestrogens on aromatase, 3 β and 17 β -hydroxysteroid dehydrogenase activities and human breast cancer cells. *Life Sciences* **66** 1281–1291.
- Lin D, Sugawara T, Strauss JF III, Clark BJ, Stocco DM, Saenger P, Rogol A & Miller WL 1995 Role of steroidogenic acute regulatory protein in adrenal and gonadal steroidogenesis. *Science* **267** 1828–1831.
- Lloyd QP, Kuhn MA & Gay CV 1995 Characterization of calcium translocation across the plasma membrane of primary osteoblasts using a lipophilic calcium-sensitive fluorescent dye, calcium green C18. *Journal of Biological Chemistry* **270** 22445–22451.
- Mamchaoui K, Makhoulfi Y & Saumon G 2002 Glucose transporter gene expression in freshly isolated and cultured rat pneumocytes. *Acta Physiologica Scandinavica* **175** 19–24.
- Manna PR & Stocco DM 2005 Regulation of the steroidogenic acute regulatory protein expression: functional and physiological consequences. *Current Drug Targets. Immune, Endocrine and Metabolic Disorders* **5** 93–108.
- Manna PR, Pakarinen P, El-Hefnawy T & Huhtaniemi IT 1999 Functional assessment of the calcium messenger system in cultured mouse Leydig tumor cells: regulation of human chorionic gonadotropin-induced expression of the steroidogenic acute regulatory protein. *Endocrinology* **140** 1739–1751.
- Massie BM 1998 Mibefradil, a T-type channel-selective calcium antagonist: clinical trials in chronic stable angina pectoris. *American Journal of Hypertension* **11** 95S–102S.
- Mesiano S, Katz SL, Lee JY & Jaffe RB 1999 Phytoestrogens alter adrenocortical function: genistein and daidzein suppress glucocorticoid and stimulate androgen production by cultured adrenal cortical cells. *Journal of Clinical Endocrinology and Metabolism* **84** 2443–2448.
- Miodini P, Fioravanti L, Di Fronzo G & Cappelletti V 1999 The two phytoestrogens genistein and quercetin exert different effects on oestrogen receptor function. *British Journal of Cancer* **80** 1150–1155.
- Mitchell SH, Zhu W & Young CY 1999 Resveratrol inhibits the expression and function of the androgen receptor in LNCaP prostate cancer cells. *Cancer Research* **59** 5892–5895.
- Mizutani K, Ikeda K, Kawai Y & Yamori Y 2000 Resveratrol attenuates ovariectomy-induced hypertension and bone loss in stroke-prone spontaneously hypertensive rats. *Journal of Nutritional Science and Vitaminology* **46** 78–83.
- Murias M, Handler N, Erker T, Pleban K, Ecker G, Saiko P, Szekeres T & Jager W 2004 Resveratrol analogues as selective cyclooxygenase-2 inhibitors: synthesis and structure-activity relationship. *Bioorganic and Medicinal Chemistry* **12** 5571–5578.
- Nakajima D, Kim CS, Oh TW, Yang CY, Naka T, Igawa S & Ohta F 2001 Suppressive effects of genistein dosage and resistance exercise on bone loss in ovariectomized rats. *Journal of Physiological Anthropology and Applied Human Science* **20** 285–291.
- O'Donnell L, Robertson KM, Jones ME & Simpson ER 2001 Estrogen and spermatogenesis. *Endocrine Reviews* **22** 289–318.
- Ohno S, Shinoda S, Toyoshima S, Nakazawa H, Makino T & Nakajin S 2002 Effects of flavonoid phytochemicals on cortisol production and on activities of steroidogenic enzymes in human adrenocortical H295R cells. *Journal of Steroid Biochemistry and Molecular Biology* **80** 355–363.
- Ohno S, Nakajima Y, Inoue K, Nakazawa H & Nakajin S 2003 Genistein administration decreases serum corticosterone and testosterone levels in rats. *Life Sciences* **74** 733–742.
- Pelletier G, Labrie C & Labrie F 2000 Localization of oestrogen receptor alpha, oestrogen receptor beta and androgen receptors in the rat reproductive organs. *Journal of Endocrinology* **165** 359–370.
- Rossier MF 1997 Confinement of intracellular calcium signaling in secretory and steroidogenic cells. *European Journal of Endocrinology* **137** 317–325.
- Sargeant P, Farndale RW & Sage SO 1993 ADP- and thapsigargin-evoked Ca²⁺ entry and protein-tyrosine phosphorylation are inhibited by the tyrosine kinase inhibitors genistein and methyl-2,5-dihydroxycinnamate in fura-2-loaded human platelets. *Journal of Biological Chemistry* **268** 18151–18156.
- Sato B, Miyashita Y, Maeda Y, Norma K, Kishimoto S & Matsumoto K 1987 Effects of estrogen and vanadate on the proliferation of newly established transformed mouse Leydig cell line *in vitro*. *Endocrinology* **120** 1112–1120.
- Saunders PT, Fisher JS, Sharpe RM & Millar MR 1998 Expression of oestrogen receptor beta (ER beta) occurs in multiple cell types, including some germ cells, in the rat testis. *Journal of Endocrinology* **156** R13–R17.
- Setchell KD & Cassidy A 1999 Dietary isoflavones: biological effects and relevance to human health. *Journal of Nutrition* **129** 758S–767S.
- Strehler EE & Treiman M 2004 Calcium pumps of plasma membrane and cell interior. *Current Molecular Medicine* **4** 323–335.
- Sugawara T, Holt JA, Driscoll D, Strauss JF III, Lin D, Miller WL, Patterson D, Clancy KP, Hart IM, Clark BJ *et al.* 1995 Human steroidogenic acute regulatory protein: functional activity in COS-1 cells, tissue-specific expression, and mapping of the structural gene to 8p11.2 and a pseudogene to chromosome 13. *PNAS* **92** 4778–4782.
- Supornsilchai V, Svechnikov K, Seidlova-Wuttke D, Wuttke W & Soder O 2005 Phytoestrogen resveratrol suppresses steroidogenesis by rat adrenocortical cells by inhibiting cytochrome P450 c21-hydroxylase. *Hormone Research* **64** 280–286.
- Svechnikov K, Supornsilchai V, Strand ML, Wahlgren A, Seidlova-Wuttke D, Wuttke W & Soder O 2005 Influence of long-term dietary administration of procymidone, a fungicide with anti-androgenic effects, or the phytoestrogen genistein to rats on the pituitary–gonadal axis and Leydig cell steroidogenesis. *Journal of Endocrinology* **187** 117–124.
- Wakeling AE, Dukes M & Bowler JA 1991 Potent specific pure antiestrogen with clinical potential. *Cancer Research* **51** 3867–3873.

- Wang X, Dyson MT, Jo Y & Stocco DM 2003 Inhibition of cyclooxygenase-2 activity enhances steroidogenesis and steroidogenic acute regulatory gene expression in MA-10 mouse Leydig cells. *Endocrinology* **144** 3368–3375.
- Wattel A, Kamel S, Mentaverri R, Lorget F, Prouillet C, Petit JP, Fardelonne P & Brazier M 2003 Potent inhibitory effect of naturally occurring flavonoids quercetin and kaempferol on *in vitro* osteoclastic bone resorption. *Biochemical Pharmacology* **65** 35–42.
- Weber KS, Setchell KD, Stocco DM & Lephart ED 2001 Dietary soy-phytoestrogens decrease testosterone levels and prostate weight without altering LH, prostate 5 α -reductase or testicular steroidogenic acute regulatory peptide levels in adult male Sprague–Dawley rats. *Journal of Endocrinology* **170** 591–599.
- Wisniewski AB, Klein SL, Lakshmanan Y & Gearhart JP 2003 Exposure to genistein during gestation and lactation demasculinizes the reproductive system in rats. *Journal of Urology* **169** 1582–1586.
- Xing N, Chen Y, Mitchell SH & Young CY 2001 Quercetin inhibits the expression and function of the androgen receptor in LNCaP prostate cancer cells. *Carcinogenesis* **22** 409–414.

Received in final form 26 November 2006

Accepted 19 December 2006

Made available online as an Accepted Preprint

3 January 2007