

# Semen quality in the 21<sup>st</sup> century

Helena E. Virtanen<sup>1</sup>, Niels Jørgensen<sup>2</sup> and Jorma Toppari<sup>1–3</sup>

**Abstract** | Although semen quality is an important determinant of fertility, defining clear thresholds for normal ranges has proven difficult. According to ‘time to pregnancy’ studies, fecundity starts to decline when sperm concentrations fall below  $30\text{--}55 \times 10^6/\text{ml}$ , whereas the WHO criterion for normal values is currently  $15 \times 10^6/\text{ml}$ . Multiple studies over the past 15 years have reported median sperm concentrations of  $41\text{--}55 \times 10^6/\text{ml}$  in young men (mean age 18–21 years) from the general population, suggesting that many of them have suboptimal semen quality. Sperm numbers remain fairly constant between 19 and 29 years of age, which points to the importance of developmental effects. Discussion on whether population semen quality has declined has continued for decades, as regional differences in trends have been noted. The reasons for poor semen quality and adverse trends are not well established, but some associations suggest a causal relationship, for example, with maternal smoking during pregnancy. The role of chemical exposures leading to endocrine disruption and detrimental reproductive effects has been in the focus of research during the past 20 years. Identification of exposures that affect fertility could provide opportunities for effective prevention of reproductive health problems.

Several original studies published since 2000 in various countries have evaluated possible time trends in sperm concentration or total sperm count in different populations (TABLE 1). These studies have suggested a decreasing trend in sperm concentration and/or total sperm count of young men<sup>1,2</sup>, male partners in infertile couples<sup>3–8</sup>, fertile men<sup>9,10</sup> and semen donors<sup>11–13</sup> or semen donor candidates<sup>14–17</sup>. However, studies reporting no significant decrease or, indeed, that found a slightly increasing trend, have also been published<sup>18–24</sup>.

Change in sperm concentration over time (1938–2013) was evaluated in a systematic review and meta-analysis investigating the effect of age on semen quality<sup>25</sup>. Altogether, 124 measurements of mean sperm concentration were included in the analysis, and a significant decline in sperm concentration over years 1938–2013 and also over years 1994–2013 was observed ( $P < 0.001$  and  $P < 0.02$ , respectively). However, the latter decline was no longer significant when possible confounding factors — such as mean age, sample source, abstinence control, and gross domestic product for the country of the study — were taken into account<sup>25</sup>. Furthermore, an analysis studying 1994–2013 only included studies that were identified when looking for data on the associations between male ageing and semen quality. Thus, a meta-analysis focusing specifically on evaluating time trends in semen quality in the 21<sup>st</sup> century is needed in the future. In addition to studies of time trends in semen quality, original studies have also suggested regional differences in semen quality in the 21<sup>st</sup> century.

## Geographical differences in semen quality

At the end of the 1990s, two coordinated cross-sectional studies using standardized investigation procedures were performed in Europe to study possible regional differences in the semen quality of young men and of partners of pregnant women<sup>26,27</sup>. Study protocols similar to those used in these two reports<sup>26,27</sup> have been used in several further studies published since 2000 (TABLE 2, FIGS 1, 2).

## Young men

One of the above mentioned coordinated studies evaluated semen quality among young men in Denmark, Finland, Estonia and Norway, and showed that sperm concentrations and total sperm counts were significantly higher among young Finnish or Estonian men than among young Danish or Norwegian men (Finland versus Denmark 95% CI for difference 1.08–1.59, Finland versus Norway 95% CI for difference 1.06–1.60, Estonia versus Denmark 95% CI for difference 1.07–1.79, Estonia versus Norway 95% CI for difference 1.05–1.82) when adjusted to the Danish laboratory level and abstinence time. No significant differences between Finnish and Estonian men or between Danish and Norwegian men were observed<sup>27</sup>. In 2002, young Swedish men aged 18–21 years were reported to have significantly higher sperm concentrations than young Danish men (mean difference  $13.4 \times 10^6/\text{ml}$ , 95% CI 4.1–22.7). However, no reference laboratory was used in the study and potential interlaboratory differences in evaluation of sperm concentration could not, therefore,

<sup>1</sup>Department of Physiology, Institute of Biomedicine, University of Turku, Kiinamyllynkatu 10, 20520 Turku, Finland.

<sup>2</sup>University Department of Growth and Reproduction and International Center for Research and Research Training in Endocrine Disruption of Male Reproduction and Child Health (EDMaRC), Rigshospitalet, Blegdamsvej 9, DK-2100 Copenhagen, Denmark.

<sup>3</sup>Department of Paediatrics, Turku University Hospital, Kiinamyllynkatu 4 – 8, PL 52, 20521 Turku, Finland.

Correspondence to J.T. [jorma.toppari@utu.fi](mailto:jorma.toppari@utu.fi)

doi:10.1038/nruiol.2016.261  
Published online 4 Jan 2017

### Key points

- Downward trends in sperm concentrations have been described in several geographical areas during this century
- In several countries, sperm concentration of a considerable proportion of young men has been described to be on a level that has been associated with prolonged time to pregnancy
- Longitudinal studies suggest that almost full sperm production capacity is achieved around the age of 20 years, which points to the importance of earlier developmental phases in establishment of spermatogenic capacity
- Environmental factors are likely to have contributed to the declining trends in sperm concentrations; however, identifying the most important factors causing adverse effects remains a challenge

be taken into account when calculating the difference in sperm concentrations between countries<sup>28,29</sup>. Punab *et al.*<sup>30</sup> studied semen quality among military conscripts in Estonia and Lithuania, and sperm concentrations seemed to be higher in Estonian men. Sperm concentrations among Latvian military conscripts have also been reported to be somewhat higher than in Swedish young men, but significantly higher than in Danish men (mean difference  $17 \times 10^6/\text{ml}$ , 95% CI for difference:  $3.2\text{--}31 \times 10^6/\text{ml}$ )<sup>28,29,31</sup>. However, these data came from a small study and the results were not adjusted to the Danish reference laboratory level<sup>31</sup>. Around the turn of the century, adjusted sperm concentrations in young men from Japan, Lithuania, and Southern Spain seemed to be higher than in young men from Norway and Denmark<sup>27,30,32,33</sup>, and German young men seemed to have similar sperm concentrations as Danish and Norwegian young men<sup>27,34</sup>. No difference in sperm concentration was observed between young men from Eastern and Western Germany<sup>34</sup>. When compared with Danish men examined in 2006–2010, Faroese young men had lower sperm concentrations ( $P < 0.001$ ) when adjusted for duration of abstinence<sup>35</sup>; however, the Faroese men were significantly older than the Danish men (median age 24 years versus 19 years, respectively,  $P < 0.001$ )<sup>35</sup>. Around 2010, sperm concentrations of young men from USA and a Western Australian birth cohort were also studied, but adjusted results were not provided and no statistical comparison to the sperm concentration in the European countries has been reported<sup>36,37</sup>.

The previously reported difference in sperm parameters between Finland and Denmark seems to be narrowing, as sperm concentrations seem to be decreasing in Finland and increasing in Denmark<sup>2,20</sup>. Similar trends have also been observed when excluding men with previous or current andrological disorders<sup>2,20</sup>. The increase in sperm numbers among the Danish men was statistically significant ( $P = 0.02$  for sperm concentration in 1996–2000 versus 2006–2010), but, from a biological point of view, rather small. Thus, future studies will have to address whether this possible trend is continuing or whether it is due to random fluctuation that is often seen in population-based studies. The difference between sperm counts in Sweden and Denmark seems to have diminished less, as no decrease in sperm concentrations

has been reported from Sweden<sup>19,20,28</sup>. Time trend studies also suggest that sperm concentrations in young Spanish men might have declined in the twenty-first century<sup>1</sup>. Thus, possible differences between birth cohorts in each country should also be taken into account, when studying geographical differences in sperm concentrations. In a longitudinal study of Finnish men who provided semen samples four times over a 10-year period, sperm concentration remained fairly similar between the ages of 19 and 29 years<sup>38</sup>. These data are in line with a previous longitudinal study, which observed no significant change in sperm concentration during 4-year follow-up period of Danish young men, with median age at entry of 19 years<sup>39</sup>. Thus, cross-sectional data on sperm concentrations of 19-year-old men do seem to represent their adult spermatogenic capacity. Semen quality varies regionally and temporally, suggesting both genetic and environmental effects behind the trends.

### Fertile men

When evaluating semen quality of male partners of pregnant women in four European cities in a cross-sectional coordinated study, Jørgensen *et al.*<sup>26</sup> observed that Danish men had the lowest sperm concentrations, followed by French and Scottish men, and Finnish men had the highest sperm concentrations when adjusted for possible confounders (age, abstinence time, and season). No significant differences between the four laboratories were observed in the assessment of sperm concentration according to a quality control study<sup>26</sup>. In a study designed in collaboration with that of Jørgensen and colleagues, significantly lower sperm concentrations were observed among male partners of pregnant women from Columbia than among partners of pregnant women from New York, Minneapolis, or Los Angeles ( $P = 0.001$ ,  $P < 0.001$ , and  $P = 0.060$ , respectively, when using haemocytometer for determination of sperm concentration and  $P = 0.002$ ,  $P < 0.001$ , and  $P = 0.005$ , respectively when using  $\mu$ -cell disposable counting chamber)<sup>40</sup>. The final data analysis of the US study included increased number of men, and also included men from Iowa; the finding that the lowest sperm concentrations in the USA are observed in men from Missouri was confirmed<sup>41</sup>. A further study on urine pesticide metabolite levels showed a significant association between high pesticide levels and risk of poor semen quality among partners of pregnant women from Missouri but not among similar men from Minnesota<sup>42</sup>. No statistical comparisons between the sperm concentrations in USA and Europe have been reported.

Iwamoto *et al.*<sup>43</sup> evaluated semen quality in Japanese fertile men using a protocol harmonized with the European study by Jørgensen<sup>26</sup>. Men from the Kawasaki and Yokohama areas of Japan had significantly lower sperm concentrations than men from Scotland (Edinburgh) and Finland (Turku) ( $P = 0.0008$  and  $P < 0.0001$ , respectively), but similar to levels of men from Denmark (Copenhagen) and France (Paris)<sup>26,43</sup>. In a subsequent study on semen quality in fertile Japanese men, statistically significant differences in sperm concentrations were observed between four urban areas (Sapporo,

Osaka, Kanazawa, and Fukuoka) in Japan ( $P=0.04$ )<sup>44</sup>. Adjusted sperm concentrations seemed higher than in the previous Japanese study and similar to the adjusted sperm concentrations in previous study on fertile European men.

Semen quality in fertile Australian men was determined as a part of the WHO surveillance study<sup>45</sup> and Haugen *et al.*<sup>46</sup> determined semen variables in fertile Norwegian men. However, no adjusted sperm concentrations have been reported and no statistical comparison with the sperm concentration in other countries has been performed. Reasons for the regional differences in semen quality are not known, but lifestyle and environmental factors are likely to have role, in addition to genetic factors.

### Lifestyle and environmental factors

Low spermatogenic capacity can be associated with developmental disorders of the male reproductive system, such as cryptorchidism, hypospadias, and testicular germ cell cancer, which are all components of testicular dysgenesis syndrome (TDS)<sup>47</sup>. Increasing rates of TDS components have been reported during past decades in western countries, and it has been suggested that

development of TDS is most often associated with exposure of the fetal testis to environmental factors such as endocrine-disrupting chemicals<sup>48</sup>. Exposure to a multitude of endocrine disruptors has emerged during the past century. Thus, besides being possibly affected by concurrent lifestyle factors, semen quality also has developmental determinants.

### Smoking

Associations between several lifestyle and environmental factors have been studied. Meta-analyses have suggested that smoking is associated negatively with semen quality<sup>49–51</sup>, and is also thought to be associated with increased frequency of sperm aneuploidy and DNA fragmentation<sup>52–54</sup>. Furthermore, prenatal exposure to maternal smoking is also associated with reduced semen quality of adult male offspring — negative associations with sperm concentration, total sperm count, percentage of motile sperm or percentage of morphologically normal sperm, or increased risk of oligozoospermia have been observed<sup>34,55–60</sup> — and men prenatally exposed to smoking are more likely to be smokers themselves<sup>58</sup>. Regular smoking of marijuana has been negatively associated with sperm concentration<sup>61</sup>. Tobacco smoke

Table 1 | 21<sup>st</sup> century trends in sperm parameters

Study	Area	Population	Mean age (years)	n	Study period	Overall change in sperm concentration	Overall change in total sperm count
Feki <i>et al.</i> <sup>5</sup> (2009)	Tunisia	Male partners of infertile couples	36	1,835*	1996–2007	NS	↓
Jiang <i>et al.</i> <sup>10</sup> (2014)	China, Sichuan province	Men examined for fertility in sperm bank	Median 32	28,213	2007–2012	↓ Fertile men ↓ Healthy men ↓ Infertile men	NA
Rao <i>et al.</i> <sup>12</sup> (2015)	China, Hubei Province, Wuhan	Semen donors (university students)	Range 22–30	1,808	2010–2013	↓	↓
Huang <i>et al.</i> <sup>17</sup> (2016)	China, Hunan Province	Semen donor candidates	21.6	3,114	2001–2005	↓	↓
			21.4	10,386	2006–2010		
			21.9	17,136	2011–2015		
Wang <i>et al.</i> <sup>13</sup> (2016)	China, Shandong Province	Semen donors	25.8	5,210	2008–2014	↓	↓
Marimuthu <i>et al.</i> <sup>23</sup> (2003)	India	Men attending infertility clinic	31.2	1,176	1990–2000	NS†	NA
Adiga <i>et al.</i> <sup>8</sup> (2008)	India	Men evaluated for infertility	NA	7,770	1993–2005	↓	NA
Mukhopadhyay <i>et al.</i> <sup>22</sup> (2010)	India	Male partners of infertile couples*	33.2	1752	1981–1985	NS	NA
			35.2	1,977	2001–2006		
Haimov-Kochman <i>et al.</i> <sup>11</sup> (2012)	Israel	Healthy semen donors	25.2	58 (2,182 samples)	1995–2009	↓	NA
Lackner <i>et al.</i> <sup>3</sup> (2005)	Austria	Men evaluated for infertility (azoospermic men excluded)	Median 31.6	7,780	1986–2003	↓	NA
Jørgensen <i>et al.</i> <sup>20</sup> (2012)	Denmark	Young men from the general population	19.4	4,867	1996–2010	↑	↑
Jørgensen <i>et al.</i> <sup>2</sup> (2011)	Finland	Young men from the general population	19	858	1998–2006	↓	↓

Table 1 (cont.) | 21<sup>st</sup> century trends in sperm parameters

Study	Area	Population	Mean age (years)	n	Study period	Overall change in sperm concentration	Overall change in total sperm count
Geoffroy-Siraudin <i>et al.</i> <sup>6</sup> (2012)	France (Marseille)	Male partners of infertile couples	35.1	10,932 7,899 <sup>§</sup>	1988–2007	↓	↓
Rolland <i>et al.</i> <sup>7</sup> (2013)	France (126 ART centres)	Male partners of infertile women undergoing first IVF or ICSI	35.2	26,609	1989–2005	↓	NA
Splingart <i>et al.</i> <sup>15</sup> (2012)	France (Tours)	Semen donor candidates	35.2	1,114	1976–2009	NA	↓
Sripada <i>et al.</i> <sup>4</sup> (2007)	UK	Male partners of subfertile couples*	Range 22–56	4,832	1994–2005	↓	NA
Romero-Otero <i>et al.</i> <sup>9</sup> (2015)	Spain	Fertile men undergoing vasectomy	37.8	992	1985–2009	↓	↓
Fernandez <i>et al.</i> <sup>32</sup> (2012), Mendiola <i>et al.</i> <sup>1</sup> (2013)	Spain	Young men in Almeria	21.3	273	2001–2002	NA	NA
		Healthy university students in Murcia	19.2	215	2010–2011	↓ (compared with Fernandez <i>et al.</i> <sup>32</sup> )	↓ (compared with Fernandez <i>et al.</i> <sup>32</sup> )
Axelsson <i>et al.</i> <sup>19</sup> (2011), Richthoff <i>et al.</i> <sup>28</sup> (2002)	Sweden	Young native men from the general population	18.2	216	2000–2001	NS	NS
			18.0	295	2008–2010		
Chen <i>et al.</i> <sup>18</sup> (2003)	USA	Male partners of infertile couples	36.3; n = 551	408	1989–2000	↑ (NS)	NA
Centola <i>et al.</i> <sup>16</sup> (2016)	USA, Boston area	Semen donor candidates	21.9–25.1	489 (9,425 samples)	2003–2013	↓	↓
Costello <i>et al.</i> <sup>21</sup> (2002)	Australia	Semen donor candidates	Range 18–40	448	1983–2001	NA	NS
Shine <i>et al.</i> <sup>14</sup> (2008)	New Zealand	Semen donor candidates	Median 36	975	1987–2007	↓	NA
Birdsall <i>et al.</i> <sup>24</sup> (2015)	New Zealand	Semen donors	Median 35	285	2008–2014	NS (when compared with period 2001–2007)	NA

NA not available, NS not significant. \*Men with sperm concentration >20 M/mL. †Men with sperm concentration <10 M/mL excluded. §Men with total sperm count >40M.

contains several hazardous substances and increased oxidative stress, DNA damage, cell apoptosis, and a direct effect on regulation of spermatogenesis have been suggested as possible mechanisms for the adverse effects of smoking on semen quality<sup>60,62</sup>. In addition, marijuana might affect hormone levels, spermatogenesis, and mature sperm cells, as CB1 cannabinoid receptors are expressed in the anterior pituitary, testis, vas deferens, and human sperm cells<sup>61</sup>.

### Alcohol

According to a meta-analysis, alcohol consumption is negatively associated with semen volume, but not with other measures of semen quality<sup>50</sup>. In a large cross-sectional study of healthy men ( $n = 8,344$ ) from Europe and the USA, no consistent association between semen quality and alcohol consumption during the week preceding the study was observed<sup>63</sup>. However, in a cross-sectional study of young Danish men<sup>64</sup>, habitual drinking was associated with reduced sperm concentration, decreased total sperm count, and reduced percentage of spermatozoa with normal morphology, especially with a typical weekly intake of >25 units

of alcohol (1 unit of alcohol  $\approx 12$  g of ethanol)<sup>64</sup>. An autopsy study of men aged 35–69 years has suggested a dose-dependent association between spermatogenic arrest and alcohol consumption, and an average daily consumption of more than 80 g was associated with a significantly increased risk of spermatogenic arrest<sup>65</sup>. Alcohol consumption might also be associated with sperm aneuploidy<sup>54</sup>. Furthermore, prenatal exposure to alcohol has been negatively associated with sperm concentration in young men. In a Danish pregnancy cohort follow-up study, sons exposed to  $\geq 4.5$  drinks per week during pregnancy had approximately one-third lower adjusted mean sperm concentration than sons exposed to <1 drink per week ( $P = 0.04$ ), and an adverse effect of alcohol on fetal Sertoli cells has been suggested as a possible mechanism<sup>66</sup>.

Individual smoking and alcohol consumption habits might correlate and, when compared to nonsmokers and nondrinkers, a combination of smoking and alcohol consumption was associated with significantly increased sperm DNA fragmentation ( $P < 0.05$ ) in a study of men who were seeking semen analysis for fertility purposes in an IVF Unit<sup>67</sup>. Promotion of apoptosis was suggested as a

Table 2 | Sperm concentrations in young and fertile men in different geographical regions\*

Study	Area	Mean age (years)	n	Study period	Observed median sperm concentration (M/ml)	Adjusted median sperm concentration (M/ml)	Adjusted subgroup median sperm concentration (M/ml)
<b>Young men</b>							
Hart et al. <sup>37</sup> (2015)	Australia	20	365	≈2010	45	NA	NA
Andersen et al. <sup>29</sup> (2000)	Denmark	19.4	708	1996–1998	• 41 • 45 (abstinence >48 h, n = 521)	NA	NA
Jørgensen et al. <sup>27</sup> (2002)	Denmark	18.9	300	1997–1999	44	41 <sup>†</sup>	45 <sup>†a</sup>
Jørgensen et al. <sup>20</sup> (2012)	Denmark	19.4	4,867	1996–2010	45	NA	NA
		19.6	1,339	1996–2000	43		
		19.3	2,254	2001–2005	45		
		19.4	1,274	2006–2010	48		
Halling et al. <sup>35</sup> (2013)	Faroe islands	Median 24.0	481	2007–2010	40	NA	NA
			241	2007–2009	38		
			240	2009–2010	41		
Jørgensen et al. <sup>27</sup> (2002)	Estonia	18.8	104	1997–1999	62	57 <sup>†</sup>	63 <sup>†a</sup>
Punab et al. <sup>30</sup> (2002)	Estonia	20.4	79	1997–1999	64	67 <sup>†</sup>	66 <sup>†a</sup>
Jørgensen et al. <sup>27</sup> (2002)	Finland	18.8	324	1998–2000	61	54 <sup>†</sup>	53 <sup>†a</sup>
Jørgensen et al. <sup>2</sup> (2011)	Finland	18.8	338	1998–1999	60	67 <sup>#</sup>	65 <sup>#c</sup>
		19.0	382	2001–2003	54	60 <sup>#</sup>	58 <sup>#c</sup>
		19.1	138	2006	50	48 <sup>#</sup>	48 <sup>#c</sup>
Paasch et al. <sup>34</sup> (2008)	Germany (Hamburg)	19.7	334	2003–2004	49	46 <sup>†</sup>	NA
	Germany (Leipzig)	18.9	457	2003–2005	45	42 <sup>†</sup>	
Iwamoto et al. <sup>33</sup> (2013)	Japan (total)	21.3	1,559	1999–2003	59	59 <sup>  </sup>	62 <sup>  b</sup>
	Japan (Kawasaki)	20.8	658	1999–2000, 2002–2003	55	57 <sup>  </sup>	58 <sup>  b</sup>
	Japan (Osaka)	21.7	300	2002–2003	60	61 <sup>  </sup>	65 <sup>  b</sup>
	Japan (Kanazawa)	21.8	300	2002–2003	60	61 <sup>  </sup>	61 <sup>  b</sup>
	Japan (Nagasaki)	21.3	301	2002–2003	64	61 <sup>  </sup>	66 <sup>  b</sup>
Tsarev et al. <sup>31</sup> (2005)	Latvia	(≈19 years)	133	2001–2002	• 63 • 69 (abstinence ≥48 h, n = 100)	NA	NA
Punab et al. <sup>30</sup> (2002)	Lithuania	20.7	196	1997–1998	65	55 <sup>†</sup>	51 <sup>†a</sup>
Jørgensen et al. <sup>27</sup> (2002)	Norway	17.9	240	1998	53	41 <sup>†</sup>	42 <sup>†a</sup>
Fernandez et al. <sup>32</sup> (2012)	Spain	21.3	273	2001–2002	51	62 <sup>s</sup>	75 <sup>sa</sup>
Mendiola et al. <sup>1</sup> (2013)	Spain	19.2	215	2010–2011	44	NA	NA
Richthoff et al. <sup>28</sup> (2002)	Sweden	18.2	305	2000–2001	• 53.8 • 51.6 (men born and raised in Sweden, n = 248) • 55 (abstinence >48 h, n = 223)	NA	NA
Axelsson et al. <sup>19</sup> (2011),	Sweden	18.2	216	2000–2001	53	NA	NA
		18.0	295	2008–2010	56		
Mendiola et al. <sup>36</sup> (2014)	USA	19.7	221	2009–2010	52	NA	NR
<b>Fertile men (partners of pregnant women)</b>							
Stewart et al. <sup>45</sup> (2009)	Australia	35 (n = 225)	223	2000–2002	96	NA	NA
Jørgensen et al. <sup>26</sup> (2001)	Denmark	31.5	349	1996–1998	61	• 98** (winter) • 69** (summer)	NA



Table 2 (cont.) | Sperm concentrations in young and fertile men in different geographical regions\*

Study	Area	Mean age (years)	n	Study period	Observed median sperm concentration (M/ml)	Adjusted median sperm concentration (M/ml)	Adjusted subgroup median sperm concentration (M/ml)
<i>Fertile men (partners of pregnant women) (cont.)</i>							
Jørgensen <i>et al.</i> <sup>26</sup> (2001)	Finland	30.0	275	1996–1998	82	• 132** (winter) • 93** (summer)	NR
Jørgensen <i>et al.</i> <sup>26</sup> (2001)	France	32.0	207	1997–1998	74	• 103** (winter) • 73** (summer)	NR
Iwamoto <i>et al.</i> <sup>43</sup> (2006)	Japan (Kawasaki/Yokohama)	32.5	324	1998	–	53**	NR
Iwamoto <i>et al.</i> <sup>44</sup> (2013)	Japan (total)	31.7	792	1999–2002	84	84 <sup>†</sup>	NR
	Japan (Sapporo)	30.6	206	2000–2002	95	89 <sup>†</sup>	
	Japan (Osaka)	32.9	250	1999–2002	76	80 <sup>†</sup>	
	Japan (Kanazawa)	30.9	233	1999–2001	84	80 <sup>†</sup>	
	Japan (Fukuoka)	32.6	103	1999–2001	89	98 <sup>†</sup>	
Haugen <i>et al.</i> <sup>46</sup> (2006)	Norway	31	99 82 <sup>††</sup>	NA	70 70	–	NR
Jørgensen <i>et al.</i> <sup>26</sup> (2001)	Scotland, UK	32.5	251	1996–1997	77	• 119** (winter) • 84** (summer)	NR
Swan <i>et al.</i> <sup>40</sup> (2003)	USA (total)	31.3	493	1999–2001	–	NR	NR
	USA (California)	29.8	124	1999–2001	64.8		
	USA (Minnesota)	32.2	155	1999–2001	81.8		
	USA (Missouri)	30.7	176	1999–2001	53.5		
	USA (New York)	36.1	38	1999–2001	88.5		
Redmon <i>et al.</i> <sup>41</sup> (2013)	USA (total)	32	763	1999–2005	67	NR	NR
	USA (California)	–	182	1999–2002	Mean 55		
	USA (Iowa)	–	137	2002–2005	Mean 62		
	USA (Minnesota)	–	206	1999–2002	Mean 72		
	USA (Missouri)	–	201	1999–2002	Mean 48		
	USA (New York)	–	37	1999–2002	Mean 85		

\*Includes studies that have used similar study protocols. <sup>†</sup>Adjusted to the Danish laboratory level and to an abstinence period of  $\geq 96$  h. <sup>‡</sup>adjusted to a period of ejaculation abstinence of 96 h. <sup>§</sup> adjusted to a period of ejaculation abstinence of 96 h for a 21-year-old man. <sup>¶</sup>Adjusted to a period of ejaculation abstinence of 96 h for a 32-year-old man. <sup>||</sup> adjusted to a period of ejaculation abstinence of 95 h for a 19-year-old man. <sup>\*\*</sup>Calculated expected sperm concentration for a 30-year old man having ejaculation abstinence of 96 h. <sup>††</sup>Abstinence time 2–7 days and time to pregnancy  $<12$  cycles. <sup>†††</sup>Men not taking any medication, and without any previous or current andrological diseases including known fertility problems. <sup>††††</sup>Men with no history of reproductive problems. <sup>†††††</sup>Men without previous or current conditions that could potentially affect the semen quality.

potential mechanism<sup>67</sup>. Also in a study of men attending an andrology laboratory, those who smoked and drank alcohol had significantly lower semen volume, sperm concentration, and percentage of motile spermatozoa when compared with men without these two habits ( $P < 0.05$ )<sup>68</sup>.

### Obesity

According to the WHO, the worldwide prevalence of obesity in men has doubled between 1980 and 2008. In 2008, 10% of men in the world were obese (BMI  $\geq 30$  kg/m<sup>2</sup>), compared with 5% for men in 1980 (REF. 69). An extensive 2013 meta-analysis investigating the association between BMI and semen quality described a J-shaped association: overweight and obese men (BMI  $>25.0$ ) had a significantly increased risk of azoospermia

or oligozoospermia compared with men of normal weight (BMI 18.5–24.9)<sup>70</sup>. The meta-analysis included varying study populations, including men from the general population and male partners of subfertile couples. The association between obesity and semen quality is likely to be multifactorial and suggested possible mechanisms include changes in the hypothalamic–pituitary–gonadal axis, direct changes in spermatogenesis and Sertoli cell function, and an increase in scrotal temperature<sup>70</sup>. A cross-sectional study including a large proportion of overweight or obese men from fertility clinics could not confirm such an association<sup>71</sup>, but a prospective population-based cohort study on couples attempting to conceive suggested an association between high BMI and prevalence of low semen volume, low sperm concentration, and low total sperm




Figure 1 | **Median sperm concentration of young men in different geographical areas in studies using similar study protocols.** The study period and sperm concentration in million/ml (M/ml) is indicated for each area.

count<sup>72</sup>. Furthermore, some more recent studies have reported a negative association between BMI and different parameters of semen quality (semen volume, sperm concentration, total sperm count, motility, normal sperm morphology)<sup>37,73–75</sup>, but these results have not been replicated in all recent reports on the topic<sup>76,77</sup>.

### Stress

A 2011 meta-analysis by Li *et al.*<sup>50</sup> suggested that different forms of psychological stress might be associated with reduced sperm concentration, reduced progressive sperm motility, and abnormal sperm morphology. A similar conclusion was also reached by a large 2016 cross-sectional study of young Danish men from the general population, which suggested a negative association between self-reported stress (stress scores above an intermediate stress level of the study subjects) and sperm concentration, total sperm count, semen volume, and total number of morphologically normal spermatozoa<sup>78</sup>.

In the study, 20% of men had been distressed, 13% had had problems in relaxing, 12% had been irritated, and 16% had been tense all the time or a large part of the time during the preceding 4 weeks<sup>78</sup>. A cross-sectional study of the effect of stress in adult men from the general population in Northern California also reported a negative association between perceived stress score and sperm concentration and morphology<sup>79</sup> and a Chinese cross-sectional study on healthy men has also reported a negative association between psychological stress and total sperm count and percentage of sperm with normal morphology<sup>80</sup>. It has been speculated that stress could affect semen quality by inducing apoptosis of sensitive germ cells via high glucocorticoid levels<sup>78</sup>. A US study suggested no significant increase in prevalence of psychological distress between 2001 and 2012 (REF. 81) and, therefore, although stress might affect semen quality, increased psychological stress is unlikely to fully explain declining trends in semen quality.




Figure 2 | **Median sperm concentration of fertile men in different geographical areas in studies using similar study protocols.** The study period and sperm concentration in million/ml (M/ml) is indicated for each area. Seasonal differences can be seen in most areas, with sperm concentrations being considerably higher in the winter months.

### Endocrine disruptors

Endocrine-disrupting chemicals (EDCs) can influence semen quality and reproductive hormone levels<sup>82–84</sup>. EDCs include environmentally persistent organic pollutants (POPs), which remain intact for long periods and are toxic and widely distributed in the environment, and also other persistent and bioaccumulative chemicals, less persistent and less bioaccumulative compounds, current-use pesticides, pharmaceuticals and personal care product ingredients, metals and organometallic chemicals, natural hormones, and phytoestrogens<sup>82,83</sup>. Humans can be exposed to EDCs via food, water, inhaled air and particles, and via dermal contact.

**Persistent chemicals.** The persistence of many of the POPs — such as polychlorinated biphenyls (PCBs) — means that they are still major global pollutants even though they have been banned or regulated for years in many

countries<sup>82</sup>. Men exposed prenatally to high levels of PCBs and their pyrolytic products (mainly polychlorinated dibenzofurans) in so-called Yu-Cheng (oil-disease) exposure in Taiwan in 1979 had a significantly higher percentage of sperm with abnormal morphology and poorer sperm motility than unexposed controls<sup>85</sup>. Similarly, men exposed prenatally and postnatally via breast milk to 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) after the Seveso accident in Italy in 1976 had lower sperm concentrations, total sperm count, and sperm motility than control men, according to a study published in 2011<sup>86</sup>. Men exposed to TCDD at Seveso when they were 1–9 years of age also had reduced sperm concentrations and motility when compared to a control group, whereas such an association was not observed among men exposed to TCDD from the Seveso accident during puberty or in adulthood<sup>87</sup>. *In utero* exposure to background levels of perfluorooctanoic acid (which is a very stable synthetic



surfactant) has been associated negatively with sperm concentration and total sperm count, whereas no significant association was seen between *in utero* exposure to background levels of perfluorooctanesulfonic acid, PCBs or dichlorodiphenyldichloroethylene (p,p'-DDE) and semen quality in adulthood<sup>88,89</sup>, which suggests that the levels associated with adverse effects varies between chemicals. A detailed review concluded that exposure to PCBs and perfluorinated compounds during adult life seems to be negatively associated with sperm motility and sperm morphology, respectively<sup>84</sup>. Adult exposure levels to PCBs and p,p'-DDE in men have also been positively associated with time to pregnancy<sup>90</sup>.

**Nonpersistent chemicals.** Prenatal exposure to phthalates has been associated negatively with semen volume of young men<sup>91</sup>. Adult exposure to phthalates has also been negatively associated with semen quality. A meta-analysis suggested that exposure to phthalates is associated with decreased sperm concentration, motility and increased sperm DNA damage<sup>92</sup>. Subsequent studies in men representing the general population have also suggested some adverse associations between semen quality and adult exposure to phthalates<sup>93,94</sup>. Increased oxidative stress has been suggested as a possible mechanism for reduced sperm motility related to adult phthalate exposure<sup>93</sup>. However, current literature suggests that the role of adult exposure to phthalates at environmental levels in determining the classical sperm parameters is likely to be small<sup>94</sup>. Bisphenol A, which is used in the production of plastics and epoxy resin liners of canned foods, is a non-persistent chemical that does not bioaccumulate<sup>82</sup>. In a recent review on bisphenol A and male reproductive health it was concluded that, although negative associations between adult urine levels of bisphenol A and various semen quality parameters have been described in some epidemiological studies, the results remain limited and inconclusive<sup>95</sup>. One limitation with studies of chemical exposures is the lack of capacity to consider several concomitant exposures — that is, the mixture effects. This might lead to underestimation of the risks, as individual chemicals might not show a significant association with an adverse outcome, although they might contribute to harmful combined exposure. Animal studies have demonstrated cumulative dose-additive adverse effects of mixtures of antiandrogenic chemicals on male reproductive tract development<sup>96,97</sup>.

#### Use of mobile phone and wireless Internet

Meta-analyses including both *in vivo* and *in vitro* studies have suggested that exposure to mobile phones is associated with reduced sperm motility and sperm viability<sup>98–100</sup>. Increased reactive oxygen species production and increased DNA fragmentation in sperm have been suggested as possible mechanisms for such adverse effects. An association of mobile phone exposure with sperm concentration has been observed in the meta-analysis of animal studies, but the data are equivocal in human studies<sup>98–100</sup> and The National Toxicology Program is currently running a comprehensive study on the effects of mobile phone exposure on health<sup>101</sup>.

One *in vitro* study has suggested a negative association between the use of wireless-internet-connected laptops and sperm motility and also suggested increased DNA fragmentation, potentially via a nonthermal effect<sup>102</sup>.

#### Effect of semen quality on fecundity

##### Semen quality and time to pregnancy

The correlation between semen quality and fecundity is complicated, and no accepted threshold values exist for any semen variables on an individual level<sup>103</sup>. In a multi-centre study conducted by the US National Cooperative Reproductive Medicine Network, sperm concentrations of  $48 \times 10^6/\text{ml}$  and  $13.5 \times 10^6/\text{ml}$ , percentage of motile sperm of 63% and 32%, and percentage of sperm with normal morphology of 12% and 9% defined the lower end of the fertile range and the upper end of the subfertile range, respectively<sup>103</sup>. The men whose sperm variables are in between these limits fall in the grey zone between fertility and subfertility. In a study of couples trying to conceive for the first time, time to pregnancy started to increase when sperm concentration was  $<40 \times 10^6/\text{ml}$ <sup>104</sup>. Percentage of sperm with normal morphology showed also a significant positive association with time to pregnancy<sup>104</sup>. In a large multinational European study of male partners of pregnant women, time to pregnancy increased when sperm concentration was  $<55 \times 10^6/\text{ml}$  or total sperm count was  $<145 \times 10^6$ , or if the proportion of morphologically normal sperm (using strict criteria suggested by Menkveld *et al.*<sup>105</sup>) was  $<19\%$ <sup>106</sup>. A prospective US study reported a decline in pregnancy rates in otherwise healthy couples when sperm concentration is  $<30 \times 10^6/\text{ml}$ <sup>107</sup>. The latter two studies included also couples who had previously conceived successfully. A remarkably high proportion of young men all over the world have sperm concentrations below the aforementioned limits (TABLE 2). Percentage of morphologically normal spermatozoa was found to be associated with reduced time to pregnancy, and the percentage of spermatozoa with coiled tail was associated with increased time to pregnancy in a subsequent US study<sup>108,109</sup>. Additional variables, beyond those measured in a conventional semen analysis, have been associated with fecundity. According to a meta-analysis, a high level of sperm DNA fragmentation is associated with decreased pregnancy rate after *in vitro* fertilization (IVF) and increased miscarriage rate after intracytoplasmic sperm injection (ICSI)<sup>110</sup>. Furthermore, studies on presumably fertile couples attempting to achieve a pregnancy and a separate study on couples attempting to conceive for the first time have described an association between sperm chromatin integrity and time to conception<sup>111,112</sup>.

#### Updated WHO criteria for normal semen quality

In 1980, the WHO published a “Laboratory Manual for Examination of Human Semen and Semen-Cervical Mucus Interaction” in response to a growing need for the standardization of procedures for the examination of human semen<sup>113</sup>. Since then the manual has been updated and revised four times and the latest edition, from 2010, listed lower reference values for semen variables<sup>114</sup>. The new values were calculated according to the

fifth centile of fertile men and represent semen quality of recent fathers. The lower limit for normal sperm concentration was reduced from  $20 \times 10^6/\text{ml}$  to  $15 \times 10^6/\text{ml}$ ; in the 1940s the normal limit was considered to be  $60 \times 10^6/\text{ml}$ <sup>115</sup>. Whether these changes in the reference values reflect improved methodology and knowledge or an actual decline in semen quality at the population level is matter of discussion. The current WHO reference levels do not distinguish between normal and abnormal testis function from a biological point of view, but might instead serve as a tool to identify men who might require fertility treatment in order to procreate.

## Conclusions

Studies published in this millennium have clearly shown regional differences in semen quality. In population-based studies of young men, semen quality in many countries is at a level that raises concerns for their fecundity, with a considerable proportion of men having sperm concentrations associated with prolonged time to first pregnancy. Semen quality has certainly declined or stabilized to this low level during the past few decades, possibly owing to modern lifestyle factors, including exposure to environmental chemicals. The search to identify and avoid these preventable causes of reproductive problems continues.

- Mendiola, J. *et al.* Sperm counts may have declined in young university students in southern Spain. *Andrology* **1**, 408–413 (2013).
- Jørgensen, N. *et al.* Recent adverse trends in semen quality and testis cancer incidence among Finnish men. *Int. J. Androl.* **34**, e37–48 (2011).
- Lackner, J. *et al.* Constant decline in sperm concentration in infertile males in an urban population: experience over 18 years. *Fertil. Steril.* **84**, 1657–1661 (2005).
- Sripada, S. *et al.* Trends in semen parameters in the northeast of Scotland. *J. Androl.* **28**, 313–319 (2007).
- Feki, N. C. *et al.* Semen quality decline among men in infertile relationships: experience over 12 years in the south of Tunisia. *J. Androl.* **30**, 541–547 (2009).
- Geoffroy-Siraudin, C. *et al.* Decline of semen quality among 10 932 males consulting for couple infertility over a 20-year period in Marseille, France. *Asian J. Androl.* **14**, 584–590 (2012).
- Rolland, M., Le Moal, J., Wagner, V., Royère, D. & De Mouzon, J. Decline in semen concentration and morphology in a sample of 26,609 men close to general population between 1989 and 2005 in France. *Hum. Reprod.* **28**, 462–470 (2013).
- Adiga, S. K., Jayaraman, V., Kalthur, G., Upadhyay, D. & Kumar, P. Declining semen quality among south Indian infertile men: a retrospective study. *J. Hum. Reprod. Sci.* **1**, 15–18 (2008).
- Romero-Otero, J. *et al.* Semen quality assessment in fertile men in Madrid during the last 3 decades. *Urology* **85**, 1333–1338 (2015).
- Jiang, M. *et al.* Semen quality evaluation in a cohort of 28213 adult males from Sichuan area of south-west China. *Andrologia* **46**, 842–847 (2014).
- Haimov-Kochman, R. *et al.* Is the quality of donated semen deteriorating? Findings from a 15 year longitudinal analysis of weekly sperm samples. *Isr. Med. Assoc. J.* **14**, 372–377 (2012).
- Rao, M. *et al.* Evaluation of semen quality in 1808 university students, from Wuhan, central China. *Asian J. Androl.* **17**, 111–116 (2015).
- Wang, L. *et al.* Decline of semen quality among Chinese sperm bank donors within 7 years (2008–2014). *Asian J. Androl.* <http://dx.doi.org/10.4103/1008-682X.179533> (2016).
- Shine, R., Peek, J. & Birdsall, M. Declining sperm quality in New Zealand over 20 years. *N. Z. Med. J.* **121**, 50–56 (2008).
- Spligart, C. *et al.* Semen variation in a population of fertile donors: evaluation in a French centre over a 34-year period. *Int. J. Androl.* **35**, 467–474 (2012).
- Centola, G. M., Blanchard, A., Demick, J., Li, S. & Eisenberg, M. L. Decline in sperm count and motility in young adult men from 2003 to 2013: observations from a U. S. sperm bank. *Andrology* **4**, 270–276 (2016).
- Huang, C. *et al.* Decline in semen quality among 30,636 young Chinese men from 2001 to 2015. *Fertil. Steril.* <http://dx.doi.org/10.1016/j.fertnstert.2016.09.035> (2016).
- Chen, Z. *et al.* Temporal trends in human semen parameters in New England in the United States, 1989–2000. *Arch. Androl.* **49**, 369–374 (2003).
- Axelsson, J., Rylander, L., Rignell-Hydbom, A. & Giwercman, A. No secular trend over the last decade in sperm counts among Swedish men from the general population. *Hum. Reprod.* **26**, 1012–1016 (2011).
- Jørgensen, N. *et al.* Human semen quality in the new millennium: a prospective cross-sectional population-based study of 4867 men. *BMJ Open* **2**, e000990 (2012).
- Costello, M. F., Sjöblom, P., Haddad, Y., Steigrad, S. J. & Bosch, E. G. No decline in semen quality among potential sperm donors in Sydney, Australia, between 1983 and 2001. *J. Assist. Reprod. Genet.* **19**, 284–290 (2002).
- Mukhopadhyay, D. *et al.* Semen quality and age-specific changes: a study between two decades on 3,729 male partners of couples with normal sperm count and attending an andrology laboratory for infertility-related problems in an Indian city. *Fertil. Steril.* **93**, 2247–2254 (2010).
- Marimuthu, P., Kapilashrami, M. C., Misro, M. M. & Singh, G. Evaluation of trend in semen analysis for 11 years in subjects attending a fertility clinic in India. *Asian J. Androl.* **5**, 221–225 (2003).
- Birdsall, M. A., Peek, J. & Valiathan, S. Sperm quality in New Zealand: is the downward trend continuing? *N. Z. Med. J.* **128**, 50–56 (2015).
- Johnson, S. L., Dunleavy, J., Gemmell, N. J. & Nakagawa, S. Consistent age-dependent declines in human semen quality: a systematic review and meta-analysis. *Ageing Res. Rev.* **19**, 22–33 (2015).
- Jørgensen, N. *et al.* Regional differences in semen quality in Europe. *Hum. Reprod.* **16**, 1012–1019 (2001).
- Jørgensen, N. *et al.* East-West gradient in semen quality in the Nordic-Baltic area: a study of men from the general population in Denmark, Norway, Estonia and Finland. *Hum. Reprod.* **17**, 2199–2208 (2002).
- Richthoff, J., Rylander, L., Hagmar, L., Malm, J. & Giwercman, A. Higher sperm counts in southern Sweden compared with Denmark. *Hum. Reprod.* **17**, 2468–2473 (2002).
- Andersen, A. G. *et al.* High frequency of sub-optimal semen quality in an unselected population of young men. *Hum. Reprod.* **15**, 366–372 (2000).
- Punab, M. *et al.* Regional differences in semen qualities in the Baltic region. *Int. J. Androl.* **25**, 243–252 (2002).
- Tsarev, I., Gagonin, V., Giwercman, A. & Erenpreiss, J. Sperm concentration in Latvian military conscripts as compared with other countries in the Nordic-Baltic area. *Int. J. Androl.* **28**, 208–214 (2005).
- Fernandez, M. F. *et al.* Semen quality and reproductive hormone levels in men from southern Spain. *Int. J. Androl.* **35**, 1–10 (2012).
- Iwamoto, T. *et al.* Semen quality of 1559 young men from four cities in Japan: a cross-sectional population-based study. *BMJ Open* **3**, e002222 (2013).
- Paasch, U. *et al.* Semen quality in sub-fertile range for a significant proportion of young men from the general German population: a co-ordinated, controlled study of 791 men from Hamburg and Leipzig. *Int. J. Androl.* **31**, 93–102 (2008).
- Halling, J. *et al.* Semen quality and reproductive hormones in Faroese men: a cross-sectional population-based study of 481 men. *BMJ Open* **3**, e001946 (2013).
- Mendiola, J. *et al.* Reproductive parameters in young men living in Rochester, New York. *Fertil. Steril.* **101**, 1064–1071 (2014).
- Hart, R. J. *et al.* Testicular function in a birth cohort of young men. *Hum. Reprod.* **30**, 2713–2724 (2015).
- Perheentupa, A. *et al.* Semen quality improves marginally during young adulthood: a longitudinal follow-up study. *Hum. Reprod.* **31**, 502–510 (2016).
- Carlsen, E., Swan, S. H., Petersen, J. H. & Skakkebaek, N. E. Longitudinal changes in semen parameters in young Danish men from the Copenhagen area. *Hum. Reprod.* **20**, 942–949 (2005).
- Swan, S. H. *et al.* Geographic differences in semen quality of fertile U. S. males. *Environ. Health Perspect.* **111**, 414–420 (2003).
- Redmon, J. B. *et al.* Semen parameters in fertile US men: the Study for Future Families. *Andrology* **1**, 806–814 (2013).
- Swan, S. H. *et al.* Semen quality in relation to biomarkers of pesticide exposure. *Environ. Health Perspect.* **111**, 1478–1484 (2003).
- Iwamoto, T. *et al.* Semen quality of 324 fertile Japanese men. *Hum. Reprod.* **21**, 760–765 (2006).
- Iwamoto, T. *et al.* Semen quality of fertile Japanese men: a cross-sectional population-based study of 792 men. *BMJ Open* **3**, e002223 (2013).
- Stewart, T. M. *et al.* Associations between andrological measures, hormones and semen quality in fertile Australian men: inverse relationship between obesity and sperm output. *Hum. Reprod.* **24**, 1561–1568 (2009).
- Haugen, T. B., Egeland, T. & Magnus, O. Semen parameters in Norwegian fertile men. *J. Androl.* **27**, 66–71 (2006).
- Skakkebaek, N. E., Rajpert-De Meyts, E. & Main, K. M. Testicular dysgenesis syndrome: an increasingly common developmental disorder with environmental aspects. *Hum. Reprod.* **16**, 972–978 (2001).
- Skakkebaek, N. E. *et al.* Male reproductive disorders and fertility trends: influences of environment and genetic susceptibility. *Physiol. Rev.* **96**, 55–97 (2016).
- Vine, M. F., Margolin, B. H., Morrison, H. I. & Hulka, B. S. Cigarette smoking and sperm density: a meta-analysis. *Fertil. Steril.* **61**, 35–43 (1994).
- Li, Y., Lin, H. & Cao, J. Association between socio-psychological behavioral factors and male semen quality: systematic review and meta-analyses. *Fertil. Steril.* **95**, 116–123 (2011).
- Sharma, R., Harlev, A., Agarwal, A. & Esteves, S. C. Cigarette smoking and semen quality: a new meta-analysis examining the effect of the 2010 World Health Organization laboratory methods for the examination of human semen. *Eur. Urol.* **70**, 635–645 (2016).
- Rubes, J. *et al.* Smoking cigarettes is associated with increased sperm disomy in teenage men. *Fertil. Steril.* **70**, 715–723 (1998).
- Taha, E. A., Ez-Aldin, A. M., Sayed, S. K., Ghandour, N. M. & Mostafa, T. Effect of smoking on sperm vitality, DNA integrity, seminal oxidative stress, zinc in fertile men. *Urology* **80**, 822–825 (2012).
- Robbins, W. A. *et al.* Effect of lifestyle exposures on sperm aneuploidy. *Cytogenet. Genome Res.* **111**, 371–377 (2005).
- Storgaard, L. *et al.* Does smoking during pregnancy affect sons' sperm counts? *Epidemiology* **14**, 278–286 (2003).
- Jensen, T. K. *et al.* Association of *in utero* exposure to maternal smoking with reduced semen quality and testis size in adulthood: a cross-sectional study of 1,770 young men from the general population in five European countries. *Am. J. Epidemiol.* **159**, 49–58 (2004).
- Jensen, M. S., Mabeck, L. M., Toft, G., Thulstrup, A. M. & Bonde, J. P. Lower sperm counts following prenatal tobacco exposure. *Hum. Reprod.* **20**, 2559–2566 (2005).

58. Ramlau-Hansen, C. H. *et al.* Is prenatal exposure to tobacco smoking a cause of poor semen quality? A follow-up study. *Am. J. Epidemiol.* **165**, 1372–1379 (2007).
59. Ravnborg, T. L. *et al.* Prenatal and adult exposures to smoking are associated with adverse effects on reproductive hormones, semen quality, final height and body mass index. *Hum. Reprod.* **26**, 1000–1011 (2011).
60. Virtanen, H. E., Sadov, S. & Toppari, J. Prenatal exposure to smoking and male reproductive health. *Curr. Opin. Endocrinol. Diabetes Obes.* **19**, 228–232 (2012).
61. Gundersen, T. D. *et al.* Association between use of marijuana and male reproductive hormones and semen quality: a study among 1,215 healthy young men. *Am. J. Epidemiol.* **182**, 473–481 (2015).
62. Dai, J. B., Wang, Z. X. & Qiao, Z. D. The hazardous effects of tobacco smoking on male fertility. *Asian J. Androl.* **17**, 954–960 (2015).
63. Jensen, T. K. *et al.* Alcohol and male reproductive health: a cross-sectional study of 8344 healthy men from Europe and the USA. *Hum. Reprod.* **29**, 1801–1809 (2014).
64. Jensen, T. K. *et al.* Habitual alcohol consumption associated with reduced semen quality and changes in reproductive hormones; a cross-sectional study among 1221 young Danish men. *BMJ Open* **4**, e005462 (2014).
65. Pajarinen, J. *et al.* Moderate alcohol consumption and disorders of human spermatogenesis. *Alcohol Clin. Exp. Res.* **20**, 332–337 (1996).
66. Ramlau-Hansen, C. H. *et al.* Maternal alcohol consumption during pregnancy and semen quality in the male offspring: two decades of follow-up. *Hum. Reprod.* **25**, 2340–2345 (2010).
67. Anifandis, G. *et al.* The impact of cigarette smoking and alcohol consumption on sperm parameters and sperm DNA fragmentation (SDF) measured by Halosperm®. *Arch. Gynecol. Obstet.* **290**, 777–782 (2014).
68. Martini, A. C. *et al.* Effects of alcohol and cigarette consumption on human seminal quality. *Fertil. Steril.* **82**, 374–377 (2004).
69. WHO. Obesity. Situation and trends. *Global Health Observatory (GHO) data* [online], [http://www.who.int/gho/ncd/risk\\_factors/obesity\\_text/en/](http://www.who.int/gho/ncd/risk_factors/obesity_text/en/) (2016).
70. Sermondade, N. *et al.* BMI in relation to sperm count: an updated systematic review and collaborative meta-analysis. *Hum. Reprod. Update* **19**, 221–231 (2013).
71. Macdonald, A. A., Stewart, A. W. & Farquhar, C. M. Body mass index in relation to semen quality and reproductive hormones in New Zealand men: a cross-sectional study in fertility clinics. *Hum. Reprod.* **28**, 3178–3187 (2013).
72. Eisenberg, M. L. *et al.* The relationship between male BMI and waist circumference on semen quality: data from the LIFE study. *Hum. Reprod.* **29**, 193–200 (2014).
73. Tsao, C. W. *et al.* Exploration of the association between obesity and semen quality in a 7630 male population. *PLoS ONE* **10**, e0119458 (2015).
74. Andersen, J. M. *et al.* Body mass index is associated with impaired semen characteristics and reduced levels of anti-Müllerian hormone across a wide weight range. *PLoS ONE* **10**, e0130210 (2015).
75. Belloc, S. *et al.* High body mass index has a deleterious effect on semen parameters except morphology: results from a large cohort study. *Fertil. Steril.* **102**, 1268–1273 (2014).
76. Ehala-Aleksejev, K. & Punab, M. The different surrogate measures of adiposity in relation to semen quality and serum reproductive hormone levels among Estonian fertile men. *Andrology* **3**, 225–234 (2015).
77. Lu, J. C. *et al.* Body mass index, waist-to-hip ratio, waist circumference and waist-to-height ratio cannot predict male semen quality: a report of 1231 subfertile Chinese men. *Andrologia* **47**, 1047–1054 (2015).
78. Nordkap, L. *et al.* Psychological stress and testicular function: a cross-sectional study of 1,215 Danish men. *Fertil. Steril.* **105**, 174–187.e2 (2016).
79. Janevic, T. *et al.* Effects of work and life stress on semen quality. *Fertil. Steril.* **102**, 530–538 (2014).
80. Li, Y. *et al.* Socio-psychobehavioural factors associated with male semen quality in China: results from 1346 healthy men in Chongqing. *J. Fam. Plann. Reprod. Health Care* **39**, 102–110 (2013).
81. Mojtabai, R. & Jorm, A. F. Trends in psychological distress, depressive episodes and mental health treatment-seeking in the United States: 2001–2012. *J. Affect. Disord.* **174**, 556–561 (2015).
82. Bergman, Å., Heindel, J., Jobling, S., Kidd, K. & Zoeller, R. *State of the science of endocrine disrupting chemicals 2012*. (United Nations Environment Programme and the World Health Organization, 2013).
83. Gore, A. C. *et al.* EDC-2: the Endocrine Society's second scientific statement on endocrine-disrupting chemicals. *Endocr. Rev.* **36**, E1–E150 (2015).
84. Vested, A., Gierwman, A., Bonde, J. P. & Toft, G. Persistent organic pollutants and male reproductive health. *Asian J. Androl.* **16**, 71–80 (2014).
85. Guo, Y. L., Hsu, P. C., Hsu, C. C. & Lambert, G. H. Semen quality after prenatal exposure to polychlorinated biphenyls and dibenzofurans. *Lancet* **356**, 1240–1241 (2000).
86. Mocarelli, P. *et al.* Perinatal exposure to low doses of dioxin can permanently impair human semen quality. *Environ. Health Perspect.* **119**, 713–718 (2011).
87. Mocarelli, P. *et al.* Dioxin exposure, from infancy through puberty, produces endocrine disruption and affects human semen quality. *Environ. Health Perspect.* **116**, 70–77 (2008).
88. Vested, A. *et al.* Associations of *in utero* exposure to perfluorinated alkyl acids with human semen quality and reproductive hormones in adult men. *Environ. Health Perspect.* **121**, 453–458 (2013).
89. Vested, A. *et al.* *In utero* exposure to persistent organochlorine pollutants and reproductive health in the human male. *Reproduction* **148**, 635–646 (2014).
90. Buck Louis, G. M. *et al.* Persistent environmental pollutants and couple fecundity: the LIFE study. *Environ. Health Perspect.* **121**, 231–236 (2013).
91. Axelsson, J. *et al.* Prenatal phthalate exposure and reproductive function in young men. *Environ. Res.* **138**, 264–270 (2015).
92. Cai, H. *et al.* Human urinary/seminal phthalates or their metabolite levels and semen quality: a meta-analysis. *Environ. Res.* **142**, 486–494 (2015).
93. Phthalate exposure and reproductive parameters in young men from the general Swedish population. *Environ. Int.* **85**, 54–60 (2015).
94. Thurston, S. W. *et al.* Phthalate exposure and semen quality in fertile US men. *Andrology* **4**, 623–628 (2016).
95. Minguez-Alarcón, L., Hauser, R. & Gaskins, A. J. Effects of bisphenol A on male and couple reproductive health: a review. *Fertil. Steril.* **106**, 864–870 (2016).
96. Christiansen, S. *et al.* Synergistic disruption of external male sex organ development by a mixture of four antiandrogens. *Environ. Health Perspect.* **117**, 1839–1846 (2009).
97. Rider, C. V., Furr, J. R., Wilson, V. S. & Gray, L. E. J. Cumulative effects of *in utero* administration of mixtures of reproductive toxicants that disrupt common target tissues via diverse mechanisms of toxicity. *Int. J. Androl.* **33**, 443–462 (2010).
98. Dama, M. S. & Bhat, M. N. Mobile phones affect multiple sperm quality traits: a meta-analysis. *F1000Res* **2**, 40 (2013).
99. Adams, J. A., Galloway, T. S., Mondal, D., Esteves, S. C. & Mathews, F. Effect of mobile telephones on sperm quality: a systematic review and meta-analysis. *Environ. Int.* **70**, 106–112 (2014).
100. Liu, K. *et al.* Association between mobile phone use and semen quality: a systemic review and meta-analysis. *Andrology* **2**, 491–501 (2014).
101. NTP. Cell phones. *National Toxicology Program* [online], <https://ntp.niehs.nih.gov/results/areas/cellphones/> (2016).
102. Avendaño, C., Mata, A., Sanchez Sarmiento, C. A. & Doncel, G. F. Use of laptop computers connected to internet through Wi-Fi decreases human sperm motility and increases sperm DNA fragmentation. *Fertil. Steril.* **97**, 39–45.e32 (2012).
103. Guzik, D. S. *et al.* Sperm morphology, motility, and concentration in fertile and infertile men. *N. Engl. J. Med.* **345**, 1388–1393 (2001).
104. Bonde, J. P. *et al.* Relation between semen quality and fertility: a population-based study of 430 first-pregnancy planners. *Lancet* **352**, 1172–1177 (1998).
105. Menkveld, R., Stander, F. S., Kotze, T. J., Kruger, T. F. & van Zyl, J. A. The evaluation of morphological characteristics of human spermatozoa according to stricter criteria. *Hum. Reprod.* **5**, 586–592 (1990).
106. Slama, R. *et al.* Time to pregnancy and semen parameters: a cross-sectional study among fertile couples from four European cities. *Hum. Reprod.* **17**, 503–515 (2002).
107. Zinaman, M. J., Brown, C. C., Selevan, S. G. & Clegg, E. D. Semen quality and human fertility: a prospective study with healthy couples. *J. Androl.* **21**, 145–153 (2000).
108. Buck Louis, G. M. *et al.* Semen quality and time to pregnancy: the Longitudinal Investigation of Fertility and the Environment Study. *Fertil. Steril.* **101**, 453–462 (2014).
109. Patel, C. J., Sundaram, R. & Buck Louis, G. M. A data-driven search for semen-related phenotypes in conception delay. *Andrology* <http://dx.doi.org/10.1111/andr.12288> (2016).
110. Zhao, J., Zhang, Q., Wang, Y. & Li, Y. Whether sperm deoxyribonucleic acid fragmentation has an effect on pregnancy and miscarriage after *in vitro* fertilization/intracytoplasmic sperm injection: a systematic review and meta-analysis. *Fertil. Steril.* **102**, 998–1005.e1008 (2014).
111. Evenson, D. P. *et al.* Utility of the sperm chromatin structure assay as a diagnostic and prognostic tool in the human fertility clinic. *Hum. Reprod.* **14**, 1039–1049 (1999).
112. Spanò, M. *et al.* Sperm chromatin damage impairs human fertility. *Fertil. Steril.* **73**, 43–50 (2000).
113. WHO. *WHO laboratory manual for the examination of human semen and sperm – cervical mucus interaction*. (Press Concern, 1980).
114. WHO. *WHO laboratory manual for the examination and processing of human semen*. 5th edn, (WHO press, 2010).
115. MacLeod, J. & Heim, L. M. Characteristics and variations in semen specimens in 100 normal young men. *J. Urol.* **54**, 474–482 (1945).

## Acknowledgements

The authors thank Dr. Wiwat Rodprasert for drawing the figures and Ms. Johanna Järvi for technical assistance with this article. This work was supported by the European Commission (FP7), the Academy of Finland, Sigrid Juselius Foundation, Novo Nordisk Foundation, Turku University Hospital Special Governmental Research Fund, and The Research Fund of Rigshospitalet (grant no. R42-A1326).

## Author contributions

All authors researched data for article, made substantial contributions to discussion of content, wrote the article, and reviewed and edited the manuscript before submission.

## Competing interests

J.T. has received honoraria for speaking and acted as a consultant for Merck and Mylan. The other authors declare no competing interests.