

The Biology of Vascular Endothelial Growth Factor

NAPOLEONE FERRARA AND TERRI DAVIS-SMYTH

Department of Cardiovascular Research, Genentech, Inc., South San Francisco, California 94080

- I. Introduction
- II. Biological Activities of VEGF
- III. Organization of the VEGF Gene
- IV. Properties of the VEGF Isoforms
- V. Regulation of VEGF Gene Expression
 - A. Hypoxia
 - B. Cytokines
 - C. Differentiation and transformation
- VI. The VEGF Receptors
 - A. Characterization and distribution of VEGF-binding sites
 - B. The Flt-1 and Flk-1/KDR tyrosine kinases
 1. Binding characteristics
 2. Signal transduction
 3. Regulation
 4. Structural requirements for ligand binding in Flt-1 and Flk-1/KDR
 5. VEGF determinants for binding Flt-1 and Flk-1/KDR
- VII. VEGF-Related Molecules
- VIII. Role of VEGF and Its Receptors in Physiological Angiogenesis
 - A. Distribution of VEGF, Flk-1/KDR and Flt-1 mRNA
 - B. Analysis of Flk-1/KDR, Flt-1 and VEGF gene knockouts
- IX. Role of VEGF in Pathological Angiogenesis
 - A. Tumor angiogenesis
 1. Expression of VEGF in human tumors
 2. Inhibition of VEGF action *in vivo*
 - B. Intraocular neovascular syndromes
 - C. Other pathological conditions
- X. Therapeutic Applications of VEGF-Induced Angiogenesis
- XI. Perspectives

I. Introduction

THE development of a vascular supply is a fundamental requirement for organ development and differentiation during embryogenesis (1, 2) as well as for wound healing and reproductive functions in the adult (3, 4). Angiogenesis is also implicated in the pathogenesis of a variety of disorders: proliferative retinopathies, age-related macular degeneration (AMD), tumors, rheumatoid arthritis, psoriasis, etc. (3, 4). In the case of proliferative retinopathies and AMD, the new blood vessels are directly responsible for many of the

destructive events characteristic of these conditions. Leakage and bleeding, followed by organization of the clot and fibrosis, may ultimately lead to retinal detachment or irreversible damage to the macula (5). Conversely, tumor-associated neovascularization, by establishing continuity with the systemic circulation, allows the tumor cells to express their critical growth advantage and also facilitates metastatic spreading (3, 4). Accordingly, a correlation has been observed between density of microvessels in primary breast carcinoma sections, nodal metastases, and survival (6–8). Similarly, a correlation has been reported between vascularity and invasive behavior in a variety of other tumors (9–12). These findings led several investigators to conclude that the number of vessels in tumor sections is an independent predictor of outcome in cancer patients (9–12).

The search for potential regulators of angiogenesis has yielded numerous candidates: acidic fibroblast growth factor (aFGF), basic fibroblast growth factor (bFGF), transforming growth factor- α (TGF- α), TGF- β , hepatocyte growth factor, tumor necrosis factor- α (TNF- α), angiogenin, interleukin-8 (IL-8), etc. (13, 14). Although these molecules are able to promote angiogenesis, at least in certain model systems, it has been difficult to correlate such activity with the physiological or pathological regulation of blood vessel growth.

Work done by several laboratories over the last few years has elucidated the pivotal role of vascular endothelial growth factor (VEGF) in the regulation of normal and abnormal angiogenesis (15). In particular, the recent finding that the loss of even a single VEGF allele results in embryonic lethality points to an irreplaceable role played by this factor in the development and differentiation of the vascular system (16, 17). Furthermore, VEGF-induced angiogenesis has been shown to result in a therapeutic effect in animal models of coronary (18–20) or limb (21–23) ischemia and, most recently, in a human patient affected by critical leg ischemia (24).

II. Biological Activities of VEGF

VEGF is a potent mitogen (ED₅₀ 2–10 pM) for micro- and macrovascular endothelial cells derived from arteries, veins, and lymphatics, but it is devoid of consistent and appreciable mitogenic activity for other cell types (25–31). The denomination of VEGF was proposed to emphasize such narrow target cell specificity (25, 26). VEGF promotes angiogenesis in tridimensional *in vitro* models, inducing confluent microvascular endothelial cells to invade collagen gels and form capillary-like structures (32). These studies provided evidence for a potent synergism between VEGF and bFGF in the induction of this effect (32). Also, VEGF induced sprouting

Address reprint requests to: Napoleone Ferrara, M.D., Department of Cardiovascular Research, Genentech, Inc., 460 Point San Bruno Boulevard, South San Francisco, California 94080.

from rat aortic rings embedded in a collagen gel (33). This model emphasizes the specificity of VEGF, as the proliferation induced by this growth factor consisted almost exclusively of vascular endothelial cells. In contrast, insulin-like growth factor-I (IGF-I) or platelet-derived growth factor (PDGF) induced endothelial cell sprouting accompanied by extensive fibroblastic proliferation (33). VEGF also elicits a strong angiogenic response in a variety of *in vivo* models including the chick chorioallantoic membrane (26, 29), the rabbit cornea (34), the primate iris (35), the rabbit bone (27), etc.

VEGF induces expression of the serine proteases urokinase-type and tissue-type plasminogen activators (PA) and also PA inhibitor 1 (PAI-1) in cultured bovine microvascular endothelial cells (36). Moreover, VEGF increases expression of the metalloproteinase interstitial collagenase in human umbilical vein endothelial cells but not in dermal fibroblasts (37). The coinduction of PA and collagenase by VEGF is consistent with a prodegradative environment that facilitates migration and sprouting of endothelial cells. Pepper and Montesano (38) proposed that PAI-1 provides a negative regulatory step that serves to balance the proteolytic process. Other studies have shown that VEGF promotes expression of urokinase receptor (uPAR) in vascular endothelial cells (39). Considering that the PA-plasmin system and in particular the interaction of uPA with uPAR is an important element in the chain of cellular processes that mediate cellular invasion and tissue remodeling (40), these findings are consistent with the proangiogenic activities of VEGF.

VEGF is known also as vascular permeability factor (VPF) based on its ability to induce vascular leakage in the guinea pig skin (41, 42). Dvorak and colleagues (43, 44) proposed that an increase in microvascular permeability is a crucial step in angiogenesis associated with tumors and wounds. According to this hypothesis, a major function of VPF/VEGF in the angiogenic process is the induction of plasma protein leakage. This effect would result in the formation of an extravascular fibrin gel, a substrate for endothelial and tumor cell growth. Recent studies have also suggested that VEGF may be a factor that induces fenestrations in endothelial cells (45). Topical administration of VEGF acutely resulted in the development of fenestrations in the endothelium of small venules and capillaries, even in regions where endothelial cells are not normally fenestrated, and was associated with increased vascular permeability (45). Interestingly, Dellian *et al.* (46) have described the quantification and long-term physiological characterization of microvessels induced by gels containing either VEGF or bFGF in transparent chambers in the dorsal skin or in the cranium of mice. These studies indicate that VEGF- or bFGF-induced vessels have similar diameter, permeability to albumin, and red cell velocities. However, permeability and red cell velocities were higher in the cranium than in the dorsal skin. These findings led to the conclusion that the steady-state physiological properties of blood vessels, including permeability, are primarily determined by the local microenvironment, rather than the initial angiogenic stimulus (46).

An additional effect of VEGF on the vascular endothelium is the stimulation of hexose transport (47). Exposure of bovine aortic endothelial cells to VEGF or TNF- α resulted in a

significant increase in the rate of hexose transport. The combination of factors had an additive effect. This action may have relevance for increased energy demands during endothelial cell proliferation or inflammation.

Recently, Melder *et al.* (48) have shown that VEGF promotes expression of VCAM-1 and ICAM-1 in endothelial cells. This induction may result in the adhesion of activated natural killer (NK) cells to endothelial cells, mediated by specific interaction of endothelial VCAM-1 and ICAM-1 with CD18 and VLA-4 on the surface of NK cells (48). It has been suggested that these effects may provide an explanation for the previously observed preferential adhesion of IL-2-activated NK cells to the tumor vasculature (49).

VEGF has been reported to have regulatory effects on certain blood cells. Clauss *et al.* (50) reported that VEGF may promote monocyte chemotaxis. More recently, Broxmeyer *et al.* (51) have shown that VEGF induces colony formation by mature subsets of granulocyte-macrophage progenitor cells that had been stimulated with a colony stimulating factor. These findings may be explained by the common origin of endothelial cells and hematopoietic cells and the presence of VEGF receptors in progenitor cells as early as hemangioblasts in blood islands in the yolk sac (see *Section VIII*). Furthermore, Gabrilovich *et al.* (52) have reported that VEGF may have an inhibitory effect on the maturation of host professional antigen-presenting cells such as dendritic cells. VEGF was found to inhibit immature dendritic cells, without having a significant effect on the function of mature cells. These findings led to the provocative hypothesis that VEGF may facilitate tumor growth also by allowing the tumor to avoid the induction of an immune response (52).

VEGF induces vasodilatation *in vitro* in a dose-dependent fashion (53) and produces transient tachycardia, hypotension, and a decrease in cardiac output when injected intravenously in conscious, instrumented rats (54). Such effects appear to be caused by a decrease in venous return, mediated primarily by endothelial cell-derived nitric oxide, as assessed by the requirement for an intact endothelium and the prevention of the effects by *N*-methyl-arginine (53, 54). Accordingly, VEGF has no direct effect on contractility or rate in isolated rat heart *in vitro* (54). These hemodynamic effects, however, are not unique to VEGF: other angiogenic factors such as aFGF and bFGF may also induce nitric oxide-mediated vasodilatation and hypotension (55).

III. Organization of the VEGF Gene

The human VEGF gene is organized in eight exons, separated by seven introns, and its coding region spans approximately 14 kb (56, 57). The human VEGF gene has been assigned to chromosome 6p21.3 (58). cDNA sequence analysis of a variety of human VEGF clones had indicated that VEGF may exist as one of four different molecular species, having, respectively, 121, 165, 189, and 206 amino acids (VEGF₁₂₁, VEGF₁₆₅, VEGF₁₈₉, VEGF₂₀₆) (26, 28, 56, 57). It is now well established that alternative exon splicing of a single VEGF gene is the basis for this molecular heterogeneity. VEGF₁₆₅ lacks the residues encoded by exon 6, while VEGF₁₂₁ lacks the residues encoded by exons 6 and 7. Compared with

VEGF₁₆₅, VEGF₁₂₁ lacks 44 amino acids; VEGF₁₈₉ has an insertion of 24 amino acids highly enriched in basic residues, and VEGF₂₀₆ has an additional insertion of 17 amino acids. Interestingly, there is no intron between the coding sequence of the 24-amino acid insertion in VEGF₁₈₉ and the additional 17-amino acid insertion found in VEGF₂₀₆. The 5'-end of the 51-bp insertion of VEGF₂₀₆ begins with GT, the consensus sequence for the 5'-splice donor necessary for mRNA processing. Therefore, the definition of the 5'-splice donor site for removal of a 1-kb intron sequence is variable (57). Analysis of the VEGF gene promoter region reveals a single major transcription start that lies near a cluster of potential Sp1 factor-binding sites. Also, several potential binding sites for the transcription factors AP-1 and AP-2 are present in the promoter region (56). VEGF₁₆₅ is the predominant molecular species produced by a variety of normal and transformed cells. Transcripts encoding VEGF₁₂₁ and VEGF₁₈₉ are detected in the majority of cells and tissues expressing the VEGF gene (56, 57). In contrast, VEGF₂₀₆ is a very rare form, so far identified only in a human fetal liver cDNA library (57). The organization of the murine VEGF gene has been also described (59). Similarly to the human gene, the coding region of the murine VEGF gene encompasses approximately 14 kb and is comprised of eight exons interrupted by seven introns. Analysis of exons suggests the generation of three isoforms, VEGF₁₂₀, VEGF₁₆₄ and VEGF₁₈₈. Therefore, murine VEGFs are shorter than human VEGF by one amino acid. A fourth isoform comparable to VEGF₂₀₆ is not predicted, since an in-frame stop codon is present in the region corresponding to the human VEGF₂₀₆ open reading frame. Analysis of the 3'-untranslated region of the rat VEGF mRNA has revealed the presence of four potential polyadenylation sites (60). A frequently used site is about 1.9 kb further downstream from the previously reported translation termination codon (30). The sequence within this 3'-untranslated region reveals a number of motifs that are known to be involved in the regulation of mRNA stability (60). (See also Section V. A.)

IV. Properties of the VEGF Isoforms

Native VEGF is a basic, heparin-binding, homodimeric glycoprotein of 45,000 daltons (61). These properties correspond to those of VEGF₁₆₅, the major isoform. VEGF₁₂₁ is a weakly acidic polypeptide that fails to bind to heparin (62). VEGF₁₈₉ and VEGF₂₀₆ are more basic and bind to heparin with greater affinity than VEGF₁₆₅ (62). Previous studies demonstrated that such differences in the isoelectric point and in affinity for heparin may profoundly affect the bioavailability of VEGF (62, 63). VEGF₁₂₁ is a freely soluble protein; VEGF₁₆₅ is also secreted although a significant fraction remains bound to the cell surface and the extracellular matrix. In contrast, VEGF₁₈₉ and VEGF₂₀₆ are almost completely sequestered in the extracellular matrix (63). However, these isoforms may be released in a soluble form by heparin or heparinase, suggesting that their binding site is represented by proteoglycans containing heparin-like moieties. Interestingly, the long forms may be released by plasmin (62, 63) after cleavage at the COOH terminus. This action generates a bioactive proteolytic

fragment having a molecular mass of ~34,000 daltons (62, 63). Plasminogen activation and generation of plasmin have been shown to play an important role in the angiogenesis cascade. Thus, proteolysis of VEGF is likely to occur also *in vivo*. Generation of bioactive VEGF by proteolytic cleavage may be especially important in the microenvironment of a tumor where increased expression of proteases, including PA, is well documented (64, 65). Keyt *et al.* (66) have shown that the bioactive product of plasmin action is comprised of the first 110 NH₂-terminal amino acids of VEGF. These findings suggest that the VEGF proteins may become available to endothelial cells by at least two different mechanisms: as freely diffusible proteins (VEGF₁₂₁, VEGF₁₆₅) or after protease activation and cleavage of the longer isoforms. However, loss of heparin binding, whether it is due to alternative splicing of RNA or plasmin cleavage, results in a substantial loss of mitogenic activity for vascular endothelial cells: compared with VEGF₁₆₅, VEGF₁₂₁ or VEGF₁₁₀ demonstrate 50-fold reduced potency when tested in endothelial cell growth assay; the VEGF_{165/110} heterodimer resulting from limited proteolysis of VEGF₁₆₅ demonstrated a 5–10-fold loss in potency when compared with wild type VEGF₁₆₅ (66). It has been suggested that the stability of VEGF-heparan sulfate-receptor complexes contributes to effective signal transduction and stimulation of endothelial cell proliferation (66). Thus, VEGF has the potential to express structural and functional heterogeneity to yield a graded and controlled biological response. Figure 1 illustrates some of the actions of the VEGF isoforms on the vascular endothelium and possible regulatory mechanisms.




FIG. 1. Schematic representation of the actions of VEGF isoforms on the vascular endothelium. Several stimuli may result in the release of the diffusible alternatively spliced VEGF isoforms (VEGF₁₆₅, VEGF₁₂₁) from a variety of cell types. These proteins may induce a complex series of effects on the vascular endothelium, including cell sprouting, induction of interstitial collagenase, plasminogen activators (PA), and plasminogen activator inhibitor I-1 (PAI-1), as well as extravasation of plasma proteins. Plasminogen activation results in generation of plasmin, which may cleave extracellular matrix-bound VEGF (VEGF₁₈₉ or VEGF₂₀₆) to release a diffusible proteolytic fragment (VEGF₁₁₀). Plasmin may also activate procollagenase. Activation of PAI-1 may constitute a negative regulatory step, by inhibiting the action of PA.

V. Regulation of VEGF Gene Expression

A. Hypoxia

Several mechanisms have been shown to participate in the regulation of VEGF gene expression. Among these, oxygen tension plays a major role, both *in vitro* and *in vivo*. VEGF mRNA expression is rapidly and reversibly induced by exposure to low pO₂ in a variety of normal and transformed cultured cell types (67–70). In glioblastoma multiforme and other tumors with significant component of necrosis, the VEGF mRNA is highly expressed in ischemic tumor cells that are juxtaposed to areas of necrosis (68), suggesting that local hypoxia is a major inducer of VEGF gene expression in the microenvironment of a tumor. Also, ischemia caused by occlusion of the left anterior descending coronary artery results in a dramatic increase in VEGF RNA levels in the pig myocardium, suggesting the possibility that VEGF may mediate the spontaneous revascularization that follows myocardial ischemia (71). Furthermore, Stone *et al.* (72) proposed that hypoxic up-regulation of VEGF mRNA in neuroglial cells, secondary to the onset of neuronal activity, plays an important physiological role in the development of the retinal vasculature.

Similarities exist between the mechanisms leading to hypoxic regulation of VEGF and erythropoietin (Epo) (73). Hypoxia inducibility is conferred on both genes by homologous sequences. By deletion and mutation analysis, a 28-base sequence has been identified in the 5'-promoter of the rat and human VEGF gene that mediated hypoxia-induced transcription in transient assays (60, 74). Such sequence reveals a high degree of homology and similar protein-binding characteristics as the hypoxia-inducible factor 1 (HIF-1) binding site within the Epo gene, which behaves like a classic transcriptional enhancer (75). HIF-1 has been purified and cloned as a mediator of transcriptional responses to hypoxia and is a basic, heterodimeric, helix-loop-helix protein (76, 77). Forsythe *et al.* (78) presented more direct evidence that HIF-1 is indeed implicated in the activation of the VEGF gene transcription during hypoxia. When reporter constructs containing the VEGF sequences that mediate hypoxia inducibility were cotransfected with expression vectors encoding HIF-1 subunits, reporter gene transcription was much greater than that observed in cells transfected with the reporter alone, both in hypoxic and normoxic conditions (78).

It has been shown that accumulation of adenosine, which occurs under hypoxic conditions, is involved in the induction of the VEGF gene during hypoxia (79). According to these studies, adenosine, by activating adenosine A₂ receptors, results in elevated cAMP concentrations that in turn increase VEGF mRNA levels, possibly through a protein kinase A-mediated pathway (79). Activation of *c-Src* also has been shown to participate in the hypoxic up-regulation of the VEGF gene (80). Hypoxia increases the kinase activity of pp60^{c-Src} and its phosphorylation on tyrosine 416. Expression of a negative dominant mutant of *c-Src* significantly reduced the hypoxic induction of VEGF (80).

It is noteworthy that several studies have shown that transcriptional activation is not the only mechanism leading to VEGF up-regulation in response to hypoxia. Increased

mRNA stability has been identified as an important post-transcriptional component (81–83). Sequences that mediate increased stability were identified in the 3'-untranslated region of the VEGF mRNA (see also *Section III*). Also, a hypoxia-induced protein that bound to such sequences was identified (81).

B. Cytokines

Several cytokines or growth factors up-regulate VEGF mRNA expression and/or induce release of VEGF protein. Exposure of quiescent human keratinocytes to serum, epidermal growth factor (EGF), TGF- β , or keratinocyte growth factor results in a marked induction of VEGF mRNA expression (84). Also, primary, nontransformed, keratinocytes show VEGF up-regulation in response to TGF- α (85, 86). EGF also stimulates VEGF release by cultured glioblastoma cells (87). In addition, treatment of quiescent cultures of several epithelial and fibroblastic cell lines with TGF- β resulted in induction of VEGF mRNA and release of VEGF protein in the medium (88). Based on these findings, it has been proposed that VEGF may function as a paracrine mediator for indirect-acting angiogenic agents such as TGF- β (88). Furthermore, IL-1 β induces VEGF expression in aortic smooth muscle cells (89). Both IL-1 α and PGE₂ have been shown to induce expression of VEGF in cultured synovial fibroblasts, suggesting the participation of such inductive mechanisms in inflammatory angiogenesis (90). IL-6 has been also shown to significantly induce VEGF expression in several cell lines (91). Not only promoter elements, but also motifs in the 5'-untranslated region of the VEGF mRNA were found to be involved in such up-regulation (91). IGF-I, a mitogen implicated in the growth of several malignancies, has also been shown to induce VEGF mRNA and protein in cultured colorectal carcinoma cells (92). The induction was mediated by a combined increase in transcriptional rate of the VEGF gene and in the stability of the mRNA. Thus, IGF-I, in addition to its direct mitogenic effects on malignant cells, may facilitate tumor growth via an increase in the vascular supply, mediated by VEGF.

C. Differentiation and transformation

Cell differentiation has been shown to play an important role in the regulation of VEGF gene expression (93). The VEGF mRNA is up-regulated during the conversion of 3T3 preadipocytes into adipocytes or during the myogenic differentiation of C2C12 cells. Conversely, VEGF gene expression is repressed during the differentiation of the pheochromocytoma cell line PC12 into nonmalignant, neuron-like cells. These studies also indicate that induction of VEGF mRNA expression in preadipocytes requires pathways mediated by both protein kinase C and protein kinase A activation (93). Consistent with the presence of AP-1 and AP-2 sites in the VEGF gene promoter, phorbol esters and forskolin, a potent activator of adenylate cyclase, induce VEGF mRNA expression (94). Accordingly, luteotrophic hormone, a known activator of adenylate cyclase, has been shown to induce expression of VEGF mRNA in cultured bovine ovarian granulosa cells (94).

Specific transforming events also result in induction of VEGF gene expression. A mutated form of the murine p53 tumor suppressor gene (Ala¹³⁵ > Val) has been shown to induce VEGF mRNA expression and potentiate phorbol ester-stimulated VEGF mRNA expression in NIH 3T3 cells in transient transfection assays (95). Likewise, oncogenic mutations or amplification of *ras* lead to VEGF up-regulation (96, 97). This effect is blocked by treatment with inhibitors of *ras* farnesyl transferase. Interestingly, expression of oncogenic *ras*, either constitutive or transient, potentiated the induction of VEGF by hypoxia (98). Also, overexpression of *v-raf* (97) or *v-Src* (99) lead to VEGF up-regulation. Moreover, the von Hippel-Lindau (VHL) tumor suppressor gene has been recently implicated in the regulation of VEGF gene expression (100). Human renal cell carcinoma cells either lacking endogenous wild type VHL gene or expressing an inactive mutant demonstrated altered regulation of VEGF gene expression, which was corrected by introduction of wild type VHL gene. Essentially all of the endothelial cells mitogenic activity released by tumor cells expressing mutant VHL gene was neutralized by anti-VEGF antibodies (100). These findings suggest that VEGF is a key mediator of the abnormal vascular proliferations and solid tumors characteristic of VHL syndrome (101). Most recently, Iliopoulos *et al.* (102) have shown that a function of the VHL protein is to provide a negative regulation of a series of hypoxia-inducible genes, including the VEGF, platelet-derived growth factor B chain, and the glucose transporter GLUT1 genes. In the presence of a mutant VHL, mRNAs for such genes were produced both under normoxic and hypoxic conditions. Reintroduction of wild type VHL cDNA resulted in inhibition of mRNA production under normoxic conditions and restored the characteristic hypoxia inducibility of those genes (102).

Taken together, these findings indicate that several, unrelated, alterations in cellular regulatory pathways result in VEGF up-regulation. Therefore, this event may be a final common pathway necessary for uncontrolled proliferation *in vivo*.

VI. The VEGF Receptors

A. Characterization and distribution of VEGF-binding sites

Two classes of high-affinity VEGF-binding sites were initially described in the surface of bovine endothelial cells, with dissociation constant (K_d) values of 10 pM and 100 pM, respectively, and molecular mass in the range of 180–220 kDa (103, 104). Lower affinity binding sites on mononuclear phagocytes were subsequently described (105). It has been suggested that such binding sites are involved in mediating chemotactic effects for monocytes by VEGF (50). Recently, it has been suggested that low-affinity, low molecular mass (120–130 kDa), receptors exist on endothelial and tumor cells (106, 107). Such receptors cross-link VEGF₁₆₅ but not VEGF₁₂₁. Thus, certain tumor and endothelial cells express lower affinity sites that bind selectively exon 7-encoded sequences. The molecular nature and biological significance of these receptors remain to be elucidated.

Ligand autoradiography studies on fetal and adult rat tissue sections demonstrated that high-affinity VEGF-bind-

ing sites are localized to the vascular endothelium of large or small vessels *in situ* (108, 109). These findings represented direct evidence for the hypothesis that the vascular endothelium is the major target of VEGF action. Interestingly, VEGF binding was apparent not only on proliferating but also on quiescent endothelial cells (108, 109). Also, the earliest developmental identification of high-affinity VEGF binding was in the hemangioblasts in the blood islands in the yolk sac, suggesting that expression of VEGF receptors is one of the earliest events in endothelial cell differentiation (109).

B. The Flt-1 and Flk-1/KDR tyrosine kinases

1. Binding characteristics. Two VEGF receptor tyrosine kinases (RTKs) have been identified (110–116). The Flt-1 (*fms*-like-tyrosine kinase) and KDR (kinase domain region) receptors bind VEGF with high affinity. Flk-1 (fetal liver kinase-1), the murine homolog of KDR, shares 85% sequence identity with human KDR (114). Both Flt-1 and KDR/Flk-1 have seven immunoglobulin (Ig)-like domains in the extracellular domain (ECD), a single transmembrane region and a consensus tyrosine kinase sequence that is interrupted by a kinase-insert domain (110–116). Figure 2 shows the alignment of the amino acid sequences of the ECD of Flt-1 and KDR. Flt-1 has the highest affinity for rhVEGF₁₆₅, with a K_d of approximately 10–20 pM (110). KDR has a somewhat lower affinity for VEGF: the K_d has been estimated to be approximately 75–125 pM (111).

A cDNA coding an alternatively spliced soluble form of Flt-1 (sFlt-1), lacking the seventh Ig-like domain, transmembrane sequence, and the cytoplasmic domain, has been identified in human umbilical vein endothelial cells (117, 118). This sFlt-1 receptor binds VEGF with high affinity (K_d 10–20 pM) and is able to inhibit VEGF-induced mitogenesis, suggesting that it may be a physiological negative regulator of VEGF action (117, 118).

An additional member of the family of RTKs with seven Ig-like domains in the ECD is Flt-4 (119–122) which, however, is not a receptor for VEGF but rather binds a newly identified ligand called VEGF-C or VEGF-related peptide (VRP) (see Section VII).

2. Signal transduction. Our understanding of the signal transduction properties of the VEGF receptors is still incomplete. VEGF has been shown to induce the phosphorylation of at least 11 proteins in bovine aortic endothelial cells (113). PLC- γ and two proteins that associate with PLC- γ were phosphorylated in response to VEGF (123). Furthermore, immunoblot analysis for mediators of signal transduction that contain SH2 domains demonstrated that VEGF induces phosphorylation of phosphatidylinositol 3-kinase, *ras* GTPase activating protein, and several others. These findings suggest that VEGF promotes the formation of multimeric aggregates of VEGF receptors with proteins that contain SH2 domains. These studies, however, did not identify which VEGF receptor(s) are involved in these events.

Several studies have indicated that Flt-1 and KDR have different signal transduction properties (124, 125). Porcine aortic endothelial cells lacking endogenous VEGF receptors display chemotaxis and mitogenesis in response to VEGF

FLT-1	1	MVS YWDTGVLLC-ALLSCLLLTGSS---SGSKLKD	PELSLKGTHIMQAGQTLHLQCRGEAAHKWSLP	EMV-----SKESERLSITKSACGRNGKQFCS
KDR	1	M-----ESKVLLAVALWLCVETRAASVGLPSVSLDL	PRLSIQKDILTIKANTTLQITCRGQRDLDLWLPNNQ	-----SGSEQRVEVTECSDG---LFCK
FLT-1	91	TLTLNTAQANHTGFYSCYLA VPTSKKKETESAIIYFI	ISDTGR	PFVEMYSEIPEIIHM--TEGRELVIPCRVTSPNITVTL-KKFPPLDTLIPDGKRIIWD
KDR	87	TLTIPKVIGNDTGAYKCFYR-----ETDLASVIYVY	QDYRS	PFIASVSDQHGVVYITENKNKT VVI PCLGSISNLNVSLCARYPEKRFVPDGNRISWD
FLT-1	188	SRKGFIIISNATYKEIGLLTCEATVNGHLYKTN-YLTHROT	NTIID	VQISTPRPVKLLRGHTLVLNCTATTPLNTRVQMTWSYPDEKNKRAS--VRRRIDQ
KDR	181	SKKGFTIPSYMISYAGMVCFEAKINDESYQSIMYIVVVV	GRIYD	DVLSPSHGIELSVGEKLVLNCTARTELVNIGIDFNWEYPPSSKHQHKLVNRDLKTQ
FLT-1	285	SNSHANIFYSVLTIDKMQNKDKGLYTCRVSRGSPFSKSVNTSVHIYDK	AFITV	KHRKQQVLETVAGKRSYRLSMKVKAFFSPFVWVKDGLPATEKSARYL
KDR	281	SGSEMKKFLSTLTIDGVTRSDQGLYTCAASSGLMTKKNSTFVRVHEK	PF	FAVFGSGMESLVEATVGER-VRIPAKYLGYPPEIKWYKNGIPLESN--HTI
FLT-1	385	TRGYSLIIKDVTEEDAGNYTILLSIKQSNVFNLTATLIVNVKP	QIY	EKAVSSFPDPALYPLGSRQILTCTAYGIPQP-TIKWFH---PCNNHSEARC
KDR	378	KAGHVLTIMEVSERDGTGNYTVILTNPISKEKQSHVSVLVVYVPP	Q	GEKSLIS--PVDSYQYGTQTCTCTVYAI PPPHHIHWYQLEECANEPQAVS
FLT-1	481	DPCSN-----NEESFILDADSNMGNRIESITQRMATIEGKNKMASTLVVADSRISGIYICIASNKVGTVGRNISFYIT	DVPN	GFHVNL--EKMPTGEDLK
KDR	476	VTNPY-----PCEEWRSVEDFQGGNKIEVNKNQFALIEGKNKT VSTLVIQAANVSALYKCEAVNKVGRGERVISFHV	TRGPE	---ITLQPDMPTEQESVS
FLT-1	575	LSCTVKNKFLYRDVTWILLR-----TVNNRTHMYSISKQ-----KMAIT-----KEHSITLNLTIMNVSLQDSGTYACRARNVYTGEELQKKEITIR	DQ	
KDR	569	LWCTADRSTFENLTWYKLGQPLPIHVGELPTPVCKNLDTL-WKLNATM--FSNSTNDILIMELKNASLQDQGDYVCLAQDRKTKRRCVVRQLTVL	ER	
FLT-1	659	EAPYLLRNLSHDHTVAISSSTLTDCHANGVPEPQITWFKNNHKIQQEPGIILGPGSSSTLFIERVTEEDGEGVYHCKATNQKGSVESSAYLTQVQGT	SDKSN--	
KDR	665	VAPTITGNLENQTTSIGESIEVSC TASGNPPQIMWFKDNETLVEDSGIVLKDGNRNLTIRVRKEDGLYTQCACSVLGCAKVEAFFIIEGA	Q	EKTNLD
FLT-1	757	-FE		
KDR	765	PFE		

FIG. 2. Alignment of the extracellular domains of human Flt-1 and KDR. The seven immunoglobulin (Ig)-like domains are shown as individual boxed areas.

when transfected with a plasmid coding for KDR (124). In contrast, transfected cells expressing Flt-1 lack such responses (124). Flk-1/KDR undergoes strong ligand-dependent tyrosine phosphorylation in intact cells, while Flt-1 reveals a weak or undetectable response (110, 124, 125). Also, VEGF stimulation results in weak tyrosine phosphorylation that does not generate any mitogenic signal in transfected NIH 3T3 cells expressing Flt-1 (125). These findings agree with other studies showing that placenta growth factor (PIGF), which binds with high affinity to Flt-1 but not to Flk-1/KDR, lacks direct mitogenic or permeability-enhancing properties or the ability to effectively stimulate tyrosine phosphorylation in endothelial cells (126) (see Section VII). Therefore, interaction with Flk-1/KDR is a critical requirement to induce the full spectrum of VEGF biological responses. In further support of this conclusion, VEGF mutants that bind selectively to Flk-1/KDR are fully active endothelial cell mitogens (see Section VI.B.5) (127). Furthermore, Kendall *et al.* (118) suggested that sFlt-1 may form heterodimeric complexes with KDR, which could potentially exert a dominant-negative effect on KDR signal transduction. These findings contributed to cast doubt on the role of Flt-1 as a truly signaling receptor. However, more recent evidence indicates that Flt-1 indeed signals, although our understanding of these processes is clearly fragmentary. Cunningham *et al.* (128), using the yeast two-hybrid system, have demonstrated an interaction between Flt-1 and the p85 subunit of phosphatidylinositol 3-kinase. Mutagenesis analysis revealed that change of a tyrosine residue at position 1213 to phenylalanine completely abolished such interaction. These data suggest that p85 couples Flt-1 to intracellular signal transduction systems and implicate elevated levels of PtdIns(3,4,5)P₃ levels in this process (128). Also, members of the *Src* family, such as *Fyn* and *Yes*, show an increased level of phosphorylation after VEGF stimulation in transfected cells expressing Flt-1

but not KDR (124). Furthermore, Barleon *et al.* (129) have shown that a specific biological response, the migration of monocytes in response to VEGF (or PlGF), is mediated by Flt-1. However, the most compelling evidence so far for an important biological role played by the Flt-1 receptor has been provided by gene knockout studies (see Section VIII. B).

3. Regulation. The expression of Flt-1 and Flk-1/KDR genes is largely restricted to the vascular endothelium (see Section VIII.A). The promoter region of Flt-1 has been cloned and characterized and a 1-kb fragment of the 5'-flanking region essential for endothelial-specific expression was identified (130). Likewise, a 4-kb 5'-flanking sequence has been identified in the promoter of KDR that confers endothelial cell-specific activation (131).

Similarly to VEGF, hypoxia has been proposed to play an important role in the regulation of VEGF receptor gene expression. Exposure of rats to acute or chronic hypoxia led to pronounced up-regulation of both Flt-1 and Flk-1/KDR genes in the lung vasculature (132). Also, Flk-1/KDR and Flt-1 mRNAs were substantially up-regulated throughout the heart after myocardial infarction in the rat (133). However, *in vitro* studies have yielded unexpected results. Even though Thieme *et al.* (134) have shown that hypoxia increases VEGF receptor number by 50% in cultured bovine retinal capillary endothelial cells, the expression of KDR is not induced but paradoxically shows an initial down-regulation (135). Brogi *et al.* (136) have proposed that the hypoxic up-regulation of KDR observed *in vivo* is not direct but requires the release of an unidentified paracrine mediator from ischemic tissues. Also, recent studies have shown that both TNF- α (137) and TGF- β (138) are able to inhibit the expression of the KDR gene in cultured endothelial cells.

4. Structural requirements for ligand binding in Flt-1 and KDR. As noted above, the VEGF receptors have seven Ig-like do-

mainly in the ECD. Until now, the significance and function of these domains for ligand binding and receptor activation were unknown. Recently, the domains in the ECD of Flt-1 and KDR responsible for specific ligand recognition were identified by constructing and analyzing a variety of receptor variants (139). These included individual Ig-like domain (140) deletions, as well as chimeras in which domains of either KDR or Flt-4 were exchanged for the homologous sequences from Flt-1. Deletion of the second Ig-like domain of human Flt-1 completely abolishes VEGF binding (Fig. 3). Introduction of the second domain of KDR into an Flt-1 mutant lacking the homologous domain restored VEGF binding. However, PlGF was unable to displace VEGF bound to such mutant, a pattern characteristic of the KDR but not the Flt-1 receptor (Fig. 3). Also, "swap" experiments in which the second Ig-like domain of Flt-1 replaced the corresponding domain in Flt-4 demonstrated that such a chimeric receptor had the ability to bind VEGF with affinity nearly identical to that of wild type Flt-1. Furthermore, transfected cells expressing this chimeric Flt-4 receptor exhibited in-

creased DNA synthesis in response to both VEGF and PlGF (139). Thus, VEGF binding to domain 2 of Flt-1 is able to initiate a signal transduction cascade, even in the context of the ECD of a foreign receptor. Further studies are required to elucidate the significance of the remaining Ig-like domains in receptor dimerization (141) and in coupling binding with signal transduction.

5. VEGF determinants for binding Flt-1 and KDR. Site-directed mutagenesis has been used to localize the determinants on VEGF that mediate binding to the KDR and Flt-1 receptors. Alanine-scanning analysis was performed to identify a positively charged surface in VEGF that mediates receptor binding (127, 142). A model based on the crystal structure of PDGF-BB was used (143). Arg⁸², Lys⁸⁴, and His⁸⁶, located in a hairpin loop, were found to be critical for binding KDR, while negatively charged residues, Asp⁶³, Glu⁶⁴, and Glu⁶⁷, were primarily responsible for Flt-1 binding. The single mutations R82A, K84A, and H86A were found to display modestly decreased KDR binding. The triple mutants involving alanine replacement or neo-glycosylation sites, R82A, K84A, H86A VEGF and R82N, I83L, K84S VEGF, exhibited minimal binding to KDR receptor. The half-maximally effective concentrations (EC_{50}) to stimulate bovine capillary endothelial cell growth for most of the VEGF mutants were similar to those observed for wild type VEGF. The most significant effect on endothelial cell proliferation was observed with mutations in the 82–86 region. The EC_{50} of R82A, K84A, H86A VEGF increased 20-fold such that mitogenic potency of this mutant was decreased to 5% of wild type VEGF. In contrast, the mutants that failed to bind Flt-1 were fully active endothelial cell mitogens (127, 142).

VII. VEGF-Related Molecules

Over the last few years, three VEGF-related genes have been identified from mammalian sources. The encoded factors are known as PlGF, VEGF-B, and VEGF-C/VRP. In addition, two sequences in the genome of the parapoxvirus *orf* virus show homology to VEGF. Figure 4 shows the alignment of the amino acid sequences of these molecules with the sequence of VEGF₁₆₅. Although the biological role of these factors is still largely unclear, their structural homology to VEGF suggests that they may play a role in the regulation of blood vessel growth. The first VEGF-related factor identified is PlGF. This molecule shares a 53% identity with the PDGF-like region of VEGF. The encoded protein was expected to have 149 amino acids, including the signal peptide (144). Subsequently, a longer form characterized by a 21-amino acid insertion was identified (145). Similar to the 24-amino acid insertion in the longer forms of VEGF, this insertion is highly enriched in basic residues. These two isoforms, which arise from alternative splicing of mRNA, are known as PlGF-1 and PlGF-2 or PlGF₁₃₁ and PlGF₁₅₂, respectively. Similar to VEGF, these molecules are dimeric glycoproteins. Park *et al.* (126) have shown that PlGF binds with high affinity ($K_d \sim 250$ pM) Flt-1 but not KDR. Purified PlGF demonstrated minimal activity in vascular endothelial cell growth and vascular permeability assays, suggesting that binding to KDR is a requirement for both activities. However, PlGF was able to




FIG. 3. The second Ig-like domain contains the major determinants for binding and ligand specificity in the VEGF receptors. In panel A, Flt-1-IgG individual domain deletion variants (5 ng per reaction) were tested for their ability to bind [¹²⁵I]VEGF₁₆₅ in the absence (striped bars) or presence of 50 ng cold VEGF₁₆₅ (solid bars). Deletion of the second Ig-like domain completely abolishes the binding of VEGF. The second Ig-like domain of KDR was cloned into the Flt-1 domain 2 deletion construct to produce "swap" mutants (panel B). Replacement of the second domain of Flt-1 with the homologous domain of KDR reestablished VEGF-binding. However, PlGF₁₅₂ (open bars) could not displace VEGF₁₆₅ bound to Flt.K2, a pattern characteristic of the KDR but not the Flt-1 receptor.

VEGF ₁₆₅	1	MNFL-----LSVHWSLALLLYLH-H-----AKWSQAAPMAE--
VEGF-B	1	MSPL-----LRRLL--LAALLQLA-P-----AQAPVSQPDAP--
VEGF-C/VRP	1	MHLGGFSVACSLAAALLPGPREAPAAAAAFESGLDLSDAEPAGEATAYASKDLEEQLRSVSSVDELMTVLYPEYWKMYKCQLRKGGWQHNRQANLN
PlGF ₁₅₂	1	M-PV-----MRLFPCLQLLAGLALP-----AVPPQQWALSA--
OrfA2R	1	M-KL-----TATLQVVVALLICMY-----NLPECVQSQN--
VEGF ₁₆₅	32	-----GGGQNHHEVVKFMD-VYQRSYCHPIETLVDIFQEYPDEIEYIFKPSVPLMFCGGCCNDEGLECVPTTESNITMQIMRIKP--HQQQHIGE
VEGF-B	30	-----G--H-QRKVVSWID-VYTRATCQPREVVVPLTVELMGTVAQQLVPSCVTVQRCGGCCDDGLECVPTGQHQVRMQLMIR--YPSSQLGE
VEGF-C/VRP	101	SRTEETIKFAAAHYNTIELKSIDNEWKRTOCMPEVCIDVGKEFGVATNTFFKPPCVSVYRCGGCCNSEGLQCMNTSTSYLSKTLFEITVPLSQGPKPVT
PlGF ₁₅₂	32	-----GNGSSEVEVVPFQE-VWGRSYCRALERLVDVVSEYPSVEEHMFSPSCVSLRLCTGCCGDENLHCVPVETANVTMQLLKIRS--GDRPSYVE
OrfA2R	29	-----DSPPSTNDWNR-TLDKSGCKPRDTVVYLGEYPESTNLQYNPRCTVTKRCSCCNGDGGQICITAVETRNTTVTV-----SVTG
VEGF ₁₆₅	120	MSFLQHNNKCECR-----PKKD-----RA--R
VEGF-B	114	MSLEEHSQCCECR-----PKKK-----DSAVK
VEGF-C/VRP	201	ISFANHTSCROMSKLDVYRQVHSIIIRSLPATLPQCQAANKTCPTNYMWNHICRCLAQEDFMFSSDAGDDSTDGFHDICGPNKELDEETCQCVCRAGLR
PlGF ₁₅₂	120	LTFSQHVRCECR-----PLRE-----KM--K
OrfA2R	105	VSSSSGTNSGV-----
VEGF ₁₆₅	139	QENPCG-----PCSERRKHLFVQDPQTCKCSC-----KNTD-SR-----CKAR-----QLE
VEGF-B	135	PDSRP-----LCPRCTQHHQRDPDPTCRRC-----RRRSFLR-----CQGR-----GLE
VEGF-C/VRP	301	PASCGPHKELDRNSCQCCKNKLFPSCQGANREFDENTCQCVCKRTCPRNQPLNPGKACACECTESPQKCLLKGGKFHHQTCSCYRRPCTNRQKACEPGFS
PlGF ₁₅₂	139	PERRRP-----K-GRGRRREKQRPDTC-----
OrfA2R	116	-----STNLQRISVTEHTKDCIG-----
VEGF ₁₆₅	179	LNERTRCCKP--RR----
VEGF-B	176	LNPDTRCRKL--RR----
VEGF-C/VRP	401	YSEEVRCVPSYWKRPQMS
PlGF ₁₅₂	161	---HLGGDAVP--RR----
OrfA2R	135	-RTTTTPTTTTREP RR----

FIG. 4. Amino acid sequence of VEGF₁₆₅ and VEGF-related molecules: VEGF-B, VEGF-C, PlGF₁₅₂, and a VEGF-like sequence identified in the genome of the parapoxvirus orf virus. The conserved cysteine residues are boxed.

potentiate the bioactivity of low, marginally efficacious, concentrations of VEGF, both on endothelial cell growth and on vascular permeability (126). The molecular basis of this effect remains to be fully elucidated. Interestingly, naturally occurring heterodimers between VEGF and PlGF have been identified in the conditioned medium of a rat glioma cell line (146). In agreement with previous studies, the PlGF homodimer demonstrated minimal mitogenic activity on endothelial cells. However, the VEGF:PlGF heterodimer was active, although its potency was approximately 7-fold lower than the VEGF homodimer. It has been suggested that the formation of heterodimers with PlGF constitutes a mechanism of negative regulation of VEGF bioactivity, by shifting the balance toward less potent molecules (147).

As previously noted, similarly to the VEGF receptors, Flt-4 is a RTK with seven Ig-like domains in the ECD (119–122). Interestingly, the expression of Flt-4 mRNA, which is initially localized to angioblasts and venules in the early embryo, becomes restricted to lymphatic endothelium at later stages of development (148). This expression pattern suggested that Flt-4 may play a role in the regulation of lymphangiogenesis (122, 148). A ligand selective for Flt-4 has been recently identified by two groups and has been named VEGF-C (149) or VRP (150). VEGF-C/VRP is a secreted protein with 399 amino acid residues and has a 32% identity to VEGF. Its COOH-terminal half contains a 180-amino acid region that is not found in VEGF. This region contains cysteine-rich motifs similar to a protein component of silk produced by the larval salivary gland of the midge *C. tentans* (149, 150). VEGF-C/

VRP has been reported to stimulate the growth of human lung endothelial cells, albeit at 100-fold less potency than VEGF₁₆₅ (150). It is still unclear whether VEGF-C/VRP is capable of high-affinity interaction with KDR (149, 150).

A newly identified member of the VEGF gene family is VEGF-B (151, 152). This molecule consists of 188 amino acids, including the signal peptide. VEGF-B has been reported to stimulate the growth of human and bovine vascular endothelial cells (151). Interestingly, VEGF-B is distributed primarily in the skeletal muscle and myocardium and is coexpressed with VEGF (151). Similar to the long forms of VEGF, VEGF-B is expressed as a membrane-bound protein that can be released in a soluble form after addition of heparin. VEGF-B and VEGF are also able to form heterodimers, when coexpressed (151). These findings led to the hypothesis that VEGF-B may participate in the regulation of angiogenesis, particularly in muscle (151). Figure 5 schematizes the interaction of VEGF and VEGF-related factors with their tyrosine kinase receptors.

Intriguingly, two sequences having a significant homology to VEGF have been identified in the genome of two different strains of orf virus, a parapoxvirus that affects goats, sheeps, and occasionally humans (153). This suggests that the viral VEGF-like gene has been acquired from a mammalian host and is undergoing genetic drift. Interestingly, the lesions of goats and humans after orf virus infection are characterized by extensive microvascular proliferation in the skin, raising the possibility that the product of the viral VEGF-like gene is responsible for such lesions.




FIG. 5. The diagram illustrates the interaction of VEGF and VEGF-related molecules with the three known members of the family of RTKs with seven Ig-like domains in the ECD. VEGF interacts with Flt-1 and KDR; PIGF binds only Flt-1 and VEGF-C/VRP binds with high affinity to Flt-4. It is unknown at the present time whether VEGF-B binds to any of these receptors.

VIII. Role of VEGF and Its Receptors in Physiological Angiogenesis

A. Distribution of VEGF, Flk-1/KDR, and Flt-1 mRNA

The proliferation of blood vessels is crucial for a wide variety of physiological processes such as embryonic development, normal growth and differentiation, wound healing, and reproductive functions. Previous studies have indicated that the VEGF mRNA is temporally and spatially related to the proliferation of blood vessels in the rat, mouse, and primate ovary and in the rat uterus, suggesting that VEGF is a mediator of the cyclical growth of blood vessels that occurs in the female reproductive tract (154–157). In fact, *in situ* hybridization studies in the rat ovary provided the first evidence that VEGF may be a regulator of physiological angiogenesis (154).

During embryonic development, VEGF expression is first detected within the first few days after implantation in the giant cells of the trophoblast (109, 158), suggesting a role for this factor in the induction of vascular growth in the decidua, placenta, and vascular membranes. At later developmental stages in the mouse or rat embryos, the VEGF mRNA is expressed in several organs, including heart, vertebral column, kidney, and along the surface of the spinal cord and brain (109, 158). In the developing mouse brain, the highest levels of mRNA expression are associated with the choroid plexus and the ventricular epithelium (158). In the human fetus (16–22 weeks), VEGF mRNA expression is detectable in virtually all tissues and is most abundant in lung, kidney, and spleen (159). VEGF protein, as assessed by immunocytochemistry, is expressed in epithelial cells and myocytes, but not vascular endothelial cells (159).

In situ hybridization studies have shown that the Flk-1 mRNA is expressed in the yolk sac and intraembryonic mesoderm and later on in angioblasts, endocardium, and small

and large vessel endothelium (115, 116). There is evidence that the Flk-1 mRNA is down-regulated in adult endothelial cells as compared with fetal endothelial cells (115, 116). These findings strongly suggested a role for Flk-1 in the regulation of vasculogenesis and angiogenesis. Other studies have demonstrated that expression of Flk-1 mRNA is first detected in the proximal-lateral embryonic mesoderm, which gives rise to the heart (160). Flk-1 is then detectable in endocardial cells of heart primordia and subsequently in the major embryonic and extraembryonic vessels (160). These studies have indicated that Flk-1 may be the earliest marker of endothelial cell precursors (160). The Flt-1 mRNA is selectively expressed in vascular endothelial cells, both in fetal and adult mouse tissues (161). Similar to the high-affinity VEGF binding (108, 109), the Flt-1 mRNA is expressed in both proliferating and quiescent endothelial cells (161), suggesting a role for Flt-1 in the maintenance of endothelial cells.

Interestingly, VEGF expression is also detectable around microvessels in areas where endothelial cells are normally quiescent, such as kidney glomerulus, pituitary, heart, lung, and brain (61, 162, 163). These findings raised the possibility that VEGF may be required not only to induce active vascular proliferation but, at least in some circumstances, also for the maintenance of the differentiated state of blood vessels (61). In agreement with this hypothesis, Alon *et al.* (164) have shown that VEGF acts as a survival factor, at least for the developing retinal vessels. They propose that hyperoxia-induced vascular regression in the retina of neonatal animals is a consequence of inhibition of VEGF production by glial cells. Accordingly, intraocular administration of VEGF to newborn rats at the onset of hyperoxia was able to prevent cell apoptosis and regression of the retinal vasculature (164).

It has been suggested that VEGF is also involved in a major pathophysiological process such as wound healing (84–86). Keratinocytes in a healing wound express VEGF mRNA.

Interestingly, a decreased expression of VEGF mRNA has been observed in the skin of genetically diabetic db/db mice (84), suggesting that an altered regulation of VEGF gene expression contributes to defective angiogenesis and impaired wound healing characteristic of this disorder.

B. Analysis of *Flk-1/KDR*, *Flt-1*, and *VEGF* gene knockouts

Recent studies have demonstrated that both *Flt-1* and *Flk-1/KDR* are essential for normal development of embryonic vasculature. However, their respective roles in endothelial cell proliferation and differentiation appear to be distinct (165, 166). Mouse embryos homozygous for a targeted mutation in the *Flt-1* locus died *in utero* between day 8.5 and 9.5 (165). Endothelial cells developed in both embryonic and extraembryonic sites but failed to organize in normal vascular channels. Mice in which the *Flk-1* gene had been inactivated lacked vasculogenesis and also failed to develop blood islands (166). Hematopoietic precursors were severely disrupted and organized blood vessels failed to develop throughout the embryo or the yolk sac, resulting in death *in utero* between day 8.5 and 9.5 (166).

However, these findings do not necessarily imply VEGF as being equally essential, since other ligands might potentially activate the *Flt-1* and *Flk-1/KDR* receptors and thus substi-

tute VEGF action. Very recent studies (16, 17) have generated direct evidence for the role played by VEGF in embryonic vasculogenesis and angiogenesis. Unexpectedly, inactivation of the VEGF gene in mice resulted in embryonic lethality in heterozygous embryos, between day 11 and 12. The *VEGF*^{+/−} embryos were growth retarded and also exhibited a number of developmental anomalies (167). The forebrain region appeared significantly underdeveloped. In the heart region, the outflow region was grossly malformed; the dorsal aortas were rudimentary, and the thickness of the ventricular wall was markedly decreased. The yolk sac revealed a substantially reduced number of nucleated red blood cells within the blood islands. Also, the vitelline veins failed to fuse with the vascular plexus of the yolk sac. Significant defects in the vasculature of other tissues and organs, including placenta and nervous system, were evidenced. For example, in the nervous system of heterozygous embryos at day 10.5, vascular elements could be demonstrated in the mesenchyme but not in the neuroepithelium (17) (Fig. 6). This failure of blood vessel ingrowth was accompanied by apoptosis and disorganization of neuroepithelial cells (Fig. 6). The *VEGF*^{+/−} embryos survive approximately 2 days longer than the *Flt-1* or *Flk-1/KDR* null embryos, presumably reflecting a partial activation of these tyrosine kinases by VEGF. *In situ* hybridization confirmed expression of




FIG. 6. Hematoxylin and eosin staining (upper panels) and CD34 immunostaining (lower panels) on sections of neuroepithelium (ne) from wild type (left) and *VEGF*^{+/−} (right) E 10.5 mouse embryos. Arrows indicate blood vessels. Blood vessel lumina can be identified in the mesenchyme adjacent to the ne in both groups. However, they are absent within the ne of the heterozygous embryos. Note also the presence of apoptotic cells in the ne of the heterozygous embryos. This contrasts with the well differentiated and vascularized ne in the wild type. [Reproduced with permission from N. Ferrara *et al.*: *Nature* 380:439–442, 1996 (17). ©1996 Macmillan Magazines Limited.]

VEGF mRNA in heterozygous embryos. Thus, the VEGF+/- phenotype is due to gene dosage and not to maternal imprinting.

Although several heterozygous phenotypes have been described (168), this may be the first report that the loss of a single allele of a gene that is not maternally imprinted can be lethal. Therefore, VEGF and its receptors are essential for blood island formation and angiogenesis such that even reduced concentrations of VEGF are inadequate to support a normal pattern of development. These findings also indicate that, in the VEGF+/- mutant, the mechanisms that normally up-regulate VEGF gene expression, such as hypoxia, are unable to provide an effective compensatory response. It is tempting to speculate that, as VEGF concentrations and angiogenic gradients fall below a threshold during critical periods, this can cause irreversible disruption of normal organogenesis.

IX. Role of VEGF in Pathological Angiogenesis

A. Tumor angiogenesis

1. Expression of VEGF in human tumors. *In situ* hybridization studies have demonstrated that the VEGF mRNA is markedly up-regulated in the vast majority of human tumors so far examined. These include: lung (169), thyroid (170), breast (171, 172), gastrointestinal tract (173, 174), kidney and bladder (175), ovary (176), and uterine cervix (177) carcinomas, angiosarcoma (178), germ cell tumors (179), and several intracranial tumors including glioblastoma multiforme (68, 180, 181) and sporadic, as well as VHL syndrome-associated, capillary hemangioblastoma (182, 183) (Table 1). Only sections of lobular carcinoma of the breast and papillary carcinoma of the bladder failed to show significant VEGF mRNA expression (184). As already indicated in Section V.B, the expression of VEGF in glioblastoma multiforme and other tumors with significant necrosis is highest in hypoxic tumor cells adjacent to necrotic areas (68, 180, 181). A correlation has been noted

between VEGF mRNA expression and vascularity of the tumor (169, 174, 177, 182, 183). In the tumors where VEGF and PlGF were coexpressed, only VEGF expression correlated with the degree of malignancy and vascularity (170, 179). In virtually all specimens examined, the VEGF mRNA was expressed in tumor cells but not in endothelial cells. In contrast, the mRNAs for Flt-1 and KDR were up-regulated in the endothelial cells associated with the tumor (173, 180, 185). These findings are consistent with the hypothesis that VEGF is primarily a paracrine mediator (186). An interesting exception may be angiosarcoma, where VEGF and Flt-1 mRNA were found to be coexpressed in angiosarcoma cells, raising the possibility that in this malignancy VEGF may play a role as an autocrine factor (178). Angiosarcoma cells, however, arise from the endothelium. Recently, Freeman *et al.* (187) have suggested that lymphocytes infiltrating the tumor may constitute an additional source of VEGF, which contributes to tumor angiogenesis. Immunohistochemical studies have localized the VEGF protein not only to the tumor cells but also to the vasculature (173, 180, 185). This finding indicates that tumor-secreted VEGF accumulates in the target cells. Ultrastructural studies have localized VEGF bound to tumor endothelial cells to the abluminal plasma membrane and to the recently described vesiculovacuolar organelles, cytoplasmic structures that are thought to be involved in macromolecular transport across the tumor endothelium (188).

Elevations in VEGF levels have been detected in the serum of some cancer patients (189). Also, a correlation has been observed between VEGF expression and microvessel density in primary breast cancer sections (190). A postoperative survey indicated that the relapse-free survival rate of patients with VEGF-rich tumors was significantly worse than that of VEGF-poor tumors, suggesting that expression of VEGF is associated with stimulation of angiogenesis and with early relapse in primary breast cancer (190). A similar correlation has been described in gastric carcinoma patients (191). VEGF positivity in tumor sections was correlated with vessel involvement, lymph node metastasis, and liver metastasis. Furthermore, patients with VEGF-positive tumors had a worse prognosis than those with VEGF-negative tumors (191).

2. Inhibition of VEGF action *in vivo*. The availability of specific monoclonal antibodies capable of inhibiting VEGF-induced angiogenesis *in vivo* and *in vitro* (192) made it possible to generate direct evidence for a role of VEGF in tumorigenesis. In a study published by Kim *et al.* in 1993 (193), such antibodies were found to exert a potent inhibitory effect on the growth of three human tumor cell lines injected subcutaneously in nude mice, the SK-LMS-1 leiomyosarcoma, the G55 glioblastoma multiforme, and the A673 rhabdomyosarcoma. The growth inhibition ranged between 70% and more than 95%. Figure 7 illustrates the effects of the anti-VEGF-neutralizing antibody on the *in vivo* growth of such cell lines. These findings provided the first direct demonstration that inhibition of the action of an endogenous endothelial cell mitogen may result in suppression of tumor growth *in vivo*. Subsequently, other tumor cell lines were found to be inhibited *in vivo* by this treatment (194–198) (Table 2).

In agreement with the hypothesis that inhibition of neo-

TABLE 1. Human tumors overexpressing the VEGF mRNA *in situ*

Tumor type	Reference
Intracranial tumors	
Glioblastoma multiforme	68, 180–182
Meningioma	182
Capillary hemangioblastoma	182, 183
Thyroid carcinomas	170
Lung carcinomas	169
Breast carcinomas	171, 172
Gastrointestinal tract tumors	
Esophageal carcinomas	173
Gastric carcinomas	173, 191
Small bowel carcinomas	173
Colorectal carcinomas	173, 194
Hepatocellular carcinoma	174
Urinary tract tumors	
Kidney carcinomas	175
Bladder carcinomas	175
Female reproductive tract tumors	
Ovarian carcinomas	176
Uterine cervix carcinomas	177
Germ cell tumors	179
Angiosarcoma	178




FIG. 7. Effects of anti-VEGF monoclonal antibody on tumor size (A, B) and weight (C). A673, G55, and SK-LMS-1 cells were injected subcutaneously in nude mice. Animals were then treated with anti-VEGF neutralizing antibody (A4.6.1) or a control antibody (5B6) twice weekly intraperitoneally, at the indicated doses. In A and B, tumor size was measured weekly. Panel C illustrates the weight of the tumors at the end of the experiment. Data shown reflect the response to 100 µg (5 mg/kg) of antibody twice weekly. Plus and minus signs denote the presence or absence of antibody treatment. A673 and G55 tumors were collected 4 weeks after tumor cell injection. SK-LMS-1 tumors were harvested after 10 weeks. [Reproduced with permission from J. Kim *et al.*: *Nature* 362:841–844, 1993 (193). © 1993 Macmillan Magazines Limited.]

vascularization is the mechanism of tumor suppression, the density of blood vessels was significantly lower in sections of tumors from antibody-treated animals as compared with controls (193, 194). Furthermore, neither the antibodies nor VEGF had any effect on the *in vitro* growth of the tumor cells (193, 194). Intravital videomicroscopy techniques have allowed a more direct verification of the hypothesis that anti-VEGF antibodies indeed block tumor angiogenesis (195). Tumor spheroids of A673 cells were implanted in dorsal skinfold chambers inserted in nude mice. Noninvasive im-

TABLE 2. Human tumor cell lines inhibited *in vivo* by anti-VEGF neutralizing antibodies

Cell line	Tumor type	Reference
A673	Rhabdomyosarcoma	193, 195
G55	Glioblastoma multiforme	193
SK-LMS-1	Leiomyosarcoma	193
HM 7	Colon carcinoma	194
LS LiM6	Colon carcinoma	194
A-431	Epidermoid carcinoma	196
HT-1080	Fibrosarcoma	197
U87	Glioblastoma multiforme	198
LS174T	Colon carcinoma	198
P-MEL	Melanoma	198

The following data represent additional human tumor cell lines inhibited *in vivo* by anti-VEGF neutralizing antibodies: MCF-7 cells from breast carcinoma (P. Borgström *et al.*, submitted); D-145 cells from prostatic carcinoma (P. Borgström *et al.*, submitted); SK-OV-3 cells from ovarian carcinoma (S. Mesiano, unpublished observations); SK-AS cells from neuroblastoma (N. Ferrara, unpublished observations); and Calu-6 cells from non-small cell lung carcinoma (F. Kabbinavar, unpublished observations).

aging of the vasculature revealed a nearly complete suppression of tumor angiogenesis in anti-VEGF treated animals as compared with controls, at all time points examined. These findings were corroborated by histological analysis that showed a dramatic difference in the density of CD34-positive vascular elements between the two groups (195). Thus, inhibition of VEGF-induced angiogenesis caused a dramatic change in growth characteristics of this cell line, from a rapidly growing malignancy to a dormant tumor seedling. Very similar findings were subsequently obtained with other tumor cell lines (P. Borgström *et al.*, submitted).

Warren *et al.* (194) have demonstrated that VEGF is a mediator of the *in vivo* growth of human colon carcinoma HM7 cells in an orthotopic nude mouse model of liver metastasis. Similar to human tumors, in this murine model the expression of Flk-1 mRNA was markedly up-regulated in the vasculature associated with liver metastases. Treatment with anti-VEGF monoclonal antibodies resulted in a dramatic decrease in the number and size of metastases. Most of the tumors in the treated group were less than 1 mm in diameter and all were less than 3 mm. Also, neither blood vessels nor Flk-1 mRNA expression could be demonstrated in such metastases. Also, administration of anti-VEGF-neutralizing antibodies inhibited primary tumor growth and metastasis of A431 human epidermoid carcinoma cells in *scid* mice (196) or HT-1080 fibrosarcoma cells implanted in BALB/c nude mice (197).

Recently, Borgström *et al.* (submitted) have shown that a combination treatment that includes anti-VEGF monoclonal antibody and doxorubicin results in a significant enhancement of the efficacy of either agent alone and led in some cases to complete regression of tumors derived from MCF-7 breast carcinoma cells in nude mice. Combination treatments that include anti-VEGF monoclonal antibody and cisplatin have resulted in similar enhancement of the efficacy of each agent (our unpublished observations).

Intravital fluorescence microscopy and video imaging analysis have been also applied to address the important

issue of the effects of VEGF on permeability and other properties of tumor vessels (198). Three different human tumor cell lines (U87, P-MEL, and LS174T) were implanted in two locations in immunodeficient mice, the cranium and the dorsal skinfold. Treatment with an anti-VEGF monoclonal antibody (192) was initiated when the tumor xenografts were already established and vascularized and resulted in time-dependent reductions in vascular permeability (198). These effects were accompanied by striking changes in the morphology of vessels, with dramatic reduction in diameter and tortuosity (198). This reduction in diameter is expected to block the passage of blood elements and eventually stop the flow in the tumor vascular network. Accordingly, a regression of blood vessels was observed after repeated administrations of anti-VEGF antibody. These findings led to the intriguing conclusion that tumor vessels require constant stimulation with VEGF to maintain not only their proliferative properties but also some key morphological features (198).

An additional verification of the hypothesis that VEGF action is required for tumor angiogenesis has been provided by the finding that retrovirus-mediated expression of a dominant negative Flk-1 mutant, which inhibits signal transduction through wild type Flk-1/KDR receptor, suppresses the growth of glioblastoma multiforme as well as other tumor cell lines *in vivo* (199, 200).

Further evidence that VEGF action is necessary for effective tumor angiogenesis has been recently obtained in an *in vivo* model of embryonic stem (ES) cell tumorigenesis (17). ES cells are able to form highly vascularized teratocarcinomas when injected in nude or syngeneic mice (201). VEGF null ES cells were dramatically impaired in their ability to form tumors in nude mice. The number of vessels in the VEGF^{-/-} group was substantially reduced and showed a much less complex branching pattern than that observed in controls. Thus, even in a pluripotent system such as the ES cells, which is expected to have the potential to activate redundant pathways, VEGF is required for *in vivo* growth.

B. Intraocular neovascular syndromes

Diabetes mellitus, occlusion of central retinal vein, or prematurity with subsequent exposure to oxygen can all be associated with intraocular neovascularization (202, 203). The new blood vessels may lead to vitreous hemorrhage, retinal detachment, neovascular glaucoma, and eventual blindness (5). Diabetic retinopathy is the leading cause of blindness in the working population (203). All of these conditions are known to be associated with retinal ischemia (202). In 1948, Michaelson proposed that a key event in the pathogenesis of these conditions is the release by the ischemic retina into the vitreous of a diffusible angiogenic factor(s) ("factor X") responsible for retinal and iris neovascularization (204). Factors such as FGF and IGF-I do not show a consistent elevation as would be expected if they played a major pathogenic role (205, 206). VEGF, by virtue of its diffusible nature and hypoxia inducibility, was an attractive candidate as a mediator of intraocular neovascularization. Accordingly, elevations of VEGF levels in the aqueous and vitreous of eyes with proliferative retinopathy have been

described (207–209). In a large series where ocular fluids from 165 patients were examined, a strong correlation was found between levels of immunoreactive VEGF in the aqueous and vitreous humors and active proliferative retinopathy (207). VEGF levels were undetectable or very low (<0.5 ng/ml) in the eyes of patients affected by nonneovascular disorders or diabetes without proliferative retinopathy. In contrast, VEGF levels were in the range of 3–10 ng/ml in the presence of active proliferative retinopathy associated with diabetes, occlusion of central retinal vein, or prematurity. Remarkably, the VEGF levels were again very low in the eyes of patients with quiescent proliferative retinopathy, a phase of vascular regression that follows the period of active vascular proliferation in diabetic and other retinopathies (207). Thus, although the involvement of other factors cannot be ruled out, VEGF is the molecule that correlates best with ocular angiogenesis (210). In agreement with these findings, *in situ* hybridization studies have demonstrated up-regulation of VEGF mRNA in the retina of patients with proliferative retinopathies secondary to diabetes, central retinal vein occlusion, retinal detachment, or intraocular tumors (211). Interestingly, VEGF mRNA expression was localized to the specific retinal layer(s) expected to be ischemic in each of these conditions (211).

More direct evidence for a role of VEGF as a mediator of intraocular neovascularization has been generated in a primate model of iris neovascularization and in a murine model of retinopathy of prematurity (212, 213). In the former, intraocular administration of anti-VEGF antibodies dramatically inhibits the neovascularization that follows occlusion of central retinal veins (214). Likewise, soluble Flt-1 or Flk-1 fused to an IgG suppresses retinal angiogenesis in the mouse model (215).

Neovascularization is a major cause of visual loss also in AMD, the overall leading cause of blindness (216). Most AMD patients have atrophy of the retinal pigment epithelial cells and characteristic formations called "drusen." A significant percentage of AMD patients (~20%) manifest the neovascular (exudative) form of the disease. In this condition, the new vessels stem from the extraretinal choriocapillary. It is well established that the appearance of such choroid neovascularization coincides with a dramatic worsening in prognosis (216). Leakage and bleeding from these vessels may lead to damage to the macula and ultimately to loss of central vision. Because of the proximity of the lesions to the macula, laser photocoagulation or surgical therapy are of very limited value. Very recent studies have documented the immunohistochemical localization of VEGF in surgically resected choroidal neovascular membranes from AMD patients (217, 218). Transdifferentiated retinal pigment epithelial cells in the highly vascularized regions of the membranes were the major sources of VEGF. Also, Kvantu (219) has shown that choroid fibroblasts are able to release VEGF and also that various cytokines are able to induce VEGF gene expression in these cells. These findings suggest a role for VEGF in the progression of AMD-related choroidal neovascularization, raising the possibility that a pharmacological treatment with monoclonal antibodies or other VEGF inhibitors may constitute a therapy for this condition.

C. Other pathological conditions

Two independent studies have suggested that VEGF is involved in the pathogenesis of rheumatoid arthritis (RA), an important disease in which angiogenesis plays a significant role (220, 221). The RA synovium is characterized by the formation of pannus, an extensively vascularized tissue that invades and destroys the articular cartilage (222). By its vascularity and rapid proliferation rate, the RA synovium has been likened to a tumor (223). Levels of immunoreactive VEGF were high in the synovial fluid of RA patients whereas they were very low or undetectable in the synovial fluid of patients affected by other forms of arthritis or by degenerative joint disease. Furthermore, anti-VEGF antibodies significantly reduced the endothelial cell chemotactic activity of the RA synovial fluid (220).

It has been shown that VEGF expression is increased in psoriatic skin (85). Increased vascularity and permeability are characteristic of psoriasis. Also, VEGF mRNA expression has been examined in three bullous disorders with subepidermal blister formation: bullous pemphigoid, erythema multiforme, and dermatitis herpetiformis (224). In all of these conditions, VEGF mRNA was markedly up-regulated not only in the epidermis over blisters, but also at a distance from blisters, in areas adjacent to dermal inflammatory infiltrates.

Angiogenesis is also important in the pathogenesis of endometriosis, a condition characterized by ectopic endometrium implants in the peritoneal cavity (225). Recently, elevation of VEGF in the peritoneal fluid of patients with endometriosis has been reported (226, 227). Immunohistochemistry indicated that activated peritoneal fluid macrophages as well as tissue macrophages within the ectopic endometrium are the main source of VEGF in this condition. Interestingly, VEGF secretion by macrophages was enhanced by ovarian steroids (226, 227). VEGF up-regulation has been also implicated in the hypervascularity of the ovarian stroma that characterizes Stein-Leventhal syndrome (228).

Moreover, Sato *et al.* (229) proposed that VEGF is implicated in the pathogenesis of Graves' disease. TSH, insulin phorbol ester, (Bu)₂cAMP, and Graves' IgG were found to stimulate VEGF mRNA expression in cultured human thyroid follicles (229). These findings suggest that VEGF, secreted by thyroid follicles via the protein kinase A and C pathways, may be responsible for the characteristic hypervascularity (230) of Graves' syndrome.

Furthermore, it has been suggested that VEGF up-regulation plays a pathogenic role in the capillary hyperpermeability that characterizes ovarian hyperstimulation syndrome (231) as well as in the dysfunctional endothelium of preeclampsia (232).

X. Therapeutic Applications of VEGF-Induced Angiogenesis

The availability of agents able to promote the growth of new collateral vessels would be potentially of major therapeutic value for disorders characterized by inadequate tissue perfusion and might constitute an alternative to surgical reconstruction procedures. For example, chronic limb ischemia, most frequently caused by obstructive atherosclerosis

affecting the superficial femoral artery, is associated with a high rate of morbidity and mortality, and treatment is currently limited to surgical revascularization or endovascular interventional therapy (233, 234). No pharmacological therapy has been shown to be effective for this condition. It has been shown that intraarterial or intramuscular administration of rhVEGF₁₆₅ may significantly augment perfusion and development of collateral vessels in a rabbit model where chronic hindlimb ischemia was created by surgical removal of the femoral artery (21, 22). These studies provided angiographic evidence of neovascularization in the ischemic limbs. Figure 8 illustrates the development of collateral vessels in this ischemic hindlimb model at various time points after a single intraarterial administration of VEGF (1 mg). Arterial gene transfer with cDNA encoding VEGF also led to revascularization in the same rabbit model to an extent comparable to that achieved with the recombinant protein (23, 235). In addition, the hypothesis that the angiogenesis initiated by the administration of VEGF improved muscle function in ischemic limbs was tested by Walder *et al.* (236). A single intraarterial injection of rhVEGF₁₆₅ augmented muscle function in this rabbit model of peripheral limb ischemia. Also, exercise-induced hyperemia was significantly increased in ischemic limbs treated with rhVEGF₁₆₅ (224). Such improvement in perfusion was, however, not seen in other non-ischemic tissues including the contralateral limb. Similarly, Bauters *et al.* (237) have shown that both maximal flow velocity and maximal blood flow are significantly increased in ischemic limbs after VEGF administration. Other studies have shown that VEGF administration also leads to a recovery of normal endothelial reactivity in dysfunctional endothelium. After obstruction of a large artery and development of collateral vessels, the increase in blood flow that normally follows acetylcholine infusion is severely blunted; serotonin paradoxically leads to a decrease in blood flow (238). Thirty days after a single intraarterial bolus of VEGF₁₆₅, restoration of normal increase in blood flow in ischemic rabbit hindlimb after acetylcholine or serotonin infusion was demonstrated (239).

Banai *et al.* (18) have shown that VEGF administration results in increased coronary blood flow in a dog model of coronary insufficiency. After occlusion of the left circumflex coronary artery, daily intraluminal injections of rhVEGF distal to the occlusion resulted in a significant enhancement in collateral blood flow over a 4-week period. In addition, Harada *et al.* (19) have demonstrated that extraluminal administration of as little as 2 µg of rhVEGF by an osmotic pump results in a significant increase in coronary blood flow in a pig model of chronic myocardial ischemia created by ameroid occlusion of the left proximal circumflex artery. Also, magnetic resonance imaging was employed to provide a noninvasive assessment of the benefits secondary to VEGF administration in the porcine model (20). Image series converted to a space-time map demonstrated reduction in the size of the ischemic zone and decreased delay in contrast arrival after VEGF treatment (20). These findings demonstrated improvement in cardiac global and regional function and reduced infarct size, resulting from enhanced collateral blood supply (20).

A further potential therapeutic application of VEGF is the




FIG. 8. Selective internal iliac angiography of control and VEGF-treated rabbit at (a) day 10 (baseline), (b) day 20, and (c) day 40. VEGF was administered as single intraarterial bolus in the iliac artery of rabbits in which the ipsilateral femoral artery had been removed to induce sustained, severe ischemia. Note the modest collateral vessel development in the control, which contrasts with the marked improvement observed after VEGF treatment. [Reproduced with permission from S. Takeshita *et al.*: *J Clin Invest* 93:662–670, 1994 (21) by copyright permission of The American Society for Clinical Investigation.]

prevention of restenosis after percutaneous transluminal angioplasty. Between 15% and 75% of patients undergoing percutaneous transluminal angioplasty for occlusive coronary or peripheral arterial disease develop restenosis within 6 months (234). It has been proposed that damage to the endothelium is a crucial event triggering fibrocellular intimal proliferation (240). Therefore, the induction of rapid reendothelialization may be an effective strategy to prevent the cascade of events leading to neointima formation and ultimately to restenosis in patients. Recent evidence shows that VEGF accelerates reendothelialization and also attenuates intimal hyperplasia in balloon-injured rat carotid artery or rabbit aorta (241, 242).

Very recently, the hypothesis that VEGF may result in therapeutically significant angiogenesis in humans has been tested by Isner *et al.* (243) in a gene therapy trial in patients with severe limb ischemia. A case report of an interim analysis of this trial has been published (24). Arterial gene transfer of 2000 μ g of naked plasmid DNA encoding VEGF₁₆₅, applied to the hydrogel polymer coating of an angioplasty balloon, resulted in angiographic and histological evidence of angiogenesis in the knee, midtibial, and ankle levels 4 weeks after the transfer. Such effects persisted at a 12-week view.

XI. Perspectives

The findings that heterozygous mutations inactivating the VEGF gene and homozygous mutations inactivating the Flt-1 or Flk-1/KDR genes result in profound deficits in vasculogenesis and blood island formation, leading to early intrauterine death, emphasize the pivotal role played by the VEGF/VEGF-receptor system in the development of the vas-

cular system. Future studies, using inducible gene knockout technology (244), should help determine when the embryo is most vulnerable to VEGF deficiency. Such studies may also be of major value in establishing the role of VEGF and its receptors in the maintenance and homeostasis of endothelial cells in the adult animal. It will be of considerable interest also to determine the role of the other members of the VEGF family, both in the context of developmental angiogenesis and in the physiology of the adult animal.

The elucidation of the signal transduction properties of the Flt-1 and Flk-1/KDR receptors may provide the key to dissection of the pathways leading to such fundamental biological events as endothelial cell differentiation, morphogenesis, and angiogenesis. Furthermore, a more complete understanding of the signaling events involving other endothelial cell-specific tyrosine kinases (122, 245, 246) as well as cell adhesion molecules (48, 247) and their interrelation with the VEGF/VEGF receptor system should provide a more integrated view of the biology of vascular cells, both in normal and abnormal circumstances. In this context, recent studies have shown that VEGF-mediated angiogenesis requires a specific vascular integrin pathway, mediated by $\alpha v\beta 5$ (247).

An attractive possibility is that recombinant VEGF or gene therapy with VEGF gene may be used to promote endothelial cell growth and collateral vessel formation. This would represent a novel therapeutic modality for conditions that frequently are refractory to conservative measures and unresponsive to pharmacological therapy. Surprisingly, gene therapy with naked plasmid DNA has resulted in a demonstrable therapeutic effect, both in experimental animals and in humans (23, 24, 235). This raises the possibility that more efficient expression systems such as adenoviral vectors (248)

may achieve even greater pharmacological effect. However, the finding that the VEGF protein is able to promote therapeutic angiogenesis even at minute concentrations (19, 20) suggests that gene therapy may not offer advantages over the recombinant protein.

The high expression of VEGF mRNA in human tumors, the presence of the VEGF protein in ocular fluids of individuals with proliferative retinopathies, and the localization of VEGF in AMD lesions strongly support the hypothesis that VEGF is a key mediator of angiogenesis associated with various disorders. Therefore, anti-VEGF antibodies or other inhibitors of VEGF action such as small molecules inhibiting signal transduction of Flk-1/KDR (249), soluble receptors (118, 126, 215), or antisense oligonucleotides (250) may be of therapeutic value for a variety of malignancies as well as for other disorders, used alone or in combination with other agents. The recent elucidation of the minimal structural elements required for VEGF binding in Flt-1 may lead to the generation of smaller soluble receptors, with improved bioavailability and *in vivo* efficacy (139). Although the safety of such treatment has not been yet established, it is tempting to speculate that an anti-VEGF therapy may have low toxicity, possibly limited to inhibition of wound healing and ovarian and endometrial function, since endothelial cells are essentially quiescent in most adult tissues. Very recently, a humanized version of a high-affinity anti-VEGF monoclonal antibody, which retains the same affinity and efficacy as the original murine antibody, has been generated and may be used in future clinical trials (L. G. Presta *et al.*, submitted).

In conclusion, in spite of the plurality of factors potentially involved in angiogenesis, one specific factor, VEGF, appears to play an irreplaceable role in a variety of physiological and pathological circumstances.

References

1. Hamilton WJ, Boyd JD, Mossman HW 1962 Human Embryology. William & Wilkins, Baltimore
2. Gilbert SF 1988 Developmental Biology. Sinauer, Sunderland, MA
3. Folkman J, Klagsbrun M 1987 Angiogenic factors. *Science* 235: 442-447
4. Folkman J 1991 What is the evidence that tumors are angiogenesis-dependent? *J Natl Cancer Inst* 82:4-6
5. Garner A 1994 Vascular diseases. In: Garner A, Klintworth GK (eds) Pathobiology of Ocular Disease. A Dynamic Approach, ed 2. Marcel Dekker, New York, pp 1625-1710
6. Weidner N, Semple JP, Welch W, Folkman J 1991 Tumor angiogenesis and metastasis. Correlation in invasive breast carcinoma. *N Engl J Med* 324:1-8
7. Horak ER, Leek R, Klenk N, Lejeune S, Smith K, Stuart M, Greenall M, Harris AI 1992 Quantitative angiogenesis assessed by anti-PECAM antibodies: correlation with node metastasis and survival in breast cancer. *Lancet* 340:1120-1124
8. Vartanian RK, Weidner N 1994 Correlation of intratumoral endothelial cell proliferation with microvessel density (tumor angiogenesis) and tumor cell proliferation in breast carcinoma. *Am J Pathol* 144:1188-1194
9. Wakui S, Furusato M, Sasaki H, Akiyama A, Kinoshita I, Asano K, Tokuda T, Aizawa S, Ushigome S 1992 Tumor angiogenesis in prostatic carcinoma with and without bone metastasis: a morphometric study. *J Pathol* 168:257-262
10. Macchiarini P, Fontanini G, Hardin MJ, Squartini F, Angeletti CA 1992 Relation of neovascularization to metastasis of non-small cell lung carcinoma. *Lancet* 340:145-146
11. Bigler SA, Deering RE, Brawer MK 1993 Comparison of microscopic vascularity in benign and malignant prostate tissue. *Hum Pathol* 24:220-226
12. Smith-McCune KS, Weidner N 1994 Demonstration and characterization of the angiogenic properties of cervical dysplasia. *Cancer Res* 54:804-808
13. Folkman J, Shing Y 1992 Angiogenesis. *J Biol Chem* 267:10931-10934
14. Klagsbrun M, D'Amore PA 1991 Regulators of angiogenesis. *Annu Rev Physiol* 53:217-239
15. Ferrara N 1993 Vascular endothelial growth factor. *Trends Cardiovasc Med* 3:244-250
16. Carmeliet P, Ferreira V, Breier G, Pollefeyt S, Kieckens L, Gertsenshtein M, Fahrig M, Vandenhoek A, Harpal K, Eberhardt C, Declercq C, Pawling J, Moons L, Collen D, Risau W, Nagy A 1996 Abnormal blood vessel development and lethality in embryos lacking a single VEGF allele. *Nature* 380:435-439
17. Ferrara N, Carver-Moore K, Chen H, Dowd M, Lu L, O'Shea KS, Powell-Braxton L, Hillan KJ, Moore MW 1996 Heterozygous embryonic lethality induced by targeted inactivation of the VEGF gene. *Nature* 380:439-442
18. Banai S, Jaklitsch MT, Shou M, Lazarous DF, Scheinowitz M, Biro S, Epstein S, Unger E 1994 Angiogenic-induced enhancement of collateral blood flow to ischemic myocardium by vascular endothelial growth factor in dogs. *Circulation* 89:2183-2189
19. Harada K, Friedman M, Lopez J, Wang S, Li J, Prasad PV, Pearlman JD, Edelmann E, Sellke FW, Simons M 1996 Vascular endothelial growth factor in chronic myocardial ischemia. *Am J Physiol* 270:H1791-1802
20. Pearlman JD, Hibberd MG, Chuang ML, Harada K, Lopez JJ, Gladstone SR, Friedman M, Sellke FW, Simons M 1995 Magnetic resonance mapping demonstrates benefits of VEGF-induced myocardial angiogenesis. *Nature Med* 1:1085-1089
21. Takeshita S, Zhong L, Brogi E, Kearney M, Pu L-Q, Bunting S, Ferrara N, Symes JF, Isner JM 1994 Therapeutic angiogenesis: a single intra-arterial bolus of vascular endothelial growth factor augments collateral vessel formation in a rabbit ischemic hindlimb model. *J Clin Invest* 93:662-670
22. Takeshita S, Pu L-Q, Stein LA, Sniderman AD, Bunting S, Ferrara N, Isner JM, Symes JF 1994 Intramuscular administration of vascular endothelial growth factor induces dose-dependent collateral artery augmentation in a rabbit model of chronic limb ischemia. *Circulation* 90:[Suppl II]228-234
23. Takeshita S, Tsurumi Y, Couffinhal T, Asahara T, Bauters C, Symes JF, Ferrara N, Isner JM 1996 Gene transfer of naked DNA encoding for three isoforms of vascular endothelial growth factor stimulates collateral development *in vivo*. *Lab Invest* 75:487-502
24. Isner JM, Pieczek A, Schainfeld R, Blair R, Haley L, Asahara T, Rosenfield K, Razvi S, Walsh K, Symes JF 1996 Clinical evidence of angiogenesis following arterial gene transfer of phVEGF₁₆₅ in patient with ischaemic limb. *Lancet* 348:370-374
25. Ferrara N, Henzel WJ 1989 Pituitary follicular cells secrete a novel heparin-binding growth factor specific for vascular endothelial cells. *Biochem Biophys Res Commun* 161:851-858
26. Leung DW, Cachianes G, Kuang W-J, Goeddel DV, Ferrara N 1989 Vascular endothelial growth factor is a secreted angiogenic mitogen. *Science* 246:1306-1309
27. Connolly DT, Heuvelman DM, Nelson R, Olander JV, Eppley BL, Delfino JJ, Siegel NR, Leimgruber RM, Feder J 1989 Tumor vascular permeability factor stimulates endothelial cell growth and angiogenesis. *J Clin Invest* 84:1470-1478
28. Keck PJ, Hauser SD, Krivi G, Sanzo K, Warren T, Feder J, Connolly DT 1989 Vascular permeability factor, an endothelial cell mitogen related to platelet derived growth factor. *Science* 246: 1309-1312
29. Plouët J, Schilling J, Gospodarowicz D 1989 Isolation and characterization of a newly identified endothelial cell mitogen produced by AtT20 cells. *EMBO J* 8:3801-3806
30. Conn G, Bayne M, Soderman L, Kwok PW, Sullivan KA, Palisi TM, Hope DA, Thomas KA 1990 Amino acid and cDNA sequence of a vascular endothelial cell mitogen homologous to platelet-derived growth factor. *Proc Natl Acad Sci USA* 87:2628-2632
31. Pepper MS, Wasi S, Ferrara N, Orci L, Montesano R 1994 *In vitro*

- angiogenic and proteolytic properties of bovine lymphatic endothelial cells. *Exp Cell Res* 210:298–305
32. **Pepper MS, Ferrara N, Orci L, Montesano R** 1992 Potent synergism between vascular endothelial growth factor and basic fibroblast growth factor in the induction of angiogenesis *in vitro*. *Biochem Biophys Res Commun* 189:824–831
 33. **Nicosia RF, Nicosia SV, Smith M** 1994 Vascular endothelial growth factor, platelet-derived growth factor and insulin-like growth factor-1 promote rat aortic angiogenesis *in vitro*. *Am J Pathol* 145:1023–1029
 34. **Phillips GD, Stone AM, Jones BD, Schultz JC, Whitehead RA, Knighton DR** 1995 Vascular endothelial growth factor (rhVEGF₁₆₅) stimulates direct angiogenesis in the rabbit cornea. *In Vivo* 8:961–965
 35. **Tolentino MJ, Miller JW, Gragoudas ES, Chatzistefanou K, Ferrara N, Adamis AP** 1996 VEGF is sufficient to produce iris neovascularization and neovascular glaucoma in a non-human primate. *Arch Ophthalmol* 114:964–970
 36. **Pepper MS, Ferrara N, Orci L, Montesano R** 1991 Vascular endothelial growth factor (VEGF) induces plasminogen activators and plasminogen activator inhibitor type 1 in microvascular endothelial cells. *Biochem Biophys Res Commun* 181:902–908
 37. **Unemori EN, Ferrara N, Bauer EA, Amento EP** 1992 Vascular endothelial growth factor induces interstitial collagenase expression in human endothelial cells. *J Cell Physiol* 153:557–562
 38. **Pepper MS, Montesano R** 1990 Proteolytic balance and capillary morphogenesis. *Cell Differ Dev* 32:319–331
 39. **Mandriota S, Montesano R, Orci L, Seghezzi G, Vassalli J-D, Ferrara N, Mignatti P, Pepper MS** 1995 Vascular endothelial growth factor increases urokinase receptor expression in vascular endothelial cells. *J Biol Chem* 270:9709–9716
 40. **Mignatti P, Tsuboi R, Robbins E, Rifkin DB** 1989 *In vitro* angiogenesis on the human amniotic membrane: requirement for basic fibroblast growth factor-induced proteinases. *J Cell Biol* 108:671–682
 41. **Senger DR, Galli SJ, Dvorak AM, Perruzzi CA, Harvey VS, Dvorak HF** 1983 Tumor cells secrete a vascular permeability factor that promotes accumulation of ascites fluid. *Science* 219:983–985
 42. **Connolly DT, Olander JV, Heuvelman D, Nelson R, Monsell R, Siegel N, Haymore BL, Leimgruber R, Feder J** 1989 Human vascular permeability factor. Isolation from U937 cells. *J Biol Chem* 264:20017–20024
 43. **Dvorak HF** 1986 Tumors: wounds that do not heal. Similarity between tumor stroma generation and wound healing. *N Engl J Med* 315:1650–1658
 44. **Dvorak HF, Harvey VS, Estrella P, Brown LF, McDonagh J, Dvorak AM** 1987 Fibrin containing gels induce angiogenesis: implications for tumor stroma generation and wound healing. *Lab Invest* 57:673–686
 45. **Roberts WG, Palade G** 1995 Increased microvascular permeability and endothelial fenestration induced by vascular endothelial growth factor. *J Cell Sci* 108:2369–2379
 46. **Dellian M, Witwer BP, Salehi HA, Yuan F, Jain RK** 1996 Quantitation and physiological characterization of angiogenic vessels in mice. Effects of basic fibroblast growth factor, vascular endothelial growth factor/vascular permeability factor, and host microenvironment. *Am J Pathol* 149:59–71
 47. **Pekala P, Marlow M, Heuvelman D, Connolly D** 1990 Regulation of hexose transport in aortic endothelial cells by vascular permeability factor and tumor necrosis factor- α , but not by insulin. *J Biol Chem* 265:18051–18054
 48. **Melder RJ, Koenig GC, Witwer BP, Safabakhsh N, Munn LL, Jain RK** 1996 During angiogenesis, vascular endothelial growth factor and basic fibroblast growth factor regulate natural killer cell adhesion to tumor endothelium. *Nature Med* 2:992–997
 49. **Sasaki A, Melder RJ, Whiteside TL, Herberman RB, Jain RK** 1991 Preferential localization of human adherent lymphokine-activated killer cells in tumor microcirculation. *J Natl Cancer Inst* 83:433–437
 50. **Clauss M, Gerlach M, Gerlach H, Brett J, Wang F, Familletti PC, Pan Y-C, Olander JV, Connolly DT, Stern D** 1990 Vascular permeability factor: a tumor-derived polypeptide that induces endothelial cell and monocyte procoagulant activity, and promotes monocyte migration. *J Exp Med* 172:1535–1545
 51. **Broxmeyer HE, Cooper S, Li ZH, Song HY, Kwon BS, Warren RS, Donner DB** 1995 Myeloid progenitor cells regulatory effects of vascular endothelial growth factor. *Int J Hematol* 62:203–215
 52. **Gabrilovich DI, Chen HL, Girgis KR, Cunningham HT, Meny GM, Nadaf S, Kavanaugh D, Carbone D** 1996 Production of vascular endothelial growth factor by human tumors inhibits the functional maturation of dendritic cells. *Nature Med* 2:1096–1103
 53. **Ku DD, Zaleski JK, Liu S, Brock T** 1993 Vascular endothelial growth factor induces EDRF-dependent relaxation of coronary arteries. *Am J Physiol* 265:H586–592
 54. **Yang R, Thomas GR, Bunting S, Ko A, Keyt B, Ferrara N, Ross J, Jin H** 1996 Effects of VEGF on hemodynamics and cardiac performance. *J Cardiovasc Pharmacol* 27:838–844
 55. **Cuevas P, Garcia-Calvo M, Carceller F, Reimers D, Zazo M, Cuevas B, Munoz-Willery I, Martinez-Coso V, Lamas S, Gimenez-Gallego G** 1996 Correction of hypertension by normalization of endothelial levels of fibroblast growth factor and nitric oxide synthase in spontaneously hypertensive rats. *Proc Natl Acad Sci USA* 93:11996–12001
 56. **Tisher E, Mitchell R, Hartmann T, Silva M, Gospodarowicz D, Fiddes J, Abraham J** 1991 The human gene for vascular endothelial growth factor. *J Biol Chem* 266:11947–11954
 57. **Houck KA, Ferrara N, Winer J, Cachianes G, Li B, Leung DW** 1991 The vascular endothelial growth factor family: identification of a fourth molecular species and characterization of alternative splicing of RNA. *Mol Endocrinol* 5:1806–1814
 58. **Vincenti V, Cassano C, Rocchi M, Persico G** 1996 Assignment of the vascular endothelial growth factor gene to the human chromosome 6p21.3. *Circulation* 93:1493–1495
 59. **Shima DT, Kuroki M, Deutsch U, Ng Y-S, Adamis AP, D'Amore PA** 1996 The mouse gene for vascular endothelial growth factor. *J Biol Chem* 271:3877–3883
 60. **Levy AP, Levy NS, Wegner S, Goldberg MA** 1995 Transcriptional regulation of the rat vascular endothelial growth factor gene by hypoxia. *J Biol Chem* 270:13333–13340
 61. **Ferrara N, Houck K, Jakeman L, Leung DW** 1992 Molecular and biological properties of the vascular endothelial growth factor family of proteins. *Endocr Rev* 13:18–32
 62. **Houck KA, Leung DW, Rowland AM, Winer J, Ferrara N** 1992 Dual regulation of vascular endothelial growth factor bioavailability by genetic and proteolytic mechanisms. *J Biol Chem* 267:26031–26037
 63. **Park JE, Keller G-A, Ferrara N** 1993 The vascular endothelial growth factor (VEGF) isoforms: differential deposition into the subepithelial extracellular matrix and bioactivity of ECM-bound VEGF. *Mol Biol Cell* 4:1317–1326
 64. **Rifkin DB, Moscatelli D, Bizik J, Quarto N, Blei F, Dennis P, Flaumenhaft R, Mignatti P** 1990 Growth factor control of extracellular proteolysis. *Cell Differ Dev* 32:313–318
 65. **Sreenath T, Matrisian LM, Stetler-Stevenson W, Gattoni-Celli S, Pozzatti RO** 1992 Expression of matrix metalloproteinases in transfected rat cell lines of high and low metastatic potential. *Cancer Res* 52:4942–4947
 66. **Keyt B, Berleau L, Nguyen H, Heinshon H, Chen H, Vandlen R, Ferrara N** 1996 The carboxyl-terminal domain (111–165) of VEGF is critical for mitogenic potency. *J Biol Chem* 271:7788–7795
 67. **Minchenko A, Bauer T, Salceda S, Caro J** 1994 Hypoxic stimulation of vascular endothelial growth factor expression *in vivo* and *in vitro*. *Lab Invest* 71:374–379
 68. **Shweiki D, Itin A, Soffer D, Keshet E** 1992 Vascular endothelial growth factor induced by hypoxia may mediate hypoxia-initiated angiogenesis. *Nature* 359:843–845
 69. **Shima DT, Adamis AP, Ferrara N, Yeo K-T, Yeo T-K, Allende M, Folkman J, D'Amore PA** 1995 Hypoxic induction of vascular endothelial cell growth factors in the retina: identification and characterization of vascular endothelial growth factor (VEGF) as the sole mitogen. *Mol Med* 2:64–71
 70. **Brogi E, Wu T, Namiki A, Isner JM** 1994 Indirect angiogenic cytokines upregulate VEGF and bFGF expression in vascular smooth muscle cells, whereas hypoxia upregulates VEGF expression only. *Circulation* 90:649–652
 71. **Banai S, Shweiki D, Pinson A, Chandra M, Lazarovici G, Keshet E** 1994 Upregulation of vascular endothelial growth factor expres-

- sion induced by myocardial ischemia: implications for coronary angiogenesis. *Cardiovasc Res* 28:1176–1179
72. Stone J, Itin A, Alon T, Pe'er J, Gnossin H, Chan-Ling T, Keshet E 1995 Development of retinal vasculature is mediated by hypoxia-induced vascular endothelial growth factor (VEGF) expression in neuroglia. *J Neurosci* 15:4738–4747
 73. Goldberg MA, Schneider TJ 1994 Similarities between the oxygen-sensing mechanisms regulating the expression of vascular endothelial growth factor and erythropoietin. *J Biol Chem* 269:4355–4361
 74. Liu Y, Cox SR, Morita T, Kourembanas S 1995 Hypoxia regulates vascular endothelial growth factor gene expression in endothelial cells. Identification of a 5' enhancer. *Circ Res* 77:638–643
 75. Madan A, Curtin PT 1993 A 24-base pair sequence 3' to the human erythropoietin contains a hypoxia-responsive transcriptional enhancer. *Proc Natl Acad Sci USA* 90:3928–3932
 76. Wang GL, Semenza GL 1995 Purification and characterization of hypoxia-inducible factor-1. *J Biol Chem* 270:1230–1237
 77. Wang GL, Jiang BH, Rue EA, Semenza GL 1995 Hypoxia-inducible factor-1 is a basic helix-loop helix-PAS heterodimer regulated by cellular O₂ tension. *Proc Natl Acad Sci USA* 92:5510–5514
 78. Forsythe JA, Jiang BH, Iyer NV, Agani F, Leung SW, Koos RD, Semenza GL 1996 Activation of vascular endothelial growth factor gene transcription by hypoxia-inducible factor 1. *Mol Cell Biol* 16:4604–4613
 79. Takagi H, King GL, Robinson GS, Ferrara N, Aiello LP 1996 Hypoxic induction of VEGF is mediated by adenosine through A₂ receptors and elevation of cAMP in retinal pericytes and endothelial cells. *Invest Ophthalmol Vis Sci* 37:2165–2176
 80. Mukhopadhyay D, Tsiokas L, Zhou X-M, Foster D, Brugge JS, Sukhatme VP 1995 Hypoxic induction of human vascular endothelial growth factor expression through c-Src activation. *Nature* 375:577–581
 81. Levy AP, Levy NS, Goldberg MA 1996 Post-transcriptional regulation of vascular endothelial growth factor by hypoxia. *J Biol Chem* 271:2746–2753
 82. Ikeda E, Achen MG, Breier G, Risau W 1995 Hypoxia-induced transcriptional activation and increased mRNA stability of vascular endothelial growth factor in C6 glioma cells. *J Biol Chem* 270:19761–19766
 83. Stein I, Neeman M, Shweiki D, Itin A, Keshet E 1995 Stabilization of vascular endothelial growth factor mRNA by hypoxia and hypoglycemia and coregulation with other ischemia-induced genes. *Mol Cell Biol* 15:5363–5368
 84. Frank S, Hubner G, Breier G, Longaker MT, Greenhalgh DG, Werner S 1995 Regulation of VEGF expression in cultured keratinocytes. Implications for normal and impaired wound healing. *J Biol Chem* 270:12607–12613
 85. Detmar M, Brown LF, Claffey KP, Yeo KT, Kocher O, Jackman RW, Berse B, Dvorak HF 1994 Overexpression of vascular permeability factor/vascular endothelial growth factor and its receptors in psoriasis. *J Exp Med* 180:1141–1146
 86. Detmar M, Yeo KT, Nagy JA, Van de Water L, Brown LF, Berse B, Elicker BM, Ledbetter S, Dvorak HF 1995 Keratinocyte-derived vascular permeability factor (vascular endothelial growth factor) is a potent mitogen for dermal microvascular endothelial cells. *J Invest Dermatol* 105:44–50
 87. Goldman CK, Kim J, Wong WL, King V, Brock T, Gillespie GY 1993 Epidermal growth factor stimulates vascular endothelial growth factor production by human malignant glioma cells: a model of glioblastoma multiforme pathophysiology. *Mol Biol Cell* 4:121–133
 88. Pertovaara L, Kaipainen A, Mustonen T, Orpana A, Ferrara N, Saksela O, Alitalo K 1994 Vascular endothelial growth factor is induced in response to transforming growth factor- β in fibroblastic and epithelial cells. *J Biol Chem* 269:6271–6274
 89. Li J, Perrella MA, Tsai JC, Yet SF, Hsieh CM, Yoshizumi M, Patterson C, Endego WO, Zhou F, Lee M 1995 Induction of vascular endothelial growth factor gene expression by interleukin-1 beta in rat aortic smooth muscle cells. *J Biol Chem* 270:308–312
 90. Ben-Av P, Crofford LJ, Wilder RL, Hla T 1995 Induction of vascular endothelial growth factor expression in synovial fibroblasts by prostaglandin E and interleukin-1: a potential mechanism for inflammatory angiogenesis. *FEBS Lett* 372:83–87
 91. Cohen T, Nahari D, Cerem LW, Neufeld G, Levi BZ 1996 Interleukin 6 induces the expression of vascular endothelial growth factor. *J Biol Chem* 271:736–741
 92. Warren RS, Yuan H, Matli MR, Ferrara N, Donner DB 1996 Induction of vascular endothelial growth factor by insulin-like growth factor 1 in colorectal carcinoma. *J Biol Chem* 271:29483–29488
 93. Claffey KP, Wilkinson WO, Spiegelman BM 1992 Vascular endothelial growth factor. Regulation by cell differentiation and activated second messenger pathways. *J Biol Chem* 267:16317–16322
 94. Garrido C, Saule S, Gospodarowicz D 1993 Transcriptional regulation of vascular endothelial growth factor gene expression in ovarian bovine granulosa cells. *Growth Factors* 8:109–117
 95. Kieser A, Weich H, Brandner G, Marme D, Kolch W 1994 Mutant p53 potentiates protein kinase C induction of vascular endothelial growth factor expression. *Oncogene* 9:963–969
 96. Rak J, Mitsuhashi Y, Bayko L, Filmus J, Shirasawa S, Sasazuki T, Kerbel RS 1995 Mutant ras oncogenes upregulate VEGF/VPF expression: implications for induction and inhibition of tumor angiogenesis. *Cancer Res* 55:4575–4580
 97. Grugel S, Finkenzeller G, Weindel K, Barleon B, Marme D 1995 Both v-Ha-Ras and v-Raf stimulate expression of vascular endothelial growth factor in NIH 3T3 cells. *J Biol Chem* 270:25915–25919
 98. Mazure NM, Chen EY, Yeh P, Laderout KR, Giaccia AJ 1996 Oncogenic transformation and hypoxia synergistically act to modulate vascular endothelial growth factor expression. *Cancer Res* 56:3436–3440
 99. Mukhopadhyay D, Tsiokas L, Sukhatme VP 1995 Wild-type p53 and v-Src exert opposing influences on human vascular endothelial growth factor gene expression. *Cancer Res* 55:6161–6165
 100. Siemester G, Weindel K, Mohrs K, Barleon B, Martiny-Baron G, Marme D 1996 Reversion of deregulated expression of vascular endothelial growth factor in human renal carcinoma cells by von Hippel-Lindau tumor suppressor protein. *Cancer Res* 56:2299–2303
 101. Larroche J-C 1984 Malformations of the nervous system. In: Adams, JH, Corsellis, JAN, Duchon, LW (eds) *Greenfield's Neuropathology*, ed 4. Wiley Medical Publishers. New York, pp 385–450
 102. Iliopoulos O, Levy AP, Jiang C, Kaelin WG, Goldberg MA 1996 Negative regulation of hypoxia-inducible genes by the von Hippel-Lindau protein. *Proc Natl Acad Sci USA* 93:10595–10599
 103. Vaisman N, Gospodarowicz D, Neufeld G 1990 Characterization of the receptors for vascular endothelial growth factor. *J Biol Chem* 265:19461–19469
 104. Ploüet J, Moukadir HJ 1990 Characterization of the receptors to vasculotropin on bovine adrenal cortex-derived capillary endothelial cells. *J Biol Chem* 265:22071–22075
 105. Shen H, Clauss M, Ryan J, Schmidt AM, Tijburg P, Borden L, Connoll DT, Stern D, Kao J 1993 Characterization of vascular permeability factor/vascular endothelial growth factor receptors in mononuclear phagocytes. *Blood* 81:2767–2773
 106. Gitay-Goren H, Cohen T, Tessler S, Soker S, Gengrinovitch S, Rockwell P, Klagsbrun M, Levi BZ, Neufeld G 1996 Selective binding of VEGF₁₂₁ to one of the three vascular endothelial growth factor receptors on vascular endothelial cells. *J Biol Chem* 271:5519–5523
 107. Soker S, Fidler IJ, Neufeld G, Klagsbrun M 1996 Characterization of novel VEGF receptors on tumor cells that bind VEGF₁₆₅ via its exon 7-encoded domain. *J Biol Chem* 271:5761–5767
 108. Jakeman LB, Winer J, Bennett GL, Altar CA, Ferrara N 1992 Binding sites for vascular endothelial growth factor are localized on endothelial cells in adult rat tissues. *J Clin Invest* 89:244–253
 109. Jakeman LB, Armanini M, Phillips HS, Ferrara N 1993 Developmental expression of binding sites and mRNA for vascular endothelial growth factor suggests a role for this protein in vasculogenesis and angiogenesis. *Endocrinology* 133:848–859
 110. deVries C, Escobedo JA, Ueno H, Houck KA, Ferrara N, Williams LT 1992 The fms-like tyrosine kinase, a receptor for vascular endothelial growth factor. *Science* 255:989–991
 111. Terman BI, Vermazen MD, Carrion ME, Dimitrov D, Armellino DC, Gospodarowicz D, Bohlen P 1992 Identification of the KDR

- tyrosine kinase as a receptor for vascular endothelial growth factor. *Biochem Biophys Res Commun* 34:1578–1586
112. Shibuya M, Yamaguchi S, Yamane A, Ikada T, Tojo T, Matsu-shime H, Sato M 1990 Nucleotide sequence and expression of a novel human receptor-type tyrosine kinase (*flt*) closely related to the *fms* family. *Oncogene* 8:519–527
 113. Terman BI, Carrion ME, Kovacs E, Rasmussen BA, Eddy RL, Shows TB 1991 Identification of a new endothelial cell growth factor receptor tyrosine kinase. *Oncogene* 6:519–524
 114. Matthews W, Jordan CT, Gavin M, Jenkins NA, Copeland NG, Lemischka IR 1991 A receptor tyrosine kinase cDNA isolated from a population of enriched primitive hematopoietic cells and exhibiting close genetic linkage to c-kit. *Proc Natl Acad Sci USA* 88: 9026–9030
 115. Quinn T, Peters KG, deVries C, Ferrara N, Williams LT 1993 Fetal liver kinase 1 is a receptor for vascular endothelial growth factor and is selectively expressed in vascular endothelium. *Proc Natl Acad Sci USA* 90:7533–7537
 116. Millauer B, Witzmann-Voos S, Schnurch H, Martinez R, Moller NP, Risau W, Ullrich A 1993 High affinity binding and developmental expression suggest Flk-1 as a major regulator of vasculogenesis and angiogenesis. *Cell* 72:835–846
 117. Kendall RL, Thomas KA 1993 Inhibition of vascular endothelial growth factor by an endogenously encoded soluble receptor. *Proc Natl Acad Sci USA* 90:10705–10709
 118. Kendall RL, Wang G, Thomas KA 1996 Identification of a soluble form of the vascular endothelial growth factor receptor FLT-1 and its heterodimerization with KDR. *Biochem Biophys Res Commun* 226:324–328
 119. Pajusola K, Aprelikova O, Korhonen J, Kaipainen A, Pertovaara L, Alitalo K 1992 FLT4 receptor tyrosine kinase contains seven immunoglobulin-like loops and is expressed in multiple human tissues and cell lines. *Cancer Res* 52:5738–5743
 120. Finnerty H, Kelleher K, Morris GE, Bean K, Merberg DM, Kriz R, Morris JC, Sookdeo H, Turner KJ, Wood CR 1993 Molecular cloning of murine FLT and FLT4. *Oncogene* 8:2293–2298
 121. Pajusola K, Aprelikova O, Pelicci G, Weich H, Claesson-Welsh L, Alitalo K 1994 Signalling properties of FLT4, a proteolytically processed receptor tyrosine kinase related to two VEGF receptors. *Oncogene* 9:3545–3555
 122. Mustonen T, Alitalo K 1995 Endothelial receptor tyrosine kinases involved in angiogenesis. *J Cell Biol* 129:895–898
 123. Guo D, Jia Q, Song HY, Warren RS, Donner D 1995 Vascular endothelial growth factor promotes tyrosine phosphorylation of mediators of signal transduction that contain SH2 domains. Association with endothelial cell proliferation. *J Biol Chem* 270:6729–6733
 124. Waltenberger J, Claesson-Welsh L, Siegbahn A, Shibuya M, Heldin C-H 1994 Different signal transduction properties of KDR and Flt1, two receptors for vascular endothelial growth factor. *J Biol Chem* 269:26988–26995
 125. Seetharam L, Gotoh N, Maru Y, Neufeld G, Yamaguchi S, Shibuya M 1995 A unique signal transduction pathway for the FLT tyrosine kinase, a receptor for vascular endothelial growth factor. *Oncogene* 10:135–147
 126. Park JE, Chen H, Winer J, Houck K, Ferrara N 1994 Placenta growth factor. Potentiation of vascular endothelial growth factor bioactivity, *in vitro* and *in vivo*, and high affinity binding to Flt-1 but not to Flk-1/KDR. *J Biol Chem* 269:25646–25654.
 127. Keyt B, Nguyen H, Berleau L, Duarte C, Park J, Chen H, Ferrara N 1996 Identification of VEGF determinants for binding Flt-1 and KDR receptors. Generation of receptor-selective VEGF variants by site-directed mutagenesis. *J Biol Chem* 271:5638–5646
 128. Cunningham SA, Waxham MN, Arrate PM, Brock TA 1995 Interaction of the Flt-1 tyrosine kinase receptor with the p85 subunit of phosphatidylinositol 3-kinase. *J Biol Chem* 270:20254–20257
 129. Barleon B, Sozzani S, Zhou D, Weich H, Mantovani A, Marmé D 1996 Migration of human monocytes in response to vascular endothelial growth factor (VEGF) is mediated via the VEGF receptor flt-1. *Blood* 87:3336–3343
 130. Morishita K, Johnson DE, Williams LT 1995 A novel promoter for vascular endothelial growth factor receptor (*flt-1*) that confers endothelial-specific gene expression. *J Biol Chem* 270:27948–27953
 131. Patterson C, Perrella MA, Hsieh C-M, Yoshizumi M, Lee M-E, Haber E 1995 Cloning and functional analysis of the promoter for KDR/Flk-1, a receptor for vascular endothelial growth factor. *J Biol Chem* 270:23111–23118
 132. Tuder RM, Flook BE, Voelkel NF 1995 Increased gene expression for VEGF and the VEGF receptors KDR/Flk and Flt in lungs exposed to acute or chronic hypoxia. *J Clin Invest* 95:1798–1807
 133. Li J, Brown LF, Hibberd MG, Grossman JD, Morgan JP, Simons M 1996 VEGF, Flk-1 and Flt-1 expression in a rat myocardial infarction model of angiogenesis. *Am J Physiol* 270:H1803–1811
 134. Thieme H, Aiello LP, Ferrara N, King GL 1995 Comparative analysis of VEGF receptors on retinal and aortic endothelial cells. *Diabetes* 44:98–103
 135. Takagi H, King GL, Ferrara N, Aiello LP 1996 Hypoxia regulates VEGF receptor KDR/Flk-1 gene expression through adenosine A₂ receptors in retinal capillary endothelial cells. *Invest Ophthalmol Vis Sci* 37:1311–1321
 136. Brogi E, Schatteman G, Wu T, Kim EA, Varticovski L, Keyt B, Isner JM 1996 Hypoxia-induced paracrine regulation of vascular endothelial growth factor receptor expression. *J Clin Invest* 97: 469–476
 137. Patterson C, Perrella MA, Endege WO, Yoshizumi M, Lee M-E, Haber E 1996 Downregulation of vascular endothelial growth factor receptors by tumor necrosis factor- α in cultured human vascular endothelial cells. *J Clin Invest* 98:4900–4906
 138. Mandriota SJ, Menoud PA, Pepper MS 1996 Transforming growth factor beta 1 down-regulates vascular endothelial growth factor receptor 2/flk-1 expression in vascular endothelial cells. *J Biol Chem* 271:11500–11505
 139. Davis-Smyth T, Chen H, Park J, Presta LG, Ferrara N 1996 The second immunoglobulin-like domain of the VEGF tyrosine kinase receptor Flt-1 determines ligand binding and may initiate a signal transduction cascade. *EMBO J* 15:4919–4927
 140. Williams AF, Barclay AN 1988 The immunoglobulin superfamily-domains for cell surface recognition. *Annu Rev Immunol* 6:381–405
 141. Heldin C-H 1995 Dimerization of cell surface receptors in signal transduction. *Cell* 80:213–223
 142. Ferrara N, Keyt B 1997 Vascular endothelial growth factor. Basic biology and clinical implications. In: Rosen E, Goldberg ID (eds) *Regulation of Angiogenesis*. Springer Verlag, New York, pp 209–232
 143. Oefner C, D'Arcy A, Winkler FK, Eggmann B, Hosang M 1992 Crystal structure of human platelet-derived growth factor BB. *EMBO J* 11:3921–3926
 144. Maglione D, Guerriero V, Viglietto G, Delli-Bovi P, Persico MG 1991 Isolation of a human placenta cDNA coding for a protein related to the vascular permeability factor. *Proc Natl Acad Sci USA* 88:9267–9271
 145. Hauser S, Weich HA 1993 A heparin-binding form of placenta growth factor (PIGF-2) is expressed in human umbilical vein endothelial cells and in placenta. *Growth Factors* 9:259–268
 146. DiSalvo J, Bayne ML, Conn G, Kwok PW, Trivedi PG, Soderman DD, Palisi TM, Sullivan KA, Thomas KA 1995 Purification and characterization of a naturally occurring vascular endothelial growth factor. placenta growth factor heterodimer. *J Biol Chem* 270:7717–7723
 147. Cao Y, Chen H, Zhou L, Chiang MK, Anand-Apte B, Weatherbee JA, Wang Y, Fang F, Flanagan JG, Tsang ML 1996 Heterodimers of placenta growth factor/vascular endothelial growth factor. Endothelial activity, tumor cell expression, and high affinity binding to Flk-1/KDR. *J Biol Chem* 271:3154–3162
 148. Kaipainen A, Korhonen K, Mustonen V, van Hinsberg G-H, Fang D, Dumont D, Breitman M, Alitalo K 1995 Expression of the FLT4 receptor tyrosine kinase becomes restricted to lymphatic endothelium during development. *Proc Natl Acad Sci USA* 92:3566–3570
 149. Joukov V, Pajusola K, Kaipainen A, Chilov D, Lahtinen I, Kukk E, Saksela O, Kalkkinen N, Alitalo K 1996 A novel vascular endothelial growth factor, VEGF-C, is a ligand for the FLT4 (VEGFR-3) and KDR (VEGFR-2) receptor tyrosine kinases. *EMBO J* 15:290–298
 150. Lee J, Gray A, Yuan J, Luoh S-M, Avraham H, Wood WI 1996 Vascular endothelial growth factor-related protein: a ligand and

- specific activator of the tyrosine kinase receptor Flt4. *Proc Natl Acad Sci USA* 93:1988–1992
151. Olofsson B, Pajusola K, Kaipainen A, VonEuler G, Joukov V, Saksela O, Orpana A, Pettersson R, Alitalo K, Eriksson U 1996 Vascular endothelial growth factor B, a novel growth factor for endothelial cells. *Proc Natl Acad Sci USA* 93:2576–2581
 152. Grimmer S, Lagencrantz J, Drinkwater C, Silins G, Townson S, Pollock P, Gotley D, Carson E, Rakar S, Nordenskjöld M, Ward L, Hayward N, Weber G 1996 Cloning and characterization of a human gene related to vascular endothelial growth factor. *Genome Res* 6:124–131
 153. Lyttle DJ, Fraser KM, Flemings SB, Mercer AA, Robinson AJ 1994 Homologs of vascular endothelial growth factor are encoded by the poxvirus orf virus. *J Virol* 68:84–92
 154. Phillips HS, Hains J, Leung DW, Ferrara N 1990 Vascular endothelial growth factor is expressed in rat corpus luteum. *Endocrinology* 127:965–968
 155. Ravindranath N, Little-Ihrig L, Phillips HS, Ferrara N, Zeleznick AJ 1992 Vascular endothelial growth factor mRNA expression in the primate ovary. *Endocrinology* 131:254–260
 156. Cullinan-Bove K, Koos RD 1993 Vascular endothelial growth factor/vascular permeability factor in the rat uterus: rapid stimulation by estrogen correlates with estrogen-induced increases in uterine capillary permeability and growth. *Endocrinology* 133:829–837
 157. Shweiki D, Itin A, Neufeld G, Gitay-Goren H, Keshet E 1993 Patterns of expression of vascular endothelial growth factor (VEGF) and VEGF receptors in mice suggest a role in hormonally regulated angiogenesis. *J Clin Invest* 91:2235–2243
 158. Breier G, Albrecht U, Sterrer S, Risau W 1992 Expression of vascular endothelial growth factor during embryonic angiogenesis and endothelial cell differentiation. *Development* 114:521–532
 159. Shifren JL, Doldi N, Ferrara N, Mesiano S, Jaffe RB 1994 In the human fetus, vascular endothelial growth factor (VEGF) is expressed in epithelial cells and myocytes, but not vascular endothelium: implications for mode of action. *J Clin Endocrinol Metab* 79:316–322
 160. Yamaguchi TP, Dumont DJ, Conlon RA, Breitman ML, Rossant J 1993 Flk-1, an Flt-related receptor tyrosine kinase, is an early marker for endothelial cell precursors. *Development* 118:489–498
 161. Peters KG, deVries C, Williams LT 1993 Vascular endothelial growth factor receptor expression during embryogenesis and tissue repair suggests a role in endothelial differentiation and blood vessel growth. *Proc Natl Acad Sci USA* 90:8915–8919
 162. Monacci W, Merrill M, Oldfield E 1993 Expression of vascular permeability factor/vascular endothelial growth factor in normal rat tissues. *Am J Physiol* 264:C995–1002
 163. Brown LF, Berse B, Tognazzi K, Manseau EJ, Van de Water L, Senger DR, Dvorak HF, Rosen S 1992 Vascular permeability factor mRNA and protein expression in human kidney. *Kidney Int* 42:1457–1461
 164. Alon T, Hemo I, Itin A, Pe'er J, Stone J, Keshet E 1995 Vascular endothelial growth factor acts as a survival factor for newly formed retinal vessels and has implications for retinopathy of prematurity. *Nature Med* 1:1024–1028
 165. Fong G-H, Rossant J, Gertenstein M, Breitman M 1995 Role of Flt-1 receptor tyrosine kinase in regulation of assembly of vascular endothelium. *Nature* 376:66–67
 166. Shalabi F, Rossant J, Yamaguchi TP, Gertenstein M, Wu X-F, Breitman ML, Schuh AC 1995 Failure of blood island formation and vasculogenesis in Flk-1 deficient mice. *Nature* 376:62–66
 167. Theiler K 1989 *The House Mouse. Atlas of Embryonic Development*. Springer-Verlag, New York
 168. Brandon EP, Idzerda RL, McKnight GS 1995 Targeting the mouse genome: a compendium of knockouts (part III). *Curr Biol* 5:873–881
 169. Mattern J, Koomagi R, Volm M 1996 Association of vascular endothelial growth factor expression with intratumoral microvessel density and tumour cell proliferation in human epidermoid lung carcinoma. *Br J Cancer* 73:931–934
 170. Viglietto G, Maglione D, Rambaldi M, Cerutti J, Romano A, Trapasso F, Fede M, Ippolito P, Chiappetta G, Botti G, Persico MG 1995 Upregulation of vascular endothelial growth factor (VEGF) and downregulation of placenta growth factor (PlGF) associated with malignancy in human thyroid tumors and cell lines. *Oncogene* 11:1569–1579
 171. Brown LF, Berse B, Jackman RW, Guidi AJ, Dvorak HF, Senger DR, Connolly JL, Schnitt SJ 1995 Expression of vascular permeability factor (vascular endothelial growth factor) and its receptors in breast cancer. *Hum Pathol* 26:86–91
 172. Yoshiji H, Gomez DE, Shibuya M, Thorgeirsson UP 1996 Expression of vascular endothelial growth factor, its receptors, and other angiogenic factors in breast cancer. *Cancer Res* 56:2013–2016
 173. Brown LF, Berse B, Jackman RW, Tognazzi K, Manseau EJ, Senger DR, Dvorak HF 1993 Expression of vascular permeability factor (vascular endothelial growth factor) and its receptors in adenocarcinomas of the gastrointestinal tract. *Cancer Res* 53:4727–4735
 174. Suzuki K, Hayashi N, Miyamoto Y, Yamamoto M, Ohkawa K, Ito Y, Sasaki Y, Yamaguchi Y, Nakase H, Noda K, Enomoto N, Arai K, Yamada Y, Yoshihara H, Tujimura T, Kawano K, Yoshikawa K, Kamada T 1996 Expression of vascular permeability factor/vascular endothelial growth factor in human hepatocellular carcinoma. *Cancer Res* 56:3004–3009
 175. Brown LF, Berse B, Jackman RW, Tognazzi K, Manseau EJ, Dvorak HF, Senger DR 1993 Increased expression of vascular permeability factor (vascular endothelial growth factor) and its receptors in kidney and bladder carcinomas. *Am J Pathol* 143:1255–1262
 176. Olson TA, Mohanraj D, Carson LF, Ramakrishnan S 1994 Vascular permeability factor gene expression in normal and neoplastic human ovaries. *Cancer Res* 54:276–280
 177. Guidi AJ, Abu-Jawdeh G, Berse B, Jackman RW, Tognazzi K, Dvorak HF, Brown LF 1995 Vascular permeability factor (vascular endothelial growth factor) expression and angiogenesis in cervical neoplasia. *J Natl Cancer Inst* 87:12137–12145
 178. Hashimoto M, Ohsawa M, Ohnishi A, Naka N, Hirota S, Kitamura Y, Aozasa K 1995 Expression of vascular endothelial growth factor and its receptor mRNA in angiosarcoma. *Lab Invest* 73:859–863
 179. Viglietto G, Romano A, Maglione D, Rambaldi M, Paoletti I, Lago CT, Califano D, Monaco C, Mineo A, Santelli G, Manzo G, Botti G, Chiappetta G, Persico MG 1996 Neovascularization in human germ cell tumors correlates with a marked increase in the expression of vascular endothelial growth factor but not placenta growth factor. *Oncogene* 13:577–587
 180. Plate KH, Breier G, Weich HA, Risau W 1992 Vascular endothelial growth is a potential tumour angiogenesis factor *in vivo*. *Nature* 359:845–847
 181. Phillips HS, Armanini M, Stavrou D, Ferrara N, Westphal M 1993 Intense focal expression of vascular endothelial growth factor mRNA in human intracranial neoplasms: Association with regions of necrosis. *Int J Oncol* 2:913–919
 182. Berkman RA, Merrill MJ, Reinhold WC, Monacci WT, Saxena A, Clark WC, Robertson JT, Ali IU, Oldfield EH 1993 Expression of the vascular permeability/vascular endothelial growth factor gene in central nervous system neoplasms. *J Clin Invest* 91:153–159
 183. Witzigmann-Voss S, Breier G, Risau W, Plate 1994 Up-regulation of vascular endothelial growth factor and its receptors in von Hippel-Lindau disease-associated and sporadic hemangioblastoma. *Cancer Res* 55:1358–1364
 184. Dvorak HF, Brown LF, Detmar M, Dvorak AM 1995 Vascular permeability factor/vascular endothelial growth factor, microvascular permeability and angiogenesis. *Am J Pathol* 146:1029–1039
 185. Plate KH, Breier G, Weich H, Mennel HD, Risau W 1994 Vascular endothelial growth factor and glioma angiogenesis: coordinate induction of VEGF receptors, distribution of VEGF protein and possible *in vivo* regulatory mechanisms. *Int J Cancer* 59:520–529
 186. Ferrara N, Winer J, Burton T, Rowland A, Siegel M, Phillips HS, Terrell T, Keller G-A, Levinson AD 1993 Expression of vascular endothelial growth factor does not promote transformation but confers a growth advantage *in vivo* to Chinese hamster ovary cells. *J Clin Invest* 91:160–170
 187. Freeman MR, Schneck FX, Gagnon ML, Corless C, Soker S, Niknejad K, Peoples GE, Klagsbrun M 1995 Peripheral blood T lymphocytes and lymphocytes infiltrating human cancers express vascular endothelial growth factor: a potential role for T cells in angiogenesis. *Cancer Res* 55:4140–4145

188. Qu-Hong J, Nagy JA, Senger DR, Dvorak HF, Dvorak AM 1995 Ultrastructural localization of VPF/VEGF to the abluminal plasma membrane and vesiculovacuolar organelles of tumor microvascular endothelium. *J Histochem Cytochem* 43:381-389
189. Kondo A, Asano M, Matsuo K, Ohmori I, Suzuki H 1994 Vascular endothelial growth factor/vascular permeability factor is detectable in the sera of tumor bearing mice and cancer patients. *Biochim Biophys Acta* 1221:211-214
190. Toi M, Hoshima S, Takayanagi T, Tominaga T 1994 Association of vascular endothelial growth factor expression with tumor angiogenesis and with early relapse in primary breast cancer. *Jpn J Cancer Res* 85:1045-1049
191. Maeda K, Chung YS, Ogawa Y, Takatsuka S, Kang SM, Ogawa M, Sawada T, Sowa M 1996 Prognostic value of vascular endothelial growth factor expression in gastric carcinoma. *Cancer* 77:858-863
192. Kim KJ, Li B, Houck K, Winer J, Ferrara N 1992 The vascular endothelial growth factor proteins: identification of biologically relevant regions by neutralizing monoclonal antibodies. *Growth Factors* 7:53-64
193. Kim KJ, Li B, Winer J, Armanini M, Gillett N, Phillips HS, Ferrara N 1993 Inhibition of vascular endothelial growth factor-induced angiogenesis suppresses tumour growth *in vivo*. *Nature* 362:841-844
194. Warren RS, Yuan H, Matli MR, Gillett NA, Ferrara N 1995 Regulation by vascular endothelial growth factor of human colon cancer tumorigenesis in a mouse model of experimental liver metastasis. *J Clin Invest* 95:1789-1797
195. Borgström P, Hillan KJ, Sriramam Rao P, Ferrara N 1996 Complete inhibition of angiogenesis and growth of microtumors by anti-vascular endothelial growth factor neutralizing antibodies. Novel concepts of angiostatic therapy from intravital videomicroscopy. *Cancer Res* 56:4032-4039
196. Melnyk O, Schuman MA, Kim KJ 1996 Vascular endothelial growth factor promotes tumor dissemination by a mechanism distinct from its effect on primary tumor growth. *Cancer Res* 56:921-924
197. Asano M, Yukita A, Matsumoto T, Kondo S, Suzuki H 1995 Inhibition of tumor growth and metastasis by an immunoneutralizing monoclonal antibody to vascular endothelial growth factor/vascular permeability factor. *Cancer Res* 55:5296-5301
198. Yuan F, Chen Y, Dellian M, Safabakhsh N, Ferrara N, Jain RK 1996 Time-dependent vascular regression and permeability changes in established xenografts induced by an anti-vascular endothelial growth factor/vascular permeability factor antibody. *Proc Natl Acad Sci USA* 93:14765-14770
199. Millauer B, Shawver LK, Plate KH, Risau W, Ullrich A 1994 Glioblastoma growth is inhibited *in vivo* by a negative dominant Flk-1 mutant. *Nature* 367:576-579
200. Millauer B, Longhi MP, Plate KH, Shawver LK, Risau W, Ullrich A 1996 Dominant-negative inhibition of Flk-1 suppresses the growth of many tumor types *in vivo*. *Cancer Res* 56:1615-1620
201. Hillberg F, Wagner EF 1992 Embryonic stem cells lacking functional c-jun. Consequences for growth and differentiation, AP-1 activation and tumorigenesis. *Oncogene* 7:2371-2380
202. Patz A 1980 Studies on retinal neovascularization. *Invest Ophthalmol Vis Sci* 19:1133-1138
203. Olk RJ, Lee CM 1993 Diabetic Retinopathy: Practical Management. Lippincott Co, Philadelphia
204. Michaelson IC 1948 The mode of development of the vascular system of the retina with some observations on its significance for certain retinal disorders. *Trans Ophthalmol Soc UK* 68:137-180
205. Hannehan A, de Juan E, Lutti GA, Fox GM, Schiffer S, Hjelmeland LM 1991 Altered distribution of basic fibroblast growth factor in diabetic retinopathy. *Arch Ophthalmol* 109:1005-1011
206. Meyer-Schwickerath R, Pfeiffer A, Blum WF, Freyberger H, Klein M, Losche C, Rollman R, Schatz H 1993 Vitreous levels of the insulin-like growth factors I and II, and the insulin-like growth factor binding proteins 2 and 3 increase in neovascular disease. *J Clin Invest* 92:2620-2625
207. Aiello LP, Avery R, Arrigg R, Keyt B, Jampel H, Shah S, Pasquale L, Thieme H, Iwamoto M, Park JE, Nguyen H, Aiello LM, Ferrara N, King GL 1994 Vascular endothelial growth factor in ocular fluid of patients with diabetic retinopathy and other retinal disorders. *N Engl J Med* 331:1480-1487
208. Adamis AP, Miller JW, Bernal M-T, D'Amico D, Folkman J, Yeo T-K, Yeo K-T 1994 Increased vascular endothelial growth factor in the vitreous of eyes with proliferative diabetic retinopathy. *Am J Ophthalmol* 118:445-450
209. Malecaze F, Clemens S, Simorer-Pinot V, Mathis A, Chollet P, Favard P, Bayard F, Ploüet J 1994 Detection of vascular endothelial growth factor mRNA and vascular endothelial growth factor-like activity in proliferative diabetic retinopathy. *Arch Ophthalmol* 112:1476-1482
210. Ferrara N 1995 Vascular endothelial growth factor: the trigger for neovascularization in the eye. *Lab Invest* 72:615-618
211. Pe'er J, Shweiki D, Itin A, Hemo I, Gnessin H, Keshet E 1995 Hypoxia-induced expression of vascular endothelial growth factor (VEGF) by retinal cells is a common factor in neovascularization. *Lab Invest* 72:638-645
212. Miller JW, Adamis AP, Shima DT, D'Amore PA, Moulton RS, O'Reilly MS, Folkman J, Dvorak HF, Brown LF, Berse B, Yeo T-K, Yeo K-T 1994 Vascular endothelial growth factor/vascular permeability factor is temporally and spatially correlated with ocular neovascularization in a primate model. *Am J Pathol* 145:574-584
213. Pierce EA, Avery RL, Foley ED, Aiello LP, Smith LE 1995 Vascular endothelial growth factor/vascular permeability factor expression in a mouse model of retinal neovascularization. *Proc Natl Acad Sci USA* 92:905-909
214. Adamis AP, Shima DT, Tolentino M, Gragoudas E, Ferrara N, Folkman J, D'Amore PA, Miller JW 1996 Inhibition of VEGF prevents retinal-ischemia associated iris neovascularization in a primate. *Arch Ophthalmol* 114:66-71
215. Aiello LP, Pierce EA, Foley ED, Takagi H, Riddle L, Chen H, Ferrara N, King GL, Smith LE 1995 Suppression of retinal neovascularization *in vivo* by inhibition of vascular endothelial growth factor (VEGF) using soluble VEGF-receptor chimeric proteins. *Proc Natl Acad Sci USA* 92:10457-10461
216. Bressler NM, Fine SL, Bressler SB 1994 Angiographic interpretation of classic and occult choroidal neovascularization. In: Lewis H, Ryan SJ (eds) Medical and Surgical Retina. Mosby, St. Louis, MO, pp 3-11
217. Lopez PF, Sipky BD, Lambert HM, Thach AB, Hinton DR 1996 Transdifferentiated retinal pigment epithelial cells are immunoreactive for vascular endothelial growth factor in surgically excised age-related macular degeneration-related choroidal neovascular membranes. *Invest Ophthalmol Vis Sci* 37:855-868
218. Kvant A, Algvere PV, Berglin L, Seregard S 1996 Subfoveal fibrovascular membranes in age-related macular degeneration express vascular endothelial growth factor. *Invest Ophthalmol Vis Sci* 37:1929-1934
219. Kvant A 1995 Expression and regulation of vascular endothelial growth factor in choroidal fibroblasts. *Curr Eye Res* 14:1015-1020
220. Koch AE, Harlow L, Haines GK, Amento EP, Unemori EN, Wong W-L, Pope RM, Ferrara N 1994 Vascular endothelial growth factor: a cytokine modulating endothelial function in rheumatoid arthritis. *J Immunol* 152:4149-4155
221. Fava RA, Olsen NJ, Spencer-Green G, Yeo T-K, Yeo K-T, Berse B, Jackman RW, Senger DR, Dvorak HF, Brown LF 1994 Vascular permeability factor/vascular endothelial growth factor (VPF/VEGF): accumulation and expression in human synovial fluids and rheumatoid arthritis. *J Exp Med* 180:340-346
222. Fassbender HJ, Simling-Annefeld M 1983 The potential aggressiveness of synovial tissue in rheumatoid arthritis. *J Pathol* 139:399-406
223. Hamilton J 1976 Hypothesis: *in vitro* evidence for the invasive and tumor-like properties of the rheumatoid pannus. *J Rheumatol* 10:845-851
224. Brown LF, Harris TJ, Yeo KT, Stahle-Backdahl M, Jackman RW, Berse B, Tognazzi K, Dvorak HF, Detmar M 1995 Increased expression of vascular permeability factor (vascular endothelial growth factor) in bullous pemphigoid, dermatitis herpetiformis and erythema multiforme. *J Invest Dermatol* 104:744-749
225. Buttram Jr VC, Betts JW 1979 Endometriosis. *Curr Probl Obstet Gynecol* 11:1-84
226. McLaren J, Prentice A, Charnock-Jones DS, Millican SA, Muller

- KH, Sharkey AM, Smith SK 1996 Vascular endothelial growth factor is produced by peritoneal fluid macrophages in endometriosis and is regulated by ovarian steroids. *J Clin Invest* 98:482–489
227. Shifren JL, Tseng JF, Zaloudek CJ, Ryan IP, Meng YG, Ferrara N, Jaffe RB, Taylor RN 1996 Ovarian steroid regulation of VEGF in human endometrium: implications for angiogenesis during the menstrual cycle and in the pathogenesis of endometriosis. *J Clin Endocrinol Metab* 81:3112–3118
 228. Kamat BR, Brown LF, Manseau EJ, Senger DR, Dvorak HF 1995 Expression of vascular permeability factor/vascular endothelial growth factor by human granulosa and theca lutein cells. Role in corpus luteum development. *Am J Pathol* 146:157–165
 229. Sato K, Yamazaki K, Shizume K, Kanaji Y, Obara T, Oshumi K, Demura H, Yamaguchi S, Shibuya M 1995 Stimulation by thyroid-stimulating hormone and Grave's immunoglobulin G of VEGF mRNA expression in human thyroid follicles *in vitro* and flt mRNA expression in the rat thyroid *in vivo*. *J Clin Invest* 96:1295–1302
 230. Wartofsky L 1994 Diseases of the thyroid. In: Isselbacher KJ, Braunwald E, Wilson JD, Martin JB, Fauci AS, Kasper DL (eds) *Harrison's Principles of Internal Medicine*. McGraw Hill, New York, pp 1930–1953
 231. McClure N, Healy DL, Rogers PA, Sullivan J, Beaton L, Haning Jr RV, Connolly DT, Robertson DM 1994 Vascular endothelial growth factor as capillary permeability agent in ovarian hyperstimulation syndrome. *Lancet* 344:235–269
 232. Baker PN, Krasnow J, Roberts JM, Yeo KT 1995 Elevated serum levels of vascular endothelial growth factor in patients with pre-eclampsia. *Obstet Gynecol* 86:815–821
 233. Thompson RW, D'Amore PA 1990 Recruitment of growth and collateral circulation. In: Zelenock GB, D'Alecy LG, Fantone JC III, Schlafer M, Stanley JC Eds. *Clinical ischemic syndromes: mechanisms and consequences of tissue injury*. CV Mosby, St. Louis, MO, pp 117–134
 234. Graor RA, Gray BH 1991 Interventional treatment of peripheral vascular disease. In: Young JR, Graor RA, Olin JW, Bartholomew JR (eds) *Peripheral Vascular Diseases*. Mosby, St. Louis, MO, pp 111–133
 235. Takeshita S, Weir L, Cheng D, Zheng LP, Riessen R, Bauters C, Symes JF, Ferrara N, Isner JM 1996 Therapeutic angiogenesis following arterial gene transfer of vascular endothelial growth factor in a rabbit model of hindlimb ischemia. *Biochem Biophys Res Commun* 227:628–635
 236. Walder CE, Errett CJ, Ogez J, Heinshon H, Bunting S, Lindquist P, Ferrara N, Thomas GR 1996 Vascular endothelial growth factor improves blood flow and function in a chronic ischemic hind-limb model. *J Cardiovasc Pharmacol* 27:91–98
 237. Bauters C, Asahara T, Zheng LP, Takeshita S, Bunting S, Ferrara N, Symes JF, Isner JM 1994 Physiologic assessment of augmented vascularity induced by VEGF in a rabbit ischemic hindlimb model. *Am J Physiol* 267:H1263–1271
 238. Sellke FW, Kagaya Y, Johnson RG, Shafique T, Schoen FJ, Grossman W, Weintraub RM 1992 Endothelial modulation of porcine coronary microcirculation perfused via immature vessels. *Am J Physiol* 262:H1669–1675
 239. Bauters C, Asahara T, Zheng LP, Takeshita S, Bunting S, Ferrara N, Symes JF, Isner JM 1995 Recovery of disturbed endothelium-dependent flow in collateral-perfused rabbit ischemic hindlimb following administration of VEGF. *Circulation* 91:2793–2801
 240. Essed CD, Brand MVD, Becker AE 1983 Transluminal coronary angioplasty and early restenosis. *Br Heart J* 49:393–402
 241. Asahara T, Bauters C, Pastore C, Bunting S, Ferrara N, Symes JF, Isner JM 1995 Local delivery of vascular endothelial growth factor accelerates re-endothelialization and attenuates intimal hyperplasia in balloon-injured rat carotid artery. *Circulation* 91:2802–2809
 242. Callow AD, Choi ET, Trachtenberg JD, Stevens SL, Connolly DT, Rodi C, Ryan US 1994 Vascular permeability factor accelerates endothelial regrowth following balloon angioplasty. *Growth Factors* 10:223–228
 243. Isner JM, Walsh K, Symes JF, Pieczek A, Takeshita S, Lowry J, Rosenfield K, Weir L, Brogi E, Jurayj D 1996 Arterial gene transfer for therapeutic angiogenesis in patients with peripheral artery disease. *Hum Gene Ther* 7:859–888
 244. Rajewsky K, Gu H, Kühn R, Betz UAK, Müller W, Roes J, Schwenk F 1996 Conditional gene targeting. *J Clin Invest* 3:600–603
 245. Dumont DJ, Gradwohl GJ, Fong GH, Auerbach R, Breitman ML 1993 The endothelial-specific receptor tyrosine kinase, tek, is a member of a new subfamily of receptors. *Oncogene* 8:1293–1301
 246. Sato TN, Tozawa Y, Deutsch U, Wolburg-Buchholz K, Fujiwara Y, Gendron-Maguire M, Gridley T, Wolburg H, Risau W, Qin Y 1995 Distinct roles of the receptor tyrosine kinases Tie-1 and Tie-2 in blood vessel formation. *Nature* 376:70–74
 247. Friedlander M, Brooks PC, Shaffer RW, Kincaid CM, Varner JA, Chersesh DA 1995 Definition of two angiogenic pathways by distinct alpha v integrins. *Science* 270:1500–1502
 248. Mulhauser J, Merrill MJ, Pili R, Maeda H, Bacic M, Bewig B, Passaniti A, Edwards NA, Crystal RG, Capogrossi MC 1995 VEGF₁₆₅ expressed by a replication-deficient recombinant adenovirus vector induces angiogenesis *in vivo*. *Circ Res* 77:1077–1086
 249. Strawn LM, McMahon G, App H, Schreck R, Kuchler WR, Longhi MP, Hui TH, Tang C, Levitzki A, Gazit A, Chen I, Keri G, Orfi L, Risau W, Flamme I, Ullrich A, Hirth KP, Shawver LK 1996 Flk-1 as a target for tumor growth inhibition. *Cancer Res* 56:3540–3545
 250. Robinson GS, Pierce EA, Rook SL, Foley E, Webb R, Smith LE 1996 Oligodeoxynucleotides inhibit retinal neovascularization in a murine model of proliferative retinopathy. *Proc Natl Acad Sci USA* 93:4851–4856

Diabetes Update

Dates: March 20, 1997

Location: Holiday Inn—Union Square, San Francisco

Sponsor: Department of Medicine, University of California, San Francisco

Contact: 415-476-5208, Postgraduate Programs, Box 0656, University of California—San Francisco, San Francisco, CA 94143-0656

Credit: 7.5 hours of Category 1 AMA credit

Fees: \$185/physicians; \$100/allied health professionals (\$430/235 if also registering for “Endocrinology and Metabolism” on March 21–22; see p. 134 for announcement)