

PREMENSTRUAL EXACERBATION OF SYMPTOMS IN MULTIPLE SCLEROSIS IS ATTENUATED BY TREATMENT WITH WEAK ELECTROMAGNETIC FIELDS

REUVEN SANDYK

*NeuroCommunication Research Laboratories, 36 Mill Plain Road, Danbury, CT 06811,
U.S.A. and the Department of Neuroscience at the Institute for Biomedical Engineering
and Rehabilitation Services of Touro College, NY, U.S.A.*

(Received in final form June 22, 1995)

It has been suspected that hormonal factors contribute to the etiology and pathogenesis of multiple sclerosis (MS). A direct relationship between MS and endocrine functions is suggested by changes in disease activity during the phases of the menstrual cycle. A subset of women with MS experience premenstrual worsening of symptoms which improve dramatically with the onset of menstruation. The biological mechanisms underlying these changes in disease activity are unexplained but may be related to cyclical fluctuations in gonadal sex steroid hormones, abrupt changes in the activity of the endogenous opioid peptides and fluctuations in plasma melatonin levels which affect neuronal excitability and immune functions. Extracerebral application of weak electromagnetic fields (EMFs) in the picotesla range intensity has been reported efficacious in the treatment of MS with patients experiencing sustained improvement in motor, sensory, autonomic, affective and cognitive functions. The present report concerns two women with chronic progressive stage MS who experienced, coincident with increasing functional disability, regular worsening of their symptoms beginning about a week before menstruation and abating with the onset of menstruation. These symptoms resolved two months after the initiation of treatment with EMFs. The report supports the association between the endocrine system and MS and indicates that brief, extracranial applications of these magnetic fields modifies the activity of neuroendocrine systems which precipitate worsening of MS symptoms premenstrually.

Keywords: electromagnetic fields; multiple sclerosis; menstrual cycle; premenstrual syndrome; endorphins; pineal gland; melatonin; immune responses

Hormonal factors have been suspected to contribute both directly and indirectly to the etiology and pathogenesis of multiple sclerosis (MS) (Birner, 1945; Tillman, 1950; Sweeney, 1955; Millar et al., 1959; Schapira et al., 1966; Fischman, 1981; Poser & Poser, 1983; Korn-Lubetzki et al., 1984; Birk & Rudick, 1986; Birk et al., 1990; Sandyk, 1993; Sandyk & Awerbuch, 1991; 1994 a; Duquette et al., 1992; Duquette & Girard, 1993). For instance, the susceptibility of MS is higher in women than men and women are even more susceptible when onset occurs at an early or delayed age (Wainerdi, 1961; Fischman, 1981; McCall et al., 1969; Duquette & Girard, 1993). Also, parent-child transmission is clearly more frequent when the mother rather than the father is affected (Duquette et al., 1992). Men, on the other hand, may experience a more rapid progression to a nonambulatory status (Detels et al., 1982). The association between MS and hormonal factors is supported further by the observations that: (a) pregnancy has both a short and long-term favorable effect on the

*Correspondence: Professor Reuven Sandyk, M.D., M.Sc., P.O. Box 203, Bedford Hills, NY 10507, U.S.A.
I am grateful to Lea C. Denn, M.D. for generous assistance in the clinical evaluation of the patients.

course of the disease (Ghezzi & Caputo, 1981; Korn-Lubetzki et al., 1984; Birk & Rudick, 1986; Davis & Maslow, 1992); (b) there is an increased rate of relapse during the postpartum period (Millar et al., 1959; Schapira et al., 1966; Korn-Lubetzki et al., 1984; Birk et al., 1990); (c) the course of MS is influenced by the menstrual cycle (Francis et al., 1990); and (d) there is an association between puberty and the timing of onset of the clinical manifestations MS (Fischman, 1981; Wainerdi, 1961; Sandyk, 1993).

A direct relationship between MS and endocrine functions in women is more clearly illustrated by changes in disease activity during the phases of the menstrual cycle. In my experience a substantial number of women experience exacerbation of MS symptoms premenstrually i.e. during the luteal phase and spontaneous symptomatic improvement coincident with the onset of menses. These patients report premenstrual exacerbation of symptoms starting at an early stage of the disease but with progression and overall decline in motor and mental functioning these exacerbations became more accentuated and debilitating. The occurrence of premenstrual reactivation of symptoms may be related to one or a combination of the following factors: (a) changes in gonadal steroid hormone plasma levels particularly estrogen and progesterone which increase during the luteal phase and then fall abruptly at the onset of menstruation (Junquiera & Carneiro, 1983; Magos & Studd, 1985); (b) changes in the levels of circulating opioid peptides which peak during the luteal phase and fall to undetectable levels at the onset of menstruation (Wehrenberg et al., 1982; Ferin, 1984; Reid, 1983); and (c) changes in melatonin plasma levels which rise during the luteal phase and reach a peak at menstruation (Wetterberg et al., 1976; Hariharasubramanian et al., 1985; Webley & Leidenberger, 1986; Brun et al., 1987; Parry et al., 1990; Sandyk, 1992 a).

Several recent reports have demonstrated that extracranial application of weak electromagnetic fields (EMFs) in the picotesla (pT) range of low frequency is an efficacious, safe, nonpharmacologic modality for symptomatic therapy of MS patients (Sandyk, 1992 b; c 1994 a; b; Sandyk & Derpapas, 1993 a; b; Sandyk & Iacono, 1993; 1994 a; b). These case studies have shown that extracranial application of pT EMFs is efficacious in ameliorating motor, sensory, autonomic, and cognitive functions in patients diagnosed with either relapsing-remitting or chronic progressive MS. This procedure has also been shown efficacious in rapidly resolving an acute relapse of MS (Sandyk & Derpapas, 1993 a). In addition, my experience over the past three years indicates that this treatment modality also stabilizes the course of the disease including in those patients classified with a chronic progressive course. I now report on two patients with MS who experienced regular exacerbations of symptoms premenstrually which were attenuated during the course of treatment with EMFs.

Case Reports

Case I. This 41 year old right-handed woman was well until the age of 19 when she first experienced an attack of left-sided optic neuritis resulting in blindness. Her vision in the left eye returned to normal after several months and she remained clinically asymptomatic until about 8 years ago when, after the birth of her third child, she developed fatigue, paresthesias and numbness in the fingertips of both hands, and weakness of her left leg. Over the ensuing years she experienced gradual progression of symptoms with increased weakness and spasticity in the legs and difficulties with her balance. In 1991 she became dependent on a walker for ambulation. By 1992 she was using ankle braces and in late 1993

she stopped driving a car and by the end of the year became confined to a wheelchair. Approximately six months prior to presentation she began to experience intermittent difficulties with swallowing and increasing weakness in the left arm and hand. Her medical and gynecological history were unremarkable.

The patient recalled that she noted some degree of premenstrual worsening of her symptoms, particularly in fatigue, early in the course of the disease. But as of 1991, when she began ambulating with a walker, she started to experience distinct premenstrual exacerbation of symptoms which included increased weakness and spasticity in the lower extremities, increasing difficulties with swallowing, more frequent episodes of daytime urinary urgency, deterioration of axial balance, difficulties rising from a chair, and generalized weakness with fatigue, mild mental depression, poor concentration and impaired short-term memory functions. On the day of menstruation she experienced abrupt improvement in motor and mental symptoms with a burst of energy, mood elevation and improved ambulation and reported to function physically and mentally best during the 3–4 days of menstruation.

The patient began treatment with EMFs in April of 1994. Using the Sandyk Electromagnetic Stimulator she received an average of 3 treatment sessions per week over the following 2 months and thereafter 1–2 sessions per week during the following 12 months. A session was comprised of two magnetic treatments each of 45 minutes duration separated by a 15 minute break. The frequency of stimulation ranged from 2–4 Hz using a sinusoidal wave in the first treatment and a trapezoidal wave in the second treatment. Since the introduction of treatment with EMFs the patient experienced improvement in a variety of symptoms including increased level of energy, improvement in mood, short-term memory and concentration span, increased strength in the upper limbs especially the left arm and hand, resolution of the dysphagia, improved balance especially trunk control, diminished spasticity in the lower limbs and reduction of nocturia to an average of once per night. The patient also recalled that as early as 2 months after the introduction of magnetic treatment she ceased to experience distinctive episodes of premenstrual reactivation of symptoms noting that her symptoms remained fairly even throughout the menstrual cycle although during some cycles she continued to experience a burst of energy on the day of menstruation.

Case II. This 50 year old right-handed woman was well until the age of 41 when in 1986, after returning from a trip to Florida, she developed an acute attack of optic neuritis on the right and a week later on the left with blurring of vision lasting about 4 weeks. After undergoing a spinal tap she was diagnosed with MS. Four months later, she experienced recurrence of bilateral optic neuritis with visual impairment lasting almost 8 months. She was hospitalized and after receiving a course of intravenous corticosteroids her vision improved immediately and returned to normal 6 weeks later. An MRI scan and visual evoked potential studies which were performed at the same time were abnormal and supportive of the diagnosis of MS. Over the ensuing 5 years she experienced intermittent episodes of urinary urgency and incontinence, difficulties with balance, fatigue, numbness and paresthesias in the feet, dizziness, and weakness in the left leg. Three years prior to presentation she began to use a cane due to increasing weakness in the left leg and additional difficulties with balance. She also experienced paresthesias in the feet, daytime urinary urgency with nocturia 2–3 times, increasing fatigue and heat intolerance, increased sensitivity to light, attacks of daytime sleepiness, episodes of urinary incontinence, constipation, and depression. Over the past year there has been marked deterioration in her balance and after falling sev-

eral times she confined herself to her house. At the time of presentation the patient was taking 4-aminopyridine (45 mg/d) for about 1 1/2 years and fluoxetine 40 mg/d for the past two years.

The patient recalled that about 3 years ago she began to experience regular and predictable exacerbation of symptoms beginning the week prior to menstruation which, at that time, were mild consisting mainly of increased fatigue with intermittent blurring of vision. However, about a year prior to presentation premenstrual exacerbation of symptoms became more intense and disabling and she often felt as though she is having an attack of MS. She began to experience intolerable fatigue forcing her to spend 5–6 hours a day confined to bed. Other symptoms included blurring of vision, increased weakness in the legs, deterioration of balance, marked intolerance to heat with profuse attacks of sweating, increased sensitivity to light, abdominal cramping, increasing constipation and occasional migraneous headaches all of which improved dramatically with the onset of menstruation.

In March of 1995 the patient began experimental treatment with EMFs using the Sandyk Electromagnetic Stimulator which emits an alternating current (AC) EMF of an amplitude of 7.5 picotesla. The patient received 3 magnetic treatment sessions per week. Each treatment session, which was applied in a quiet and artificially illuminated room that was magnetically unshielded, comprised two successive treatments each of 25 minutes duration separated by a break of 15 minutes. The frequency of stimulation used in this patient was 4 Hz for the first AC pulse and 4.5 Hz for the second pulse using sinusoidal and trapezoidal waves, respectively. The patient's eyes were shielded during each treatment. On this regimen the patient made a dramatic recovery during the following four months with improvement in balance, level of energy and mood, resolution of daytime sleepiness, improved strength in the left leg, diminished heat and light sensitivity and diminished daytime and nocturnal urinary frequency. After two months of magnetic therapy the patient ceased to experience worsening of her symptoms premenstrually and reported that her symptoms remained fairly even throughout the cycle. She continued, however, to experience a burst of energy with the onset of menstruation.

DISCUSSION

In the presence of normal menstrual cycles both patients experienced regular exacerbation of their MS symptoms beginning about a week before menstruation and spontaneous improvement in symptoms coincident with the onset of menstruation thus supporting the notion that the course of MS is influenced by fluctuations in gonadal sex steroid hormone levels (Birk & Rudick, 1986; Birk et al., 1990; Francis et al., 1990; Davis & Maslow, 1992; Duquette & Girard, 1993). Both patients, who at the time of presentation were classified as having a chronic progressive MS, noted during the course of the disease worsening of their symptoms premenstrually. However, with progression of the disease and increasing level of disability these exacerbations predictably became more accentuated and debilitating suggesting that in some women premenstrual worsening of symptoms may parallel deterioration of MS disability and may serve as a useful clinical marker of increased disease activity including in patients with chronic progressive MS. In both patients periodic, brief extracerebral applications of EMFs resulted, within two months, in resolution of the premenstrual reactivation of symptoms which occurred along with improvement in motor and mental functions. These findings clearly suggest that in patients with MS application of

EMFs in the picotesla range intensity is capable of attenuating premenstrual reactivation of symptoms conceivably by modifying the activity of neuroendocrine system which ultimately are responsible for exacerbation of the disease premenstrually.

The menstrual cycle reflects the expression of a cyclical process involving the interaction between the hypothalamic-pituitary axis and the ovaries. These interactions, which require integrated neural control mechanisms governed by a hypothalamic "transducer" located within the medial basal hypothalamus (Yen, 1986), are central to the sexual differentiation of the brain, to control of brain-pituitary-gonadal axis, and to the establishment of normal patterns of sexual and aggressive behavior in both sexes (Schipper, 1986). The potent influences exerted by sex steroids on catecholamine and serotonin (5-HT) turnover (Algeri et al., 1977) and the colocalization of labeled estradiol within catecholaminergic luteinizing hormone-releasing hormone (LHRH) positive perikarya (Stumpf & Sar, 1977) suggest that many of the physiologic effects of sex steroids are mediated by modulation of specific monoaminergic and peptidergic pathways. The impact of gonadal sex steroids on monoaminergic and peptidergic neuronal systems may explain the influence of the menstrual cycle on the course of diverse neurological disorders such as migraine, Parkinsonism, chorea, myoclonus, Tourette's syndrome and MS (Schipper, 1986; Quinn & Marsden, 1986; Sandyk et al., 1987; Francis et al., 1990; Davis & Maslow, 1992).

Premenstrual exacerbation of MS symptoms may be considered part of the broader manifestations of the premenstrual syndrome (PMS). PMS, which lately has been defined more rigorously as late luteal-phase dysphoric disorder (American Psychiatric Association, 1987), is a complex psychoneuroendocrine disorder which temporarily disrupts the personal and professional lives of 10%–30% of women throughout their reproductive years (Coppen & Kessel, 1963; Yen, 1986). There's increasing evidence indicating a relationship between PMS and major depressive disorder (Halbreich & Endicott, 1985; Hallman, 1986). PMS patients often have a lifetime history of major depressive disorder (Halbreich & Endicott, 1985) and a subgroup may develop major depressive disorder at a later stage (Schuckit et al., 1975). Features often experienced with PMS include fatigue, irritability, anxiety, overeating, sleep disturbances, disordered temperature regulation, carbohydrate craving, constipation, and lethargy (Dalton, 1964; Reid, 1983). These symptoms are also common in patients with seasonal affective disorder (SAD), a condition characterised by seasonal cyclical mood changes with characteristic symptoms of hypersomnia, fatigue, increase in appetite and weight, and carbohydrate craving occurring in the winter months (Rosenthal et al., 1985; Garvey et al., 1988; Wehr & Rosenthal, 1989).

The pathophysiology of PMS remains unexplained. The popular neuroendocrine hypothesis proposed in 1938 by Israel that PMS results from gonadal steroid hormonal disequilibrium remains controversial (Reid, 1983; Rubinow et al., 1987). Patients with PMS have no abnormalities in secretion or in blood levels of estrogen, progesterone, or prolactin suggesting that the syndrome may involve an abnormal response to physiological levels of sex steroid hormones (Martin & Reichlin, 1987). Nevertheless, menstrual-related fluctuations in sex steroid hormone plasma levels with an abrupt decline in estrogen and progesterone concentrations in the late luteal phase is relevant to the clinical manifestations of MS since estrogens have been shown to influence immune responses (Schuurs & Verheul, 1990; Duquette & Girard, 1993). For instance, estrogen receptors have been demonstrated on macrophages (Gulshan et al., 1990) and estradiol has been shown to increase phagocytic activity in macrophage and the production of interleukin-1 *in vivo* by mouse peritoneal macrophages (Duquette & Girard, 1993). In women, interleukin-1 activity is con-

sistently higher in plasma during the luteal phase (Duquette & Girard, 1993) and natural killer cell activity is decreased during the peri-ovulatory period (Duquette et al., 1992). In addition, estrogens influence the activity of the pineal gland (Semm et al., 1983) and that of 5-HT neurons (Pranzatelli & Snodgrass, 1985), which have been ascribed neuroimmunomodulatory functions (Roszman et al., 1985) and a pathogenetic role in MS (Claveria et al., 1974; Davidson et al., 1977; Sonninen et al., 1973; Tabaddor et al., 1978; Luca & Hategan, 1986; Kissler & Harrer, 1974; Lenon & Carnegie, 1971; Hyypä et al., 1975; Markianos & Sfagos, 1988; Sandyk & Awerbuch, 1991; 1994 *b*).

Reid and Yen (1981; 1983) proposed that cyclical gonadal steroid-induced changes in endogenous opioid peptide activity might be the central neuroendocrine event that precipitates the varied manifestations of PMS. As levels of circulating opioid peptides in primates peak during the luteal phase and subsequently fall to undetectable levels at the onset of menstruation (Wehrenberg et al., 1982; Reid, 1983; Ferin, 1984), it was suggested that opioid withdrawal premenstrually, which may lead to rebound hyperactivity of noradrenergic neural pathways owing to slowly acquired receptor supersensitivity (Walker et al., 1984), might precipitate in susceptible individuals the development of the mental changes of PMS (Reid, 1983; Halbreich et al., 1988). This hypothesis is supported by the observation that in humans opiate withdrawal is associated with depression, irritability, a low threshold for stressful stimuli, altered appetite and disturbances of temperature regulation (Halbreich et al., 1986).

Changes in the activity of endogenous opioid peptides have been demonstrated in patients with MS (Rosenberg & Appenzeller, 1988; Gusev et al., 1994), but it remains unknown whether these are related causally to the pathogenesis and clinical manifestations of the disease. Circumstantial evidence suggests that the endogenous opioids are involved in the pathophysiology of MS since: (a) stress, which causes the release of beta-endorphins (Cohen et al., 1982), adversely affects the symptoms and course of the disease (Walton, 1977; Warren et al., 1982; Grant et al., 1989), (b) chronic pain, which is associated with disruption of opioid peptide activity (Terentus, 1982), is a common feature of MS occurring in a range 20% to 80% of patients at some point during the course of the disease (Albert, 1969; Shibasaki & Kuroiwa, 1974; Clifford & Trotter, 1984) (c) MS patients who experienced diminished fatigability during treatment with amantadine (Symmetrel®) were found to have higher plasma beta-endorphin levels compared to patients whose fatigue was unaffected by the drug (Rosenberg & Appenzeller, 1988); (d) beta endorphins are known to exert a neuroimmunomodulatory function and therefore may be involved in some pathogenetic aspects of the disease (Weber & Pert, 1984; Gusev et al., 1994); and (e) opioid peptides and opiates influence the propagation of action potential discharges of single neurons by increasing potassium conductance (Williams et al., 1982) which is implicated in impulse transmission in demyelinated fibers (van Diemen et al., 1993; Russell et al., 1995). Thus, the rise in opioid peptide activity during the luteal phase and its rapid decline with the onset of menstruation might be involved in the pathogenesis of premenstrual reactivation of MS symptoms.

PMS also has been proposed to be related causally to changes in pineal melatonin secretion (Clare, 1985; Parry et al., 1987, 1990; Sandyk, 1992 *a*). Plasma melatonin levels are low during ovulation and high premenstrually (Wetterberg et al., 1976; Arendt, 1978; Penny, 1982; Law, 1986; Webley & Leidenberger, 1986; Hariharasubramanian et al., 1985; Brun et al., 1987). Wetterberg et al. (1976) found plasma melatonin levels during men-

stration to be 45 times higher than during ovulation. In addition, altered circadian nocturnal melatonin secretion was reported in women with PMS and evening administration of bright light, which inhibits melatonin secretion (Lewy et al., 1980) and resets its circadian secretion, has shown beneficial effects in patients with premenstrual depression (Parry et al., 1987, 1989). On the other hand, melatonin may cause worsening of dysphoric mood in depressed patients and exacerbation of symptoms in MS patients (Carman et al., 1976; Sandyk, 1992 *b*). Since nocturnal melatonin plasma levels are diminished in MS patients during periods of symptom exacerbation and in those with a history of mental depression (Sandyk & Awerbuch, 1992; Sandyk & Awerbuch, 1993 *a, b*; and since melatonin is a neuroimmunomodulator (Maestroni et al., 1987 *a; b*; Cardarelli, 1990; Pierpaoli & Maestroni, 1987, 1988; Fraschin & Reiter 1990; Guerrero & Reiter, 1992), it is conceivable that abnormal circadian melatonin secretion premenstrually may contribute to the development of premenstrual reactivation of symptoms in these patients.

The mechanisms by which treatment with EMFs attenuate premenstrual exacerbation of symptoms in these patients remain unexplained. The neuroendocrine system, particularly the hypothalamus and pineal gland, are sensitive to the influences of the geomagnetic field and to exposure to artificial EMFs (Jakovleva, 1975; Kholodov, 1966; Semm, 1983; Semm et al., 1980; Free et al., 1981; Stoupel et al., 1983, Michaelson & Lu, 1992; Wilson et al., 1992). For instance, in rats chronic exposure to electric fields has been reported to produce changes in corticosteroids, testosterone, and growth hormone levels (Hackman & Graves, 1981; Free et al., 1981) which in part involve the mediation of the endogenous opioid peptides and melatonin (Kavaliers & Ossenkopp, 1992). That EMFs alter opioid functions is suggested by observations in experimental animal studies which have shown magnetic stimuli to: (a) suppress the analgesic and locomotor effects of morphine (Kavaliers et al., 1984; Kavaliers & Ossenkopp, 1986 *a; b*), (b) modify the rate of tolerance development to morphine-induced analgesia (Kavaliers & Ossenkopp, 1985), and (c) reduce stress-induced opioid analgesia (Kavaliers & Ossenkopp, 1986 *c*). Magnetic stimuli also have been shown to alter the circadian secretion of pineal melatonin (Semm et al., 1980; Welker et al., 1983; Rudolph et al., 1988) which is altered in women with premenstrual depression (Parry et al., 1990). Melatonin in turn, has been shown to modify immune responses (Jankovic et al., 1970; Rella & Lapin, 1976; Maestroni et al., 1987 *a; b*; Pierpaoli & Maestroni, 1987, 1988; Angeli et al., 1988; Maestroni et al., 1988; Guerrero & Reiter, 1992), inhibit the activity of endogenous opioid peptides (Lissoni et al., 1986), affect the release of gonadotropin (Reiter, 1991), and increase 5-HT functions (Anton-Tay et al., 1968; Moszkowska et al., 1971; Aldegunde et al., 1985) which are thought to participate in the pathogenesis of PMS symptoms (Halbreich et al., 1988; Pies, 1990; Rickels et al., 1990; Stone et al., 1991).

In conclusion I have presented two women with MS who experienced regular premenstrual reactivation of symptoms which became increasingly more disabling in temporal association with overall deterioration in disability. Since all ovulating women have cyclical hormonal fluctuations, it remains unclear why only a subset of women develop PMS and with respect to MS patient, why some experience premenstrual reactivation of the disease followed by spontaneous improvement of symptoms with the onset of menstruation. Nevertheless, the observation that periodic applications EMFs led to attenuation of premenstrual exacerbation of symptoms in these patients adds an additional facet in support of the unique efficacy of this treatment modality in the management of MS.

REFERENCES

- Albert, M. L. (1969). Treatment of pain multiple sclerosis. *New England Journal of Medicine*, 280, 395.
- Aldegunde, M., Miguez, I. and Veira, J. (1985). Effects of pinealectomy on regional brain serotonin metabolism. *International Journal of Neuroscience*, 26, 9–13.
- Algeri, S., Bonati, M. and Curcio, M. (1977). Biochemical effects of steroid contraceptive drugs on some neurotransmitters in the central nervous system. In Garattini, S. and Berendes, H. W. (Eds.), *Pharmacology of steroid contraceptive drugs*. New York: Raven Press.
- American Psychiatric Association (1987). *Diagnostic and statistical manual of mental disorders*. 3rd edition. Washington, DC.
- Angeli, A., Gatti, G., Sartori, M. L., DelPonte, D. and Garagnola, R. (1988). Effect of exogenous melatonin on human natural killer (NK) cell activity: an approach to the immunomodulatory role of the pineal gland. In D. Gupta, A. Attanasio and R. J. Reiter (Eds). *The pineal gland and cancer*. London Brain Research Promotion.
- Anton-Tay, F., Chou, C., Anton, S., and Wurtman R. J. (1968). Brain serotonin concentration elevation following intraperitoneal administration of melatonin. *Science*, 162, 277–278.
- Arendt, J. (1978). Melatonin assays in body fluids. *Journal of Neural Transmission*, 13, (suppl), 265–278.
- Birk, K. and Rudick, R. (1986). Pregnancy and multiple sclerosis. *Archives of Neurology*, 43, 738–742.
- Birk, K., Ford, C., Smeltzer, S., Ryan, D., Miller, R. and Rudick, R. A. (1990). The clinical course of multiple sclerosis during pregnancy and the puerperium. *Archives of Neurology*, 47, 738–742.
- Birner, I. M. (1945). Pregnancy and multiple sclerosis. Case report *New York State Journal of Medicine*, 45, 634–635.
- Brun, J., Clausturat, B. and David, M. (1987). Urinary melatonin, LH, oestradiol, progesterone excretion during the menstrual cycle or in women taking oral contraceptive. *Acta Endocrinologica* (Copenh.), 116, 145–149.
- Cardarelli, N. F. (1990). The role of a thymus-pineal axis in an immune mechanism of aging. *Journal of Theoretical Biology*, 45, 397–405.
- Carman, J. S., Post, R. M., Buswell, R. and Goodwin, F. K. (1976). Negative effects of melatonin on depression. *American Journal of Psychiatry*, 133, 1181–1186.
- Clare, A. W. (1985). Hormones, behavior and the menstrual cycle. *Journal of Psychosomatic Research*, 29, 225–232.
- Claveria, L. E., Curzon, G., Harrison, M. J. G., and Kantamareni, B. D. (1974). Amine metabolites in the cerebrospinal fluid of patients with disseminated sclerosis. *Journal of Neurology, Neurosurgery and Psychiatry*, 37, 715–718.
- Clifford, D. B. and Trotter, J. L. (1984). Pain in multiple sclerosis. *Archives of Neurology*, 41, 1270–1272.
- Cohen, M. R., Pickar, D., Dubois, M., Nurenberger, J., Roth, Y., Cohen, R. M., Gershon, E. and Bunney, W. E. (1982). Clinical and experimental studies of stress and the opioid system. *Annals of the New York Academy of Sciences*, 398, 711–719.
- Coppen, A. and Kessel, N. (1963). Menstruation and personality. *British Journal of Psychiatry*, 109, 711–719.
- Dalton, K., (1964). *The premenstrual syndrome*. Springfield, IL: Charles C. Thomas.
- Davidson, D., Pullar, I. A., Mawdsley, C., Kinloch, N. and Yates, M. (1977). Monoamine metabolites in cerebrospinal fluid in multiple sclerosis. *Journal of Neurology, Neurosurgery and Psychiatry*, 40, 741–745.
- Davis, R. K. and Maslow, A. S. (1992). Multiple sclerosis in pregnancy: a review. *Obstetrics and Gynecological Survey*, 47, 290–296.
- Detels, R., Clark, V. A., Valdiviezo, N. L., Visscher, B. R., Malmgren, R. M. and Dudley, J. P. (1982). Factors associated with a rapid course of multiple sclerosis. *Archives of Neurology*, 39, 337–341.
- Duquette, P. and Girard, M. (1993). Hormonal factors in susceptibility to multiple sclerosis. *Current Opinion in Neurology and Neurosurgery*, 6, 195–201.
- Duquette, P., Pleines, J., Charest, L., Senecal-Quevillon, M. and Masse, C. (1992). The increased susceptibility of women to multiple sclerosis. *Canadian Journal of Neurological Sciences*, 19, 466–471.
- Ferin, M. (1984). Endogenous opioid peptides and the menstrual cycle. *Trends in Neurosciences*, 8, 194–196.
- Fischman, H. R. (1981). Multiple sclerosis: a two stage process? *American Journal of Epidemiology*, 114, 244–252.
- Francis, G. S., Antel, J. P. and Duquette, P. (1990). Inflammatory demyelinating diseases of the CNS in W. G. Bradley, B. Daroff, G. M. Fenichel and C. D. Marsden (Eds.), *Neurology in clinical practice* (p. 1154). Boston: Butterworth.
- Fraschini, F. and Reiter, R. J. (1990). *Role of melatonin and pineal peptides in neuroimmunomodulation* (p. 329). New York: Plenum.
- Free, M. J., Kaune, W. T., Phillips, R. D. and Chen, H. C. (1981). Endocrinological effects of strong 60-Hz electric fields on rats. *Bioelectromagnetics*, 2, 105–122.
- Garvey, M. J., Wesner, R. and Godes, M. (1988). Comparison of seasonal and nonseasonal affective disorder. *American Journal of Psychiatry*, 145, 100–102.
- Ghezzi, A. and Caputo, D. (1981). Pregnancy: a factor influencing the course of multiple sclerosis? *European Neurology*, 20, 115–117.

- Grant, I., Brown, G. and Harris, T. (1989). Severely threatening events and marked life difficulties preceding onset and exacerbation of MS. *Journal of Neurology Neurosurgery and Psychiatry*, 52, 8–13.
- Guerrero, J. M. and Reiter, R. J. (1992). A brief survey of pineal gland-immune system interrelationships. *Endocrine Research*, 18, 91–113.
- Gulshan, S., McCruden, A. B. & Stimson, W. H. (1990). Estrogen receptors in macrophages. *Scandinavian Journal of Immunology*, 31, 691–697.
- Gusev, E. I., Demina, T. L., Boiko, A. N. and Pinegin, B. V. (1994). Prolonged dynamic clinico-immunological observation of 85 patients with definite multiple sclerosis: first steps towards monitoring process activity. *Journal of Neurology*, 24, 500–510.
- Hackman, R. M. and Graves, H. B. (1981). Corticosterone levels in mice exposed to high intensity electric fields. *Behavioral Neural Biology*, 32, 201–213.
- Halbreich, U. and Endicott, J. (1985). Relationship of dysphoric premenstrual changes to depressive disorders. *Acta Psychiatrica Scandinavica*, 71, 331–338.
- Halbreich, U., Holtz, I. and Paul, L. (1986). Premenstrual changes. Impaired hormonal homeostasis. *Neurologic Clinics*, 6, 173–194.
- Hallman, J. (1986). The premenstrual syndrome: an equivalent of depression? *Acta Psychiatrica Scandinavica*, 73, 403–411.
- Hariharasubramanian, N., Nair, N. P. V. and Pilapil, C. (1985). Circadian rhythm of plasma melatonin and cortisol during the menstrual cycle. In G. M. Brown and S. D. Wainwright (Eds.), *The pineal gland endocrine aspects* (pp. 31–36). Oxford: Pergamon Press.
- Hyypä, M. T., Joima, T., Riekkinen, P. and Rinne, U. K. (1975). Effects of L-tryptophan on central indoleamine metabolism and short-lasting neurologic disturbances in multiple sclerosis. *Journal of Neural Transmission*, 37, 297–304.
- Israel, S. I. (1938). Premenstrual tension. *Journal of the American Medical Association*, 110, 1721–1726.
- Jakovleva, M. I. (1975). *The physiological mechanisms of electromagnetic fields*. Medicine, Leningrad.
- Jankovic, B. D., Isakovic, K. and Petrovic, S. (1970). Effect of pinealectomy on immune reactions in the rat. *Immunology*, 18, 1–5.
- Junquiera, L. C. and Carneiro, J. (1983). *Basic histology*, 4th ed. Los Altos, CA: Lange Medical Publications.
- Kavaliers, M. and Ossenkopp, K. P. (1985). Tolerance to morphine-induced analgesia in mice: magnetic fields function as environmental specific cues and reduce tolerance development. *Life Sciences*, 37, 1125–1135.
- Kavaliers, M. and Ossenkopp, K. P. (1986 a). Stress-induced opioid analgesia and activity in mice: inhibitory influences of exposure to magnetic fields. *Psychopharmacology*, 89, 440–443.
- Kavaliers, M. and Ossenkopp, K. P. (1986 b). Magnetic fields differentially inhibit mu, delta, kappa and sigma opiate-induced analgesia in mice. *Peptides*, 7, 449–453.
- Kavaliers, M. and Ossenkopp, K. P. (1986 c). Magnetic field inhibition of morphine-induced analgesia and behavioral activity in mice: evidence for involvement of calcium ions. *Brain Research*, 379, 30–38.
- Kavaliers, M. and Ossenkopp, K. P. (1992). Magnetic fields, opioid systems, and day-night rhythms of behavior. In Moore-Ede, M. C., Campbell, S. S. and Reiter, R. J. (Eds.), *Electromagnetic fields and circadian rhythmicity* (pp. 95–117). Boston: Birkhauser.
- Kavaliers, M., Ossenkopp, K. P. and Hirst, M. (1984). Magnetic fields abolish the enhanced nocturnal analgesic response to morphine in mice. *Physiology & Behavior*, 32, 261–264.
- Kholodov, I. A. (1966). The influence of electromagnetic and magnetic fields on the nervous system. *Joint research service: IPRS-37102, TT-66-33531*. Washington.
- Kisser, W. and Harrer, G. (1974). Amino acid determinations in serum of patients with multiple sclerosis. *Wiener Klinische Wochenschrift*, 86, 80–82.
- Korn-Lubetzki, I., Kahana, E., Cooper, G. and Abramsky, O. (1984). Activity of multiple sclerosis during pregnancy and puerperium. *Annals of Neurology*, 16, 229–231.
- Law, S. (1986). The regulation of the menstrual cycle and its relationship to the moon. *Acta Obstetrica Gynaecologica Scandinavica*, 65, 45–49.
- Lennon, V. A. and Carnegie, P. R. (197). Immunopharmacological disease: a break in tolerance to receptor sites. *Lancet*, 2, 630–633.
- Lewy, A. J., Wehr, T. A. and Goodwin, F. K. (1980). Light suppresses melatonin secretion in humans. *Science*, 210, 1267–1269.
- Lissoni, P., Esposti, D., Esposti, G., Mauri, R., Resentini, M., Morabito, F., Fumagali, P., Santagostino, A., Delitala, G. and Fraschini, A. (1986). A clinical study on the relationship between the pineal gland and the opioid system. *Journal of Neural Transmission*, 65, 63–73.
- Luca, N. and Hategan, D. (1986). Determination of serotonin content and ceruloplasmin activity of blood and CSF amino acid level in multiple sclerosis. *Reveu Roumania Medica Neurologica Psychiatrica*, 24, 153–159.
- Maestroni, G. J. M., Conti, A. and Pierpaoli, W. (1987 a). The pineal gland and the circadian, opiateergic, immunoregulatory role of melatonin. *Annals of the New York Academy of Science*, 496, 67–77.
- Maestroni, G. J. M., Conti, A. and Pierpaoli, W. (1987 b). Melatonin counteracts the effect of acute stress on the

- immune system via an opiate system. In G. P. Trentini, C. De Gaetani and P. Pevet (Eds.), *Fundamentals and clinics in pineal research* (pp. 277–279). New York: Raven Press.
- Maestroni, G. J. M., Conti, A. and Pierpaoli, W. (1988). Role of the pineal gland in immunity. III: Melatonin antagonizes the immunosuppressive effect of acute stress via an opiate mechanism. *Immunology*, 63, 465–469.
- Magos, A. & Studd, J. (1985). Effects of the menstrual cycle on medical disorders. *British Journal of Hospital Medicine*, 33, 68–77.
- Markianos, M. & Sfagos, C. (1988). Altered serotonin uptake kinetics in multiple sclerosis. *Journal of Neurology*, 235, 236–237.
- Martin, J. B. and Reichlin, S. (1987). *Clinical neuroendocrinology* (pp. 639–655). Philadelphia: F. A. Davis.
- McCall, M., Sutherland, J. and Acheson, E. (1969). The frequency of multiple sclerosis in Western Australia. *Acta Neurologica Scandinavica*, 45, 151–165.
- Michaelson, S. M. and Lu, S. T. (1992). Factors in postmeridian hormone changes among rats in electric field exposure studies. In M. C. Moore-Ede, S. S. Campbell and R. J. Reiter (Eds.), *Electromagnetic fields and circadian rhythmicity* (pp. 77–94). Boston: Birkhauser.
- Millar, J. H. D., Allison, R. S., Cheeseman, E. A. and Merrett, J. D. (1959). Pregnancy as a factor influencing relapse in disseminated sclerosis. *Brain*, 82, 417–426.
- Moszkowska, A., Kordon, C. and Ebels, I. (1971). Biochemical fractions and mechanisms involved in the pineal modulation of pituitary gonadotropin release. In G. E. W. Wolstenholme & J. Knight (Eds.), *The pineal gland* (pp. 241–258). London: Churchill-Livingstone.
- Parry, B. L., Rosenthal, N., Tamarin, L. and Wehr, T. (1987). Treatment of a patient with season premenstrual syndrome. *American Journal of Psychiatry*, 144, 762–766.
- Parry, B. L., Berga, S. L., Mostofi, N., Sependa, P. A., Kripke, D. F. and Gillin, J. C. (1989). Morning versus evening bright light treatment of late luteal phase dysphoric disorder. *American Journal of Psychiatry*, 146, 1215–1217.
- Parry, B. L., Berga, S. L., Kripke, D. F., Klauber, M. R., Laughlin, G. A., Yen, S. S. C. and Gillin, J. C. (1990). Altered wave form of plasma nocturnal melatonin secretion in premenstrual depression. *Archives of General Psychiatry*, 47, 1139–1146.
- Penny, R. (1982). Melatonin excretion in normal males and females: increase during pregnancy. *Metabolism*, 31, 816–822.
- Pierpaoli, W. and Maestroni, G. J. M. (1987). Melatonin: a principal neuroimmunoregulatory and anti-stress hormone: its anti-aging effects. *Immunology Letters*, 16, 355–362.
- Pierpaoli, W. and Maestroni, G. J. M. (1988). Neuroimmunomodulation: some recent views and findings. *International Journal of Neuroscience*, 39, 25–33.
- Pies, R. W. (1990). Fluoxetine treatment of premenstrual syndrome. *Journal of Clinical Psychiatry*, 51, 348.
- Poser, S. & Poser, W. (1983). Multiple sclerosis and gestation. *Neurology*, 33, 1422–1427.
- Pranzatelli, M. and Snodgrass, S. (1985). The pharmacology of myoclonus. *Clinical Neuropharmacology*, 8, 99–130.
- Quinn, N. P. and Marsden, C. D. (1986). Menstrual-related fluctuations in Parkinson's disease. *Movement Disorders*, 1, 85–87.
- Reid, R. L. (1983). Endogenous opioid activity and the premenstrual syndrome. *Lancet*, 1, 786.
- Reid, R. L. and Yen, S. S. C. (1981). Premenstrual syndrome. *American Journal of Obstetrics and Gynecology*, 139, 85–104.
- Reid, R. L. and Yen, S. S. C. (1983). The premenstrual syndrome. *Clinical Obstetrics and Gynecology*, 26, 709–716.
- Reiter, R. J. (1991). Pineal gland. Interface between the photoperiodic environment and the endocrine system. *Trends in Endocrinology and Metabolism*, 2, 13–19.
- Rella, W. and Lapin, V. (1976). Immunocompetence of pinealectomized and simultaneously pinealectomized and thymectomized rats. *Oncology*, 33, 3–6.
- Rickels, K., Freeman, E. W., Sondheim, S. and Albert, J. (1990). Fluoxetine in the treatment of premenstrual syndrome. *Current Therapeutic Research*, 48, 161–166.
- Rosenberg, G. A. and Appenzeller, O. (1988). Amantadine, fatigue, and multiple sclerosis. *Archives of Neurology*, 45, 1104–1106.
- Rosenthal, N. E., Sack, D. A., Carpenter, C. J., Parry, B., Mendelson, W. B. and Wehr, T. A. (1985). Antidepressant effects of light in seasonal affective disorder. *American Journal of Psychiatry*, 142, 163–170.
- Roszman, T. L., Jackson, J. C., Cross, R. J., Titus, M. J., Markesbery, W. R. and Brooks, W. H. (1985). Neuroanatomic and neurotransmitter influences on immune function. *The Journal of Immunology*, 135, 769S–772S.
- Rubinow, D. R., Hoban, M. C. and Grover, G. N. (1987). Menstrually-related mood disorders. In Nerozzi, D., Goodwin, F. K. and Costa, E. (Eds.), *Hypothalamic dysfunction in neuropsychiatric disorders* (pp. 335–346). New York: Raven Press.
- Rudolph, K., Wirz-Justice, A., Krauchi, K. and Feer, H. (1988). Static magnetic fields decrease nocturnal pineal cAMP in the rat. *Brain Research*, 446, 159–160.

- Russell, J. W., Windebank, A. J. and Harper, C. M. (1995). Treatment of stable chronic demyelinating polyneuropathy with 3, 4-diaminopyridine. *Mayo Clinic Proceedings*, 70, 532–539.
- Sandyk, R. (1992 a). The Pineal gland and the menstrual cycle. *International Journal of Neuroscience*, 63, 197–204.
- Sandyk, R. (1992 b). Successful treatment of multiple sclerosis with magnetic fields. *International Journal of Neuroscience*, 66, 237–250.
- Sandyk, R. (1992 c). The pineal gland and multiple sclerosis. (Editorial). *International Journal of Neuroscience*, 63, 206–215.
- Sandyk, R. (1993). Multiple sclerosis: the role of puberty and the pineal gland in its pathogenesis. *International Journal of Neuroscience*, 68, 209–225.
- Sandyk, R. (1994 a). Further observations on the effects of external picotesla range magnetic fields on visual memory and visuospatial functions in multiple sclerosis. *International Journal of Neuroscience*, 77, 203–227.
- Sandyk, R. (1994 b). Rapid normalization of visual evoked potentials by picotesla range magnetic fields in chronic progressive multiple sclerosis. *International Journal of Neuroscience*, 77, 243–259.
- Sandyk, R. and Awerbuch, G. I. (1991). The pineal gland in multiple sclerosis. *International Journal of Neuroscience*, 61, 61–67.
- Sandyk, R. and Awerbuch, G. I. (1992). Nocturnal plasma melatonin and alpha-melanocyte stimulating hormone levels during exacerbation of multiple sclerosis. *International Journal of Neuroscience*, 67, 173–186.
- Sandyk, R. and Awerbuch, G. I. (1993 a). Nocturnal melatonin secretion in multiple sclerosis patients with affective disorders. *International Journal of Neuroscience*, 68, 227–240.
- Sandyk, R. and Awerbuch, G. I. (1993 b). Nocturnal melatonin secretion in suicidal patients with multiple sclerosis. *International Journal of Neuroscience*, 71, 173–182.
- Sandyk, R. and Awerbuch, G. I. (1994 a). Relationship of nocturnal melatonin levels to duration and course of multiple sclerosis. *International Journal of Neuroscience*, 75, 229–237.
- Sandyk, R. and Awerbuch, G. I. (1994 b). The co-occurrence of multiple sclerosis and migraine headaches: the serotonergic link. *International Journal of Neuroscience*, 76, 249–257.
- Sandyk, R. and Derpapas, K. (1993 a). Successful treatment of an acute exacerbation of multiple sclerosis by external magnetic fields. *International Journal of Neuroscience*, 70, 97–105.
- Sandyk, R. and Derpapas, K. (1993 b). Magnetic fields normalize visual evoked potentials and brainstem auditory evoked potentials in multiple sclerosis. *International Journal of Neuroscience*, 68, 241–253.
- Sandyk, R. and Iacono, R. P. (1993). Resolution of longstanding symptoms of multiple sclerosis by application of picotesla range magnetic fields. *International Journal of Neuroscience*, 70, 255–269.
- Sandyk, R. and Iacono, R. P. (1994 a). Multiple sclerosis: Improvement of visuo-perceptive functions by picotesla range magnetic fields. *International Journal of Neuroscience*, 74, 177–189.
- Sandyk, R. and Iacono, R. P. (1994 b). Improvement by picotesla range magnetic fields of perceptual-motor performance and visual memory in a patient with chronic progressive multiple sclerosis. *International Journal of Neuroscience*, 78, 53–66.
- Sandyk, R., Bamford, C. R. and Iacono, R. P. (1987). Menstrual-related fluctuations in Tourette's syndrome. *International Journal of Neuroscience*, 35, 237–239.
- Schapira, K., Poskanzer, D. C., Newell, D. J. and Miller, H. (1966). Marriage, pregnancy and multiple sclerosis. *Brain*, 89, 419–428.
- Schipper, H. M. (1986). Neurology of sex steroids and oral contraceptives. *Neurologic Clinics*, 4, 721–751.
- Schuckit, M. A., Daly, V., Herman, G. and Hineman, S. (1975). Premenstrual symptoms and depression in a university population. *Diseases of the Nervous System*, 36, 516–517.
- Schuurs, A. H. and Verheul, H. A. (1990). Effects of gender and sex steroids on the immune response. *Journal of Steroid Biochemistry*, 35, 157–172.
- Semm, P. (1983). Neurobiological investigations on the magnetic sensitivity of the pineal gland in rodents and pigeons. *Comparative Biochemistry and Physiology*, 76A, 683–689.
- Semm, P., Schneider, T. and Vollrath, L. (1980). Effects of an earth-strength magnetic field on electrical activity of pineal cells. *Nature*, 288, 607–608.
- Semm, P., Demaine, C. and Vollrath, L. (1981). The effects of sex hormones, prolactin, and chorionic gonadotropin on pineal electrical activity in guinea pigs. *Cellular and Molecular Neurobiology*, 1, 259–269.
- Shibasaki, H. and Kuroiwa, Y. (1974). Painful tonic seizures in multiple sclerosis. *Archives of Neurology*, 30, 47–51.
- Sonninen, V., Riekkinen, P. and Rinne, U. K. (1973). Acid monoamine metabolites in cerebrospinal fluid in multiple sclerosis. *Neurology*, 23, 760–763.
- Stone, A. B., Pearlstein, T. B. and Brown, W. A. (1991). Fluoxetine in the treatment of late luteal phase dysphoric disorder. *Journal of Clinical Psychiatry*, 52, 290–293.
- Stoupe, E., Keret, R., Assa, S., Kaufman, H., Shimshoni, M. and Laron, Z. (1983). Secretion of growth hormone, prolactin and corticosteroids during different levels of geomagnetic activity. *Neuroendocrinology Letters*, 5, 365–368.

- Stumpf, W. E. and Sar, M. (1977). Steroid hormone target cells in the periventricular brain: relationship to peptide hormone producing cells. *Federation Proceedings*, 36, 1973–1977.
- Sweeney, W. J. (1955). Pregnancy and multiple sclerosis. *American Journal of Obstetrics and Gynecology*, 66, 124–130.
- Tabaddor, K., Wolfson, L. I. and Sharpless, N. S. (1978). Ventricular fluid homovanillic acid and 5-hydroxyindoleacetic acid concentrations in patients with movement disorders. *Neurology*, 28, 1249–1253.
- Terenius, L. (1982). Clinical aspects. In J. B. Malick & R. M. S. Bell (Eds.), *Endorphins* (pp. 229–255). New York: Marcel Dekker.
- Tillman, A. J. B. (1950). The effect of pregnancy on multiple sclerosis and its management. *Research Publications of the Association for Research in Nervous and Mental Disease*, 28, 548–582.
- Van Diemen, H. A. M., Polman, C. H., van Dongen, M. M. M., Nauta, J. J. P., Strijers, R. L. M., van Loenen, A. C., Bertelsmann, F. W. and Koetsier, J. C. (1993). 4-aminopyridine induces functional improvement in multiple sclerosis patients: a neuropsychological study. *Journal of the Neurological Sciences*, 116, 220–226.
- Wainerdi, H. R. (1961). Does the multiple-sclerosis syndrome begin at puberty? *Boston Medical Quarterly*, 12, 44–47.
- Walker, J. M., Khachaturian, H. and Watson, S. J. (1984). Some anatomical and physiological interactions among noradrenergic systems and opioid peptides. *Frontiers in Clinical Neurosciences*, 2, 74–82.
- Walton, J. N. (1977). *Brain's diseases of the nervous system* (p. 548). Oxford: Oxford University Press.
- Warren, S., Greenhill, S. and Warren, K. (1982). Emotional stress and the development of MS: case-control evidence of a relationship. *Journal of Chronic Disease*, 35, 821–831.
- Weber, R. J. and Pert, C. B. (1984). Opiatergic modulation of the immune system. In E.E. Miller & A.R. Genazzani (Eds.), *Central and peripheral endorphins: basic and clinical aspects* (pp. 35–42). New York: Raven Press.
- Webley, G. E. & Leidenberger, F. (1986). The circadian pattern of melatonin and its positive relationship with progesterone in women. *Journal of Clinical Endocrinology*, 63, 323–328.
- Wehr, T. A. and Rosenthal, N. E. (1989). Seasonality and affective illness. *American Journal of Psychiatry*, 146, 829–839.
- Wehrenberg, W. B., Wardlaw, S. L., Frantz, A. G. and Ferin, M. (1982). Beta-endorphin in hypophyseal portal blood variations throughout the menstrual cycle. *Endocrinology*, 111, 879–881.
- Welker, H. A., Semm, P., Willig, R. P., Commentz, J. C., Wiltshko, W. and Vollrath, L. (1983). Effects of an artificial magnetic field on serotonin N-acetyltransferase activity and melatonin content of the rat pineal gland. *Experimental Brain Research*, 50, 426–432.
- Wetterberg, L., Arendt, J., Paunier, L., Sizonenko, P. C., Von Donselaer, W. and Heyden, T. (1976). Human serum melatonin changes during the menstrual cycle. *Journal of Clinical Endocrinology and Metabolism*, 42, 185–188.
- Wilson, B. W., Stevens, R. G. and Anderson, L. E. (1992). Effects of electromagnetic field exposure on neuroendocrine function. In M.C. Moore-Ede, S.S. Campbell & R.J. Reiter (Eds.), *Electromagnetic fields and circadian rhythmicity* (pp. 29–50). Boston: Birkhauser.
- Williams, J. T., Egan, T. M. and North, R. A. (1982). Enkephalin opens potassium channels on mammalian central neurones. *Nature*, 299, 74–77.
- Yen, S. S. C. (1986). The human menstrual cycle. In S. S. C. Yen and R. B. Jaffe (Eds.), *Reproductive endocrinology, physiology, pathophysiology and clinical management* (pp. 200–236). Philadelphia: W.B. Saunders.