



Serotonin-induced Decrease in Brain ATP, Stimulation of Brain Anaerobic Glycolysis and Elevation of Plasma Hemoglobin; the Protective Action of Calmodulin Antagonists

NOA KOREN-SCHWARTZER, MALCA CHEN-ZION,
HANNA BEN-PORAT and RIVKA BEITNER*

Health Sciences Research Center, Department of Life Sciences, Bar-Ilan University, Ramat Gan 52900,
Israel [Tel: 972-3-5318224; Fax: 972-3-5351824]

(Received 4 January 1994)

Abstract—1. Injection of serotonin (5-hydroxytryptamine) to rats, induced a dramatic fall in brain ATP level, accompanied by an increase in P_i . Concomitant to these changes, the activity of cytosolic phosphofructokinase, the rate-limiting enzyme of glycolysis, was significantly enhanced. Stimulation of anaerobic glycolysis was also reflected by a marked increase in lactate content in brain.

2. Brain glucose 1,6-bisphosphate level was decreased, whereas fructose 2,6-bisphosphate was unaffected by serotonin.

3. All these serotonin-induced changes in brain, which are characteristic for cerebral ischemia, were prevented by treatment with the calmodulin (CaM) antagonists, trifluoperazine or thioridazine.

4. Injection of serotonin also induced a marked elevation of plasma hemoglobin, reflecting lysed erythrocytes, which was also prevented by treatment with the CaM antagonists.

5. The present results suggest that CaM antagonists may be effective drugs in treatment of many pathological conditions and diseases in which plasma serotonin levels are known to increase.

Key Words: Calmodulin antagonists, glycolysis, serotonin (5-hydroxytryptamine), phosphofructokinase, glucose 1,6-bisphosphate, fructose 2,6-bisphosphate, brain (cerebral) ischemia, plasma hemoglobin

INTRODUCTION

Plasma serotonin (5-hydroxytryptamine, 5-HT) levels are known to increase in many pathological conditions and diseases. For example, a 15-fold increase in plasma serotonin levels have been demonstrated during cerebrovascular thrombosis (Wester *et al.*, 1992). Elevation of plasma serotonin level by intravenous infusion of this amine into rats, significantly increased blood-brain barrier permeability and reduced cerebral blood flow (Sharma *et al.*, 1990). These findings prompted us to investigate the effects of serotonin on ATP levels and glycolysis in rat brain. Previous experiments in our laboratory have revealed that serotonin exerts a

deleterious effect on muscle and skin glucose metabolism, which could be prevented by treatment with calmodulin (CaM) antagonists (Beitner *et al.*, 1982, 1983; Kaplansky and Beitner, 1984). We also found that CaM antagonists are most effective drugs for treatment of skin trauma (Beitner, 1987; Beitner *et al.*, 1989a, b, 1991) and muscle damage (Beitner and Lilling, 1993), where serotonin is known to play a pathogenic role.

We show here that injection of serotonin to rats induces in brain a dramatic decrease in ATP level, accompanied by stimulation of the activity of cytosolic phosphofructokinase (PFK), the rate-limiting enzyme of glycolysis, and accumulation of lactate. We also show that injection of serotonin causes a marked increase in plasma hemoglobin. All these pathological changes induced by serotonin were prevented by

*To whom all correspondence should be addressed.

treatment with the CaM antagonists, trifluoperazine (TFP) or thioridazine (TRZ).

MATERIALS AND METHODS

Materials

5-hydroxytryptamine hydrochloride was obtained from Sigma Chemical Co. Trifluoperazine dihydrochloride was purchased from ICN Flow Co. and Thioridazine hydrochloride from Taro Pharmaceutical Co., Inc., Haifa, Israel. Other chemicals and enzymes were either from Sigma Chemical Co. or from Boehringer Mannheim GmbH.

Injection of serotonin and CaM antagonists

Charles River albino rats (90–120 g) fed *ad libitum* were injected i.p. with serotonin (40 mg/kg) in 0.1 ml saline. Identical controls received 0.1 ml saline. 10 min later the rats were anesthetized with sodium nembutal (60 mg/kg) and the brains were removed.

In the experiments in which effects of CaM antagonists were studied, trifluoperazine (TFP) or thioridazine (TRZ) at a concentration of 80 mg/kg body wt in 0.1 ml water were injected i.p. 100 min prior to injection of serotonin.

Brains were removed and rapidly frozen between a pair of aluminum tongs precooled in liquid N₂ and stored in liquid N₂ until used (24 hr). Frozen brains were powdered in a mortar cooled in liquid N₂. The powder was used for extraction of metabolites and enzymes.

Preparation and assay of bound and soluble PFK

Cytoskeleton-bound and soluble PFK were separated as described previously (Lilling and Beitner, 1990; Lilling *et al.*, 1991). PFK was assayed under maximal (optimal) conditions (pH 8.2), as described by Mansour (1972). PFK was also assayed under allosteric (suboptimal) conditions (pH 7.35), in which it is subject to allosteric regulation, in an assay mixture similar to that described by Mansour (1972), except that ATP concentration was 0.5–1 mM. One unit of PFK activity catalyzed the formation of 1 μ mol of fructose 1,6-bisphosphate per min at 25°C.

Extraction and determination of metabolites

All metabolites except fructose 2,6-bisphosphate (Fru-2,6-P₂) were extracted in 2 ml perchloric acid as described previously (Beitner *et al.*, 1978). Glucose 1,6-bisphosphate (Glc-1,6-P₂) was measured by the fluorometric method of Passonneau *et al.* (1969). ATP and inorganic phosphate were measured by the methods of Lowry *et al.* (1964). Lactate was measured with the kits from Sigma Chemical Co. Fru-2,6-P₂ was extracted and assayed by the method described by Van Schaftingen *et al.* (1982) with the modification introduced by Van Schaftingen and Hers (1983).

Hemoglobin determination

Hemoglobin in plasma was measured with the kit no. 527 from Sigma Chemical Co.

Protein determination

Protein was measured by the method of Lowry *et al.* (1951) with crystalline bovine serum albumin as a standard.

Statistical analysis

Experimental results are expressed as mean \pm SE. Student's *t*-test was used to determine statistical significance of differences between groups; *P* < 0.05 was considered significant.

RESULTS

The experiments presented in Table 1 reveal that injection of serotonin to rats induced a drastic fall in the level of brain ATP (to 24% of normal level), accompanied by an increase in P_i. The levels of Glc-1,6-P₂, the potent regulator of carbohydrate metabolism (for reviews, see Beitner, 1979, 1984, 1985, 1990, 1993), were significantly decreased, whereas Fru-2,6-P₂ remained unchanged. Lactate content was markedly (more than 3-fold) elevated, suggesting stimulation of glycolysis.

Table 2 shows the effect of serotonin on brain cytoskeleton-bound and soluble (cytosolic) PFK, the rate-limiting enzyme of glycolysis. The allosteric activity was assayed under regulatory (suboptimal) conditions (pH 7.35), in which the enzyme is subject

Table 1. Effects of serotonin on the levels of ATP, P_i, Glc-1,6-P₂, Fru-2,6-P₂ and lactate in brain

Conditions	ATP (mmol/kg)	P _i	Glc-1,6-P ₂ (μ mol/kg)	Fru-2,6-P ₂ (μ mol/kg)	Lactate (mg/g)
Control	1.24 \pm 0.24	5.89 \pm 0.81	84.0 \pm 6.1	7.34 \pm 1.05	0.39 \pm 0.16
Serotonin	0.30 \pm 0.11	9.54 \pm 1.00	46.7 \pm 12.7	6.74 \pm 0.89	1.22 \pm 0.07
% change	-76	+62	-44	—	+213
<i>P</i> value	<i>P</i> < 0.005	<i>P</i> < 0.005	<i>P</i> < 0.005	N.S.	<i>P</i> < 0.005

Values are means \pm SE for 6–12 experiments.
N.S., not significant.

Table 2. Effects of serotonin on cytoskeleton-bound and soluble PFK in brain

Conditions	PFK allosteric activity (units/g)		PFK maximal activity (units/g)	
	Cytoskeleton-bound	Soluble	Cytoskeleton-bound	Soluble
Control	0.27 ± 0.13	1.65 ± 0.21	2.18 ± 0.25	13.33 ± 0.72
Serotonin	0.29 ± 0.08	3.60 ± 0.60	2.32 ± 0.27	12.33 ± 1.36
% change	—	+118%	—	—
P value	N.S.	$P < 0.005$	N.S.	N.S.

Values are means \pm SE for 6–12 experiments.
N.S., not significant.

to regulation by its allosteric effectors, and the maximal activity was assayed under maximal (optimal) conditions (pH 8.2), in which the enzyme is not sensitive to its allosteric effectors. It can be seen that injection of serotonin induced a significant enhancement of the allosteric activity of soluble but not cytoskeleton-bound PFK. The maximal activity of the enzyme remained unchanged in both cell fractions. These results reveal that serotonin did not affect the intracellular distribution of PFK, and the marked stimulation of cytosolic PFK resulted most

probably from changes in the concentration of its allosteric effectors.

In the experiments presented in Figs 1–3 we studied whether treatment with the CaM antagonists, trifluoperazine (TFP) or thioridazine (TRZ), would prevent the pathological changes induced by serotonin. It can be seen that both CaM antagonists reversed the pathological decrease in ATP [Fig. 1(a)] as well as the increase in P_i [Fig. 1(b)], to normal control levels. These compounds also prevented the increase in soluble PFK activity [Fig. 2(a)] and lactate




Fig. 1. The antagonistic effect of CaM inhibitors to the action of serotonin on ATP (a) and P_i (b) levels in brain. Values are mean \pm SE for 6–12 experiments. * $P < 0.005$ (5-HT vs control, or 5-HT + CaM inhibitor vs 5-HT).

Fig. 2. The antagonistic effect of CaM inhibitors to the action of serotonin on soluble PFK allosteric activity (a) and lactate levels (b) in brain. Values are mean \pm SE for 6–12 experiments. * $P < 0.005$; ** $P < 0.05$ (5-HT vs control, or 5-HT + CaM inhibitor vs 5-HT).




Fig. 3. The antagonistic effect of CaM inhibitors to the action of serotonin on Glc-1,6-P₂ levels in brain. Values are mean \pm SE for 6–12 experiments. * $P < 0.005$ (5-HT vs control, or 5-HT + CaM inhibitor vs 5-HT).

content [Fig. 2(b)]. The CaM antagonists also reversed the reduction in Glc-1,6-P₂ level induced by serotonin (Fig. 3). The CaM antagonists alone had no effect on any of the parameters studied.

During these studies, we observed that injection of serotonin causes blood hemolysis. We therefore measured plasma hemoglobin obtained from rats following the injection of serotonin. As shown in Fig. 4, injection of serotonin induced a marked (about 5-fold) increase in plasma hemoglobin. This increase was prevented by treatment with TFP or TRZ. The CaM antagonists alone had no effect on plasma hemoglobin.

DISCUSSION

The present results reveal that injection of serotonin to rats induces a dramatic fall in ATP level in brain, accompanied by an increase in P_i (Table I).




Fig. 4. The antagonistic effect of CaM inhibitors to the action of serotonin on plasma hemoglobin levels. Values are mean \pm SE for 6–12 experiments. * $P < 0.005$ (5-HT vs control, or 5-HT + CaM inhibitor vs 5-HT).

Concomitant to the decrease in ATP, which is an allosteric inhibitor of PFK, and the increase in P_i an allosteric activator of this enzyme, the allosteric activity of PFK in cytosol was markedly enhanced (Table 2). In contrast to cytosolic (soluble) PFK, the allosteric activity of cytoskeleton-bound enzyme was not affected by serotonin, since it is less sensitive to allosteric inhibition by ATP and activation by P_i (results not shown). The enhancement of cytosolic PFK, the rate-limiting enzyme of glycolysis, leads to stimulation of glycolysis, which is reflected by the marked increase in lactate (Table 1).

The serotonin-induced changes in ATP, P_i, PFK and glycolysis, as well as the reduction in Glc-1,6-P₂, are typical changes which we found to occur in tissue during anoxia (Beitner *et al.*, 1979). Serotonin was reported to induce cerebral arteriolar constriction (Thompson *et al.*, 1984). It also increases blood–brain barrier permeability and reduces regional cerebral blood flow (Olesen, 1985; Sharma *et al.*, 1990). Through these actions, serotonin may induce brain ischemia, which is reflected here by the fall in ATP and the stimulation of anaerobic glycolysis, changes that are characteristic for cerebral ischemia (Sims, 1992). Serotonin has been implicated in the pathogenesis of neuronal damage during ischemia (Globus *et al.*, 1992). Serotonin or ischemia is known to induce accumulations of intracellular calcium which causes cell damage. It was recently found that the calmodulin antagonists, phenothiazines, reduce brain damage after ischemia (Zivin *et al.*, 1989; Yu *et al.*, 1992). Calmodulin antagonists are also cardioprotective in ischemic heart, preserving myocardial ATP (Sargent *et al.*, 1992). We show here that the changes induced by serotonin in brain ATP, P_i, PFK, lactate and Glc-1,6-P₂, were prevented by treatment with the calmodulin antagonists, TFP or TRZ (Figs 1–3).

An interesting observation is the increase in plasma hemoglobin induced by serotonin (Fig. 4). This increase, which reflects lysed erythrocytes, is compatible with the *in vitro* experiments, which have shown that exposure of erythrocytes to serotonin increases their osmotic fragility (Kirshtein and Gilboa-Garber, 1975). The increase in plasma hemoglobin may be a useful parameter to measure, in various pathological conditions in which plasma serotonin concentration is increased. The calmodulin antagonists, TFP or TRZ, prevented the serotonin-induced elevation in plasma hemoglobin (Fig. 4), suggesting a protective action on erythrocyte membrane against the damaging effect of serotonin. We have also previously found that serotonin induces pathological changes in glucose metabolism in muscle and skin, which were prevented by treatment with calmodulin antagonists (Beitner *et al.*, 1982, 1983;

Kaplansky and Beitner, 1984). Calmodulin antagonists were also found in our laboratory to be effective drugs in treatment of skin injuries, e.g. burns or frostbite (Beitner, 1987; Beitner *et al.*, 1989a, b, 1991), as well as in treatment of muscle damage (Beitner and Lilling, 1993). It has recently become evident that calmodulin antagonists are a new generation of drugs with broad therapeutic applications (for review, see Mannhold and Timmerman, 1992).

The present findings which have revealed that calmodulin inhibitors antagonize the pathological effects of serotonin in brain, as well as the serotonin-induced raise in plasma hemoglobin, suggest that these compounds may be effective drugs in various pathological conditions and diseases in which plasma serotonin levels are increased.

Acknowledgements—The skillful technical assistance of Mrs H. Morgenstern is acknowledged with thanks. We also thank Ms Sharon Victor for typing this manuscript. This work was supported in part by the ALSAM Foundation, Los Angeles, Calif.; the Migdal Insurance Company, Israel; the Health Sciences Research Center and the Research Committee, Bar-Ilan University.

REFERENCES

- Beitner R. (1979) The role of glucose-1,6-diphosphate in the regulation of carbohydrate metabolism in muscle. *Trends Biochem. Sci.* **4**, 228–230.
- Beitner R. (1984) Control of levels of glucose 1,6-bisphosphate. *Int. J. Biochem.* **16**, 579–585.
- Beitner R. (1985) Glucose 1,6-bisphosphate—the regulator of carbohydrate metabolism. In *Regulation of Carbohydrate Metabolism* (Edited by Beitner R.), Vol. 1, pp. 1–27. CRC Press, Boca Raton, Fla.
- Beitner R. (1987) Method and composition for the therapeutic and prophylactic treatment of trauma to the skin. U.S. Patent No. 4,654,323; No. 4,777,171 and No. 4,910,197.
- Beitner R. (1990) Regulation of carbohydrate metabolism by glucose-1,6-bisphosphate in extrahepatic tissues; comparison with fructose-2,6-bisphosphate. *Int. J. Biochem.* **22**, 553–557.
- Beitner R. (1993) Control of glycolytic enzymes through binding to cell structures and by glucose 1,6-bisphosphate under different conditions. The role of Ca^{2+} and calmodulin. *Int. J. Biochem.* **25**, 297–305.
- Beitner R. and Lilling G. (1993) Treatment of muscle damage, induced by high intracellular Ca^{2+} , with calmodulin antagonists. *Gen. Pharmac.* **24**, 847–855.
- Beitner R., Haberman S. and Nordenberg J. (1978) The effect of epinephrine and dibutyl cyclic AMP on glucose 1,6-bisphosphate levels and the activities of hexokinase, phosphofructokinase and phosphoglucosyltransferase in the isolated rat diaphragm. *Molec. Cell. Endocr.* **10**, 135–147.
- Beitner R., Cohen T. J., Nordenberg J. and Haberman S. (1979) Glucose-1,6-bisphosphate and the mechanism of Pasteur effect in diaphragm muscle. The regulation of hexokinase and phosphofructokinase activities. *Biochim. Biophys. Acta* **586**, 266–277.
- Beitner R., Kaplansky M. and Frucht H. (1982) Effect of serotonin on glucose-1,6-diphosphate levels in rat muscle and skin. *IRCS Med. Sci.* **10**, 300–301.
- Beitner R., Frucht R. and Kaplansky M. (1983) Changes in the levels of glucose 1,6-diphosphate and cyclic GMP, and in the activities of phosphofructokinase and phosphoglucosyltransferase induced by serotonin in muscle. *Int. J. Biochem.* **15**, 935–940.
- Beitner R., Chen-Zion M., Sofer-Bassukevitz Y., Oster Y., Ben-Porat H. and Morgenstern H. (1989a) Therapeutic and prophylactic treatment of skin burns with several calmodulin antagonists. *Gen. Pharmac.* **20**, 165–173.
- Beitner R., Chen-Zion M., Sofer-Bassukevitz Y., Morgenstern H. and Ben-Porat H. (1989b) Treatment of frostbite with the calmodulin antagonists thioridazine and trifluoperazine. *Gen. Pharmac.* **20**, 641–646.
- Beitner R., Chen-Zion M. and Bassukevitz Y. (1991) Effect of the calmodulin antagonist CGS 9343B on skin burns. *Gen. Pharmac.* **22**, 67–72.
- Globus M. Y.-T., Wester P., Busto R. and Dietrich W. D. (1992) Ischemia-induced extracellular release of serotonin plays a role in CA1 neuronal cell death in rats. *Stroke* **23**, 1595–1601.
- Kaplansky M. and Beitner R. (1984) The antagonistic effect of trifluoperazine to the action of serotonin on glucose 1,6-bisphosphate and ATP levels in rat skin and muscle. *IRCS Med. Sci.* **12**, 174–175.
- Kirshtein R. and Gilboa-Garber N. (1975) Effect of serotonin and its derivatives on human erythrocyte osmotic fragility. *Isr. J. Med. Sci.* **11**, 1222.
- Lilling G. and Beitner R. (1990) Decrease in cytoskeleton-bound phosphofructokinase in muscle induced by high intracellular calcium, serotonin and phospholipase A_2 *in vivo*. *Int. J. Biochem.* **22**, 857–863.
- Lilling G., Bassukevitz Y. and Beitner R. (1991) Lithium-induced changes in cytoskeleton-bound phosphofructokinase and mitochondrially-bound hexokinase in rat brain and muscles. *Lithium* **2**, 103–108.
- Lowry O. H., Rosebrough N. J., Farr A. L. and Randall R. J. (1951) Protein measurement with the Folin phenol reagent. *J. biol. Chem.* **193**, 265–275.
- Lowry O. H., Passonneau J. V., Hasselberger F. X. and Schulz D. W. (1964) Effect of ischemia on known substrates and cofactors of the glycolytic pathway in brain. *J. biol. Chem.* **239**, 18–30.
- Mannhold R. and Timmerman H. (1992) Putative therapeutic applications of calmodulin antagonists. *Pharm. Weekbl. [Sci.]* **14**, 161–166.
- Mansour T. E. (1972) Phosphofructokinase activity in skeletal muscle extracts following administration of epinephrine. *J. biol. Chem.* **247**, 6059–6066.
- Olesen S.-P. (1985) A calcium-dependent reversible permeability increase in microvessels in frog brain, induced by serotonin. *J. Physiol.* **361**, 103–113.
- Passonneau J. V., Lowry O. H., Schulz D. W. and Brown J. G. (1969) Glucose 1,6-diphosphate formation by phosphoglucosyltransferase in mammalian tissues. *J. biol. Chem.* **244**, 902–909.
- Sargent C. A., Sleph P. G., Pzwonczyk S., Smith M. A. and Grover G. J. (1992) Effect of calmodulin and protein kinase C inhibitors on globally ischemic rat hearts. *J. Cardiovasc. Pharmac.* **20**, 251–260.
- Sharma H. S., Olsson Y. and Dey P. K. (1990) Changes in blood-brain barrier and cerebral blood flow following elevation of circulating serotonin level in anesthetized rats. *Brain Res.* **517**, 215–223.
- Sims N. R. (1992) Energy metabolism and selective neuronal vulnerability following global cerebral ischemia. *Neurochem. Res.* **17**, 923–931.
- Thompson J. A., Wei E. P. and Kontos H. A. (1984) Inhibition by ketanserin of serotonin-induced cerebral arteriolar constriction. *Stroke* **15**, 1021–1024.
- Van Schaftingen E. and Hers H. G. (1983) Fructose 2,6-bisphosphate in relation with the resumption of metabolic activity in slices of Jerusalem artichoke tubers. *FEBS Lett.* **164**, 195–200.
- Van Schaftingen E., Lederer B., Bartrons R. and Hers H. G. (1982) A kinetic study of pyrophosphate: fructose-6-

- phosphate phosphotransferase from potato tubers. Application to a microassay of fructose 2,6-bisphosphate. *Eur. J. Biochem.* **129**, 191–195.
- Wester P., Dietrich W. D., Prado R., Watson B. D. and Globus M. Y.-T. (1992) Serotonin release into plasma during common carotid artery thrombosis in rats. *Stroke* **23**, 870–875.
- Yu M. J., McCowan J. R., Smalstig E. B., Bennet D. R., Roush M. E. and Clemens J. A. (1992) A phenothiazine derivative reduces rat brain damage after global or focal ischemia. *Stroke* **23**, 1287–1291.
- Zivin J. A., Kochhar A. and Saitoh T. (1989) Phenothiazines reduce ischemic damage to the central nervous system. *Brain Res.* **482**, 189–193.