

# Control of Hepatocyte DNA Synthesis by Intracellular pH and Its Role in the Action of Tumor Promoters

CHOW H. LEE,<sup>1\*</sup> EDWARD J. CRAGOE Jr.,<sup>2</sup> AND ANTHONY M. EDWARDS<sup>3</sup>

<sup>1</sup>Chemistry Program, University of Northern British Columbia, Prince George, Canada

<sup>2</sup>Nacodoches, Texas

<sup>3</sup>Department of Medical Biochemistry, School of Medicine, Flinders University, Adelaide, South Australia, Australia

The mechanisms of tumor promotion in liver by various xenobiotics of diverse structure are not well understood. However, these tumor promoters share the ability to exert growth-stimulatory effects on hepatocytes. Our laboratory has been utilizing normal rat hepatocytes under defined conditions of primary cultures, to investigate growth-stimulatory actions of liver tumor promoters. We have shown that most, if not all, of the liver tumor promoters tested stimulate hepatocyte DNA synthesis when added in combination with epidermal growth factor (EGF), insulin, and glucocorticoids. In the present study, we sought evidence for the role of the  $\text{Na}^+/\text{H}^+$  antiporter and cytoplasmic alkalization in the direct growth-stimulatory actions of tumor promoters on hepatocytes. Hepatocytes cultured under conditions (bicarbonate-buffered medium) where intracellular pH ( $\text{pH}_i$ ) was independent of extracellular pH ( $\text{pH}_e$ ), EGF- and insulin-stimulated rates of DNA synthesis were unaffected by modest changes in  $\text{pH}_e$ . However, under conditions (HEPES-buffered medium) where  $\text{pH}_i$  varied in a linear fashion with  $\text{pH}_e$ , rates of EGF- and insulin-stimulated DNA synthesis were highly dependent on  $\text{pH}_e$ . Similarly, 12-O-tetradecanoylphorbol-13-acetate (TPA) and  $\alpha$ -hexachlorocyclohexane (HCH)-stimulated DNA synthesis were  $\text{pH}_e$ -dependent but were stimulatory over different  $\text{pH}_e$  ranges, suggesting that these promoters may act by distinct mechanisms. Chemicals that are capable of inducing rapid cytoplasmic alkalization, ammonium chloride (1 and 15 mM) and monensin (0.5  $\mu\text{M}$ ), were found to stimulate hepatocyte DNA synthesis. The role of the  $\text{Na}^+/\text{H}^+$  antiport in controlling  $\text{pH}_i$  of hepatocytes was demonstrated by artificially acidifying 2',7'-bis(carboxyethyl)-5,6-carboxyfluorescein acetoxymethyl (BCECF)-loaded isolated hepatocytes with 20 mM sodium acetate and the use of specific inhibitors. Amiloride and its analogues inhibited  $\text{pH}_i$  recovery from the acid load in a dose dependent manner and the relative potency of these inhibitors paralleled their  $K_i$  values for the  $\text{Na}^+/\text{H}^+$  antiport. At concentrations that stimulate hepatocyte DNA synthesis, some liver tumor promoters phenobarbital (PB) and HCH, were found to cause a rapid rise  $\text{pH}_i$  in isolated hepatocytes which was sensitive to amiloride and its analogues. Taken together, our data suggest that activation of  $\text{Na}^+/\text{H}^+$  antiport activity may be one mechanism whereby some liver tumor promoters stimulate hepatocytes DNA synthesis. This study has implications for the mechanisms of tumor promotion in liver carcinogenesis. J. Cell. Physiol. 195: 61–69, 2003. © 2003 Wiley-Liss, Inc.

The  $\text{Na}^+/\text{H}^+$  antiporter is the most widely studied intracellular pH ( $\text{pH}_i$ ) regulator in vertebrate cells. It is now known to be a ubiquitous feature of the plasma membrane of mammalian cells (reviewed by Madhus, 1988; Grinstein et al., 1989), including rat hepatocytes (Arias and Forgac, 1984; Moseley et al., 1986a; Anwer and Nolan, 1988; Renner et al., 1989a). In response to a fall in  $\text{pH}_i$ , the  $\text{Na}^+/\text{H}^+$  antiporter functions to expel  $\text{H}^+$  from the cytoplasm, using the inwardly directed  $\text{Na}^+$ -gradient created by  $\text{Na}^+/\text{K}^+$ -ATPase.

The  $\text{Na}^+/\text{H}^+$  antiporter is believed to play a role in regulating various metabolic events that are dependent on  $\text{pH}_i$ , including DNA synthesis. The first evidence to

Contract grant sponsor: Anti-cancer Foundation of South Australia.

\*Correspondence to: Chow H. Lee, Chemistry Program, University of Northern BC, 3333 University Way, Prince George, BC V2N 4Z9, Canada. E-mail: leec@unbc.ca

Received 25 June 2002; Accepted 5 November 2002

DOI: 10.1002/jcp.10225

associate activation of  $\text{Na}^+/\text{H}^+$  antiporter and initiation of growth came from studies on the activation of the sea urchin eggs (Johnson et al., 1976). A rapid increase in  $\text{pH}_i$  after fertilization was seen as a result of the activation of amiloride-sensitive exchange of extracellular  $\text{Na}^+$  for intracellular  $\text{H}^+$ . Substantial evidence now suggests that activation of the antiport and/or subsequent cytoplasmic alkalinization may be required for cell proliferation. Many mitogens and co-mitogens, including growth factors, serum, and phorbol esters, have been shown to rapidly (within 1 min) activate the antiporter in many cell types, thus producing an intracellular alkalinization of about 0.15–0.3 pH units (Frelin et al., 1988; Quinn et al., 1996; Di Sario et al., 2001). The observation that many growth factors increase  $\text{pH}_i$  has led to the notion that increased  $\text{pH}_i$  may be a necessary early signaling event in growth factor action. Interestingly,  $\text{Na}^+/\text{H}^+$  antiporter-dependent intracellular alkalinization is also considered to be an early event in malignant transformation, and appears to play an essential role in the development of transformation-associated phenotypes (Reshkin et al., 2000). Further evidence to link  $\text{Na}^+/\text{H}^+$  antiport with growth comes from studies using specific inhibitors of the antiporter. Amiloride and its analogues have been shown to inhibit cell division in sea urchin (Johnson et al., 1976) and growth factor-induced DNA synthesis in several mammalian cell types (reviewed by Soltoff and Cantley, 1988; Grinstein et al., 1989). Koch and Leffert (1980) have further shown that DNA synthesis can be re-initiated upon removal of amiloride, consistent with the notion that  $\text{Na}^+/\text{H}^+$  antiporter plays a role in events preceding DNA synthesis.

Koch and Leffert (1979, 1980) provided the initial evidence for a possible role of  $\text{Na}^+/\text{H}^+$  antiport in hepatocyte growth. Addition to long-term hepatocyte cultures of a mixture of insulin, glucagon and epidermal growth factor (EGF), was found to stimulate rapid influx of  $^{22}\text{Na}^+$ , followed 24–28 h later by incorporation of thymidine into DNA. Such stimulation of DNA synthesis was blocked by removal of  $\text{Na}^+$  or administration of amiloride (Koch and Leffert, 1979, 1980).  $\text{Na}^+/\text{H}^+$  antiport activity has been shown to be increased in liver plasma membranes from hepatectomized and neonatal rats, further suggesting a link between the antiporter and hepatocyte growth (Moseley et al., 1986a; Goodrich and Suchy, 1990). Growth-stimulatory agents such as EGF, hepatocyte growth factor, and TPA, which are known to increase DNA synthesis in hepatocytes, have also been shown to activate the  $\text{Na}^+/\text{H}^+$  antiport (Smith and Boyer, 1986; Moule and McGivan, 1990; Tanaka et al., 1994; Kaneko et al., 1995). Other related findings in liver include inhibition by amiloride of nafenopin-stimulated DNA synthesis (Ochsner et al., 1990) and mitogen-induced protooncogenes expression in primary rat hepatocytes (Kruijer et al., 1986; Leffert et al., 1990). Similarly, inhibition by amiloride and its analogues of intracellular alkalinization and DNA synthesis have been observed in hepatoma cells (Strazzabosco et al., 1995; Garcia-Canero et al., 1999).

Our laboratory has been utilizing normal rat hepatocytes under defined conditions of primary cultures, to investigate growth-stimulatory actions of tumor promoters, an important element in the promotion stage of

carcinogenesis (Yusof and Edwards, 1990; Lee and Edwards, 2001a,b). The aims of the present study are: (1) to clarify the significance of  $\text{pH}_i$  changes in control of hepatocyte DNA synthesis; and (2) to investigate whether liver tumor promoters act like growth factors to rapidly activate the  $\text{Na}^+/\text{H}^+$  antiport, thereby inducing intracellular alkalinization.

## MATERIALS AND METHODS

### Animals and materials

Male wistar rats (Porton strain), weighing about 200 g, given free access to laboratory chow and water, were used for hepatocyte preparation in all experiments. Williams' Medium E (WME) was obtained from Flow Laboratories (Irvine, Scotland), Swim's S-77 medium and antibiotics were from GIBCO (Grand Island). LHAS medium was prepared as described by Oliver et al. (1978). Collagenase was from Boehringer Mannheim GmbH (Penzberg, Germany);  $\alpha$ -hexachlorocyclohexane (HCH) and 1,1-bis(*p*-chlorophenyl)-2,2,2-trichloroethane (DDT) were from Serva Feinbiochemica (Heidelberg, Germany) or from Aldrich (Milwaukee, WI); phenobarbital (PB) or its sodium salt were from Prosanna Laboratories (Carole Park, Queensland, Australia). Pregnenolone-16 $\alpha$ -carbonitrile (PCN), 12-*o*-tetradecanoylphorbol-13-acetate (TPA) and all other chemicals were from Sigma (St. Louis, MO). Amiloride analogues, 5-(*N*-ethyl-*N*-isopropyl)-amiloride (EIA), 5-(*N*,*N*-hexamethylene)-amiloride (HMA), 5-(*N*-methyl-*N*-(guanidino-carbonylmethyl)-amiloride (MGA), and 5-(*N*-methyl-*N*-tert-butyl)-amiloride (MBA), were provided by Dr. Edward J. Cragoe, Jr., as part of a collaborative study. The fluorescent dye, 2',7'-bis(carboxyethyl)-5,6-carboxyfluorescein acetoxymethyl ester (BCECF-AM) was from Molecular Probes (Eugene, OR). Type I collagen was extracted from rat tails, essentially as described by Michalopoulos and Pitot (1975). Culture dishes were collagen-coated using a solution of 0.3 mg collagen/ml 0.1% acetic acid. Stock solutions of test agents were prepared weekly. The final concentration of dimethylsulfoxide (DMSO) when added to cultures was 0.2% or less and had no effect on DNA synthesis.

### Hepatocyte isolation and culture conditions

Hepatocytes were isolated using a two-step procedure as previously described (Whiting and Edwards, 1979). Low centrifugal speeds (50  $\times g$ ) were used during washing of cells with LHAS medium to minimize the contamination of hepatocytes with smaller non-parenchymal cells. The washed hepatocyte pellet was resuspended in warm WME containing antibiotics (penicillin, 100 U/ml, streptomycin sulphate 100  $\mu\text{g}/\text{ml}$ ), 300 nM insulin, and 3% fetal bovine serum. For DNA synthesis experiments, cells were plated on 50 mm collagen-coated dishes at a density of about  $3.5\text{--}4 \times 10^4$  cells/cm<sup>2</sup>. Higher cell density, as indicated below, were used for other types of experiments. Cells were allowed to attach for 3–4 h at 37°C in a humidified atmosphere of 5% CO<sub>2</sub> in air. After the attachment period, cells were maintained in fresh WME (serum-free) containing antibiotics as above. This is termed 'standard culture medium'. Insulin (300 nM), dexamethasone (30 nM), and EGF (10 ng/ml) were added

depending on the nature of the experiments as indicated in each figure legend. Other test compounds were added either after attachment, or with subsequent daily medium changes, or both, as indicated in the figure legends.

### Measurements of the rate of hepatocyte DNA synthesis

After various times in culture, medium was replaced with fresh standard medium containing  $0.08 \mu\text{Ci/ml}$  [methyl- $^3\text{H}$ ]thymidine ( $20 \text{ Ci/mmol}$ , New England Nuclear, Boston, MA), and cells incubated for 3 h. Following extensive washing, DNA was extracted (Yusof and Edwards, 1990), with one aliquot taken for scintillation counting and another to determine DNA concentration (Burton, 1956). The values shown in each figure are means  $\pm$  SD from groups of four dishes per treatment obtained with a single hepatocyte preparation. Results were confirmed using independent cell preparations.

### Measurements of changes in intracellular $\text{pH}$

Buffers used were based on those described by Moolenaar et al. (1983) for studies with fibroblasts. Freshly isolated hepatocytes ( $50 \text{ mg wet weight/ml}$ ) were resuspended in cold Buffer A ( $140 \text{ mM NaCl}$ ,  $5 \text{ mM KCl}$ ,  $2 \text{ mM CaCl}_2$ ,  $1 \text{ mM MgCl}_2$ ,  $10 \text{ mM HEPES}$ , and  $10 \text{ mM glucose}$ , adjusted to  $\text{pH } 7.1$ ), and incubated with  $5 \mu\text{M BCECF}$ -acetoxymethyl ester ( $\text{pK}_a 6.98$ ) for 30 min at  $37^\circ\text{C}$ . Cells were continuously gassed with oxygen and shaken constantly ( $100 \text{ rpm}$ ) during this period. The non-fluorescent BCECF-AM readily entered the cells where cytosolic esterases cleaved the ester bonds, releasing the anionic BCECF which exhibits  $\text{pH}$ -sensitive fluorescence. Cells were pelleted ( $30 \text{ sec at } 50 \times g$ ) and resuspended at a density of  $30 \text{ mg wet weight/ml}$  in Buffer B (same as Buffer A except at  $\text{pH } 7.4$ ). A further 15–20 min incubation at  $37^\circ\text{C}$  was necessary to allow complete intracellular de-esterification of BCECF-AM. Hepatocytes were then pelleted ( $30 \text{ sec at } 50 \times g$ ) and resuspended ( $30 \text{ mg wet weight/ml}$ ) in  $2.5 \text{ ml}$  of warm Buffer B at  $\text{pH } 7.4$  and put into a pre-warmed magnetically-stirred cuvette, maintained at  $30$ – $32^\circ\text{C}$ , unless otherwise indicated. The fluorescence emission was monitored using the an Aminco Bowman Spectrofluorimeter (in earlier experiments) or a Perkin Elmer LS 50 Spectrometer (in later experiments) with excitation at  $500 \text{ nm}$  and emission at  $530 \text{ nm}$ . Calibration of cytoplasmic dye fluorescence as a function of  $\text{pH}_i$  was done by the  $\text{H}^+$  equilibration method (Thomas et al., 1979) using the  $\text{K}^+/\text{H}^+$  ionophore nigericin ( $10 \mu\text{M}$ ) and Buffer C, which has high  $\text{K}^+$  concentration ( $150 \text{ mM KCl}$ ,  $10 \text{ mM HEPES}$ ,  $\text{pH } 7.0$ ) approximating intracellular  $\text{K}^+$  concentration. Nigericin was added to cells after determining the position of the baseline. Under this condition,  $[\text{H}^+]_i/[\text{H}^+]_o = [\text{K}^+]_i/[\text{K}^+]_o$ . Changes in fluorescence were recorded with successive additions of  $1 \mu\text{l}$  aliquots of  $1 \text{ M KOH}$ , until the  $\text{pH}$  of the solution was  $7.6$ . External  $\text{pH}_i$  was measured using a Corning  $\text{pH}$  meter. For measurements of changes in  $\text{pH}_i$  upon treatment with test agents, a baseline  $\text{pH}_i$  was typically achieved first. Results presented in Figures 5–7 are representatives of at least three experiments from the same and different batches of isolated hepatocytes.

## RESULTS

### Effects of varying $\text{pH}$ of HEPES- and bicarbonate-buffered medium on the ability of hepatocyte to synthesize DNA

Studies by Pollock (1984) on primary cultured rat hepatocytes have shown an almost linear relationship between  $\text{pH}_i$  and external  $\text{pH}$  of HEPES-buffered medium in the  $\text{pH}$  range  $6.8$ – $8.0$ . In contrast,  $\text{pH}_i$  of cultured hepatocytes in bicarbonate:  $\text{CO}_2$  medium was relatively constant over the  $\text{pH}_i$  range tested. Such findings provided an opportunity to monitor possible changes in the rate of DNA synthesis upon modulation of  $\text{pH}_i$  of cultured hepatocytes.

Previous studies conducted in this laboratory used bicarbonate:  $\text{CO}_2$  medium, and it is not known whether HEPES itself has any effect on hepatocyte DNA synthesis. Therefore, the first step taken was to test directly, effects of HEPES on the ability of cultured hepatocytes (in bicarbonate-buffered medium) to synthesize DNA. A concentration-dependent inhibitory effect of HEPES on [ $^3\text{H}$ ]thymidine incorporation into DNA is shown in Figure 1. At concentrations up to  $10 \text{ mM}$ , HEPES had no effect on DNA synthesis. Higher concentrations were apparently inhibitory although statistically significant inhibition was seen only at  $25 \text{ mM}$  HEPES. For the studies below with HEPES-buffered media,  $10 \text{ mM}$  HEPES was employed.

Bicarbonate buffer (WME) was equilibrated with  $5\%$   $\text{CO}_2$  at  $37^\circ\text{C}$  and adjusted to desired  $\text{pH}$ . Similarly, HEPES-buffered (bicarbonate-free) WME was equi-




Fig. 1. Concentration-dependent effect of HEPES on hepatocyte DNA synthesis in bicarbonate-buffered medium. Freshly isolated hepatocytes were plated onto collagen-coated dishes ( $35 \times 10^3$  cells/cm $^2$ ), and after 4 h for cell attachment in the presence of insulin ( $300 \text{ nM}$ ) and  $3\%$  fetal calf serum, medium was replaced with 'fresh standard culture medium'. Cultures were maintained with daily medium changes. After 40 h in culture, [ $^3\text{H}$ ]thymidine was added to determine rates of DNA synthesis as described in Materials and Methods. Each value is mean  $\pm$  SD ( $n = 8$ ). Where treatment with HEPES resulted in a mean value significantly different ( $P < 0.001$ ) from control (without HEPES), this is shown by an asterisk.




Fig. 2. The effects of varying pH of HEPES- and bicarbonate-buffered media on the ability of hepatocytes to synthesize DNA. Hepatocytes were plated as described in Materials and Methods, and maintained after a 4-h attachment period with either 'standard culture medium' (bicarbonate-buffered) incubated under an atmosphere of 5%  $\text{CO}_2$  in air (■) or 10 mM HEPES-buffered bicarbonate-free Williams' medium E (WME) and incubated with an air atmosphere (□). For each type of medium, initial pHs were adjusted to the values shown. EGF (10 ng/ml), dexamethasone (30 nM), and insulin (300 nM) were present in all cultures and daily medium changes were conducted. Rates of DNA synthesis in all cultures were measured after 40 h using the 'standard culture medium' (pH 7.4) containing titrated thymidine, as described for all other DNA synthesis assays, according to the protocol in Materials and Methods.

brated at  $37^\circ\text{C}$  and adjusted to the desired pH. Figure 2 shows the effects of varying pH of HEPES- and bicarbonate-buffered medium on hepatocyte DNA synthesis measured after 40 h in culture. There were no significant changes in the rate of DNA synthesis over the external pH range of 6.8–7.7 when bicarbonate:  $\text{CO}_2$  medium was used. In contrast, [ $^3\text{H}$ ]thymidine incorporation into DNA was seen to increase with increasing pH of HEPES-buffered medium.

#### Effects of insulin or tumor promoters on hepatocyte DNA synthesis in HEPES-buffered media or varying pH

Since the rate of DNA synthesis was found to increase with increasing pH of HEPES-buffered medium, it was deemed useful to test the effects of promoters on hepatocyte DNA synthesis at these various pHs. Results of this study are presented in Figure 3. None of the compounds tested had an effect on DNA synthesis at external pH of 6.8. As in the previous study, the EGF-stimulated rate of DNA synthesis increased with increasing pH and the ability of insulin, TPA, or HCH to stimulate was also pH-dependent. However, the pH dependence of HCH action differed from that of TPA and insulin. Low pH (6.8–7.2) blocked the stimulatory effect of HCH whereas near-maximal stimulatory effects of insulin or TPA over the effect for EGF alone were seen at pH 7.2. There was little additional stimulation by insulin or TPA as the pH increased from 7.2 to 7.8 whereas stimulation by HCH increased markedly over this range.




Fig. 3. Effects of test agents on hepatocyte DNA synthesis in HEPES-buffered media of varying pH. Hepatocytes were isolated and plated as described in Materials and Methods. After 4 h for cell attachment, medium was replaced with 10 mM HEPES-buffered (bicarbonate-free) WME at various pHs as shown. Also present in the medium were antibiotics, EGF (10 ng/ml) and dexamethasone (30 nM). Some cultures had no addition (■), insulin (300 nM) (○), TPA (300 nM) (△), or HCH (30  $\mu\text{M}$ ) (□). Daily medium changes were conducted and rates of DNA synthesis were measured after 40 h in culture as described in Materials and Methods. Each value is mean  $\pm$  SD ( $n=4$ ).

#### Effects of monensin and ammonium chloride on hepatocyte DNA synthesis

Monensin, a  $\text{Na}^+$ -ionophore (Pressman, 1976), capable of collapsing  $\text{Na}^+$  and  $\text{H}^+$  gradients has been shown to raise  $\text{pH}_i$  of many cells (Mills et al., 1985; Church et al., 1989) including hepatocytes in this study (data not shown). Ammonium chloride ( $\text{NH}_4\text{Cl}$ ) is also known to raise  $\text{pH}_i$  of cells immediately upon addition (Moolenaar et al., 1983; Madhus, 1988). This (result not shown) and previous studies from this laboratory (results not shown) showed similar effects of  $\text{NH}_4\text{Cl}$  on  $\text{pH}_i$  of hepatocytes. It was thus of interest to test the effects of both monensin and  $\text{NH}_4\text{Cl}$  on hepatocyte DNA synthesis. Bicarbonate:  $\text{CO}_2$  medium was used, and EGF and insulin were present throughout the experiments. Rates of DNA synthesis were measured after 40 h in culture. High concentrations of  $\text{NH}_4\text{Cl}$  (15 mM) and monensin (0.5–5  $\mu\text{M}$ ), when present throughout the experiments had non-specific effects causing cells to detach after 1 day in culture (data not shown). Results in Figure 4 show that maintenance of hepatocytes with media containing a lower initial concentration of  $\text{NH}_4\text{Cl}$  (1 mM) or brief exposure (5 h) to a high concentration of  $\text{NH}_4\text{Cl}$  (15 mM), resulted in significant stimulation of DNA synthesis. Low concentrations of monensin (0.05–0.1  $\mu\text{M}$ ) had no significant effect while modest stimulation was seen in cultures exposed for 5 h to 0.5  $\mu\text{M}$  monensin.




Fig. 4. Effects of ammonium chloride and monensin on hepatocyte DNA synthesis. Hepatocytes were isolated and plated as described in Materials and Methods. After 4 h for cell attachment, medium was changed to 'standard culture medium' containing EGF (10 ng/ml), insulin (300 nM), and additions of  $\text{NH}_4\text{Cl}$  and monensin as indicated. All cultures had  $\text{NH}_4\text{Cl}$  and monensin throughout the experiment except for cultures treated with 15 mM  $\text{NH}_4\text{Cl}$  and 0.5  $\mu\text{M}$  monensin, where treatment was only for 5 h after which medium was replaced with 'standard culture medium' containing EGF and insulin. All cultures were subjected to daily medium changes and rates of DNA synthesis were measured after 40 h in culture as described in Materials and Methods. Each bar is mean  $\pm$  SD ( $n=4$ ) where treatment resulted in a mean value significantly different from control (no addition), this is shown by superscript: \* $P < 0.001$ , \*\* $P < 0.02$ .

#### Inhibition by amiloride analogues of hepatocyte $\text{pH}_i$ recovery from an acid load: evidence for a role of $\text{Na}^+/\text{H}^+$ antiporter

The role of  $\text{Na}^+/\text{H}^+$  antiporter in experimental cells can be shown by artificially acidifying cells, either by removal of cells from  $\text{NH}_4\text{Cl}$ -containing media or by addition of sodium acetate or isobutyrate (Moolenaar et al., 1984, 1986; Isom et al., 1987). Figure 5 shows the effect of adding 20 mM sodium acetate to BCECF-loaded hepatocytes. In aqueous solution, acetic acid exists in its dissociated and undissociated forms. The undissociated form readily diffuses across the plasma membrane into the cells. Once inside the cells, dissociation of acetic acid occurs, resulting in the accumulation of  $\text{H}^+$ , thereby causing a sharp drop in  $\text{pH}_i$ . In the absence of inhibitors, this is followed by a slow recovery of  $\text{pH}_i$ , due to the role of  $\text{Na}^+/\text{H}^+$  antiport, extruding excess  $\text{H}^+$  out of the cells (control cells in Fig. 5). Figure 5 shows that EIA, a specific inhibitor of the  $\text{Na}^+/\text{H}^+$  antiport inhibited  $\text{pH}_i$  recovery in a dose-dependent manner, with maximal inhibition at 30  $\mu\text{M}$ . Similar dose-dependent inhibition by other amiloride analogues (HMA, MBA, and MGA) was observed (data not shown). The relative potency of these analogues as inhibitors of  $\text{pH}_i$  recovery and their




Fig. 5. Effects of amiloride analogues on  $\text{pH}_i$  recovery from an acid load in isolated hepatocytes. Hepatocytes were loaded with BCECF, washed and resuspended with warm Buffer B, pH 7.4 as described in Materials and Methods. Inhibitors as shown by the arrow ( $\downarrow$ ), were added, 1 min before the addition of 20 mM sodium acetate ( $\text{CH}_3\text{COONa}$ ). At this time, DMSO (solvent for the inhibitors) was added to control cells (final concentration = 0.1%). The tracings of different treatments were superimposed to allow comparison of  $\text{pH}_i$  recoveries. A: Dose dependent inhibition by EIA of  $\text{pH}_i$  recovery. B: Relative effects of various amiloride analogues as inhibitors of  $\text{pH}_i$  recovery. The inhibitors were: 1, 30  $\mu\text{M}$  MGA; 2, 30  $\mu\text{M}$  MBA; 3, 30  $\mu\text{M}$  EIA; 4, 10  $\mu\text{M}$  HMA.

maximal-effective concentrations (Fig. 5) can be summarized as follows: 10  $\mu\text{M}$  HMA > 30  $\mu\text{M}$  EIA > 30  $\mu\text{M}$  MBA > 30  $\mu\text{M}$  MGA. This parallels the  $K_i$  values of the inhibitors for  $\text{Na}^+/\text{H}^+$  antiport (Simchowicz and Cragoe, 1986). Such data shows that  $\text{pH}_i$  recovery observed from an acid load was mediated predominantly by the  $\text{Na}^+/\text{H}^+$  antiport. The failure of recovery curves to reach a plateau probably reflects the leakage of dye as described below.

#### Use of probenecid to prevent dye leakage

Various investigators had reported that hepatocytes can secrete BCECF (Petzinger et al., 1988; Renner et al., 1989; Ochsner et al., 1990). Similar problems have also been encountered in this lab. This is problematic in experiments where a reproducible stable  $\text{pH}_i$  baseline is required. Probenecid, a blocker of organic anion transport (Di Virgilio et al., 1988), has been shown to block efflux of the fluorescent calcium indicators fura-2

and indo-1 in pancreatic and astrocytoma cells (Arkhammar et al., 1989, 1990; McDonough and Button, 1989). Probenecid was, therefore, tested for effects on hepatocytes pre-loaded with BCECF. Results showed that addition of probenecid blocked the progressive increase in baseline fluorescence in hepatocyte suspensions and had no effect on recovery from an acid load produce by treatment with sodium acetate (data not shown). Similarly, probenecid did not prevent  $\text{pH}_i$  recovery from an acid load produced by treating cells with 15 mM  $\text{NH}_4\text{Cl}$  then transferring cells to  $\text{NH}_4\text{Cl}$ -free medium (data not shown). Figure 6 shows measurement of  $\text{pH}_i$  changes in response to TPA in the presence of probenecid with BCECF secretion blocked by probenecid. TPA caused a clear elevation of  $\text{pH}_i$  and this effect was inhibited by EIA, suggesting that the effect of TPA was mediated by the  $\text{Na}^+/\text{H}^+$  antiport. In experiments where stable  $\text{pH}_i$  baseline was achieved in the absence of probenecid treatment, TPA also reproducibly elevate  $\text{pH}_i$  (Fig. 7).

#### Effects of test agents on $\text{pH}_i$ of isolated hepatocytes

Various agents, including liver tumor promoters, were tested for effects on hepatocytes  $\text{pH}_i$ . Figure 7 shows results from an experiment, typical of many with other separate cell preparations. The phorbol ester, TPA, was found to increase  $\text{pH}_i$  of about 0.1 pH units compared to the control over a 6.5-mins period. Both PB (0.08 pH units) and HCH (0.05 pH units) were also found to further increase  $\text{pH}_i$  compared to the control (Fig. 7) over similar time period. In some cases, control cells (0.1% saline, 0.1% DMSO, or no addition) showed a slight increase in fluorescence over a 15 min period, reflecting dye leakage (see above). Other compounds tested for effects on hepatocytes  $\text{pH}_i$  include monensin and prolactin. 0.5  $\mu\text{M}$  monensin and 20 mU/ml prolactin were found to raise the  $\text{pH}_i$  approximately 0.05 pH units




Fig. 6. Effect of TPA on  $\text{pH}_i$  of hepatocytes, previously treated with probenecid. Freshly isolated hepatocytes were loaded with BCECF (5  $\mu\text{M}$ ), washed and resuspended in warm Buffer B, pH 7.4 as described in Materials and Methods. One millimolar probenecid was added 1 min after an initial fluorescence was obtained and followed by an addition of 600 nM TPA 2 min later, as shown. Fluorescence was monitored over a period of 15 min. Thirty micrometer EIA was added 2 min prior to termination of fluorescence measurement.




Fig. 7. Effects of test agents on  $\text{pH}_i$  of isolated hepatocytes. Freshly isolated hepatocytes were loaded with BCECF (5  $\mu\text{M}$ ), washed and resuspended in warm Buffer B, pH 7.4 as described in Materials and Methods. Test agents were added after 2 min of initial fluorescence, and were monitored for a period of 15 min. The tracings were from the same batch of isolated hepatocytes and are representative from at least 3 experiments. The additions were: (a) 0.1% DMSO; (b) 600 nM TPA; (c) 2 mM PB; (d) 30  $\mu\text{M}$  HCH.

(data not shown). Figure 7 also shows the ability of EIA to decrease the continuous rise in  $\text{pH}_i$  following treatment with PB and HCH, suggesting that the  $\text{Na}^+/\text{H}^+$  antiport is involved in the elevation of  $\text{pH}_i$  by these liver tumor promoters. EIA alone had no effect on the baseline  $\text{pH}_i$  (data not shown).

#### DISCUSSION

This laboratory is interested in determining the mechanisms by which liver tumor promoters stimulate hepatocyte DNA synthesis (Yusof and Edwards, 1990; Petronijevic and Edwards, 1993; Lee and Edwards, 2001a,b). Based on previous studies that the skin tumor promoter, TPA, and growth factors caused rapid elevation of  $\text{pH}_i$  in hepatocytes (Smith and Boyer, 1986; Tanaka et al., 1994; Kaneko et al., 1995), and that early elevation of  $\text{pH}_i$  is frequently associated with subsequent stimulation of DNA synthesis in many cell types (see Introduction), we hypothesized that liver tumor promoters may elevate hepatocyte  $\text{pH}_i$  and this may be relevant to their stimulatory effects on DNA synthesis. There are two major parts in this study. The first section focuses on evidence that hepatocyte DNA synthesis is  $\text{pH}$ -dependent. The second part focuses on determining the role of the  $\text{Na}^+/\text{H}^+$  antiport in controlling hepatocyte  $\text{pH}_i$ . The latter section includes measuring the effects liver tumor promoters on  $\text{pH}_i$  of isolated hepatocytes.

Pollock (1984) showed that when hepatocytes were incubated without  $\text{HCO}_3^-$  (so that  $\text{HCO}_3^-$ -dependent mechanisms are inoperative) and in HEPES-buffered media, the ability of HEPES to move through the hepatocyte membrane results in conditions where  $\text{pH}_i$  varies in linear fashion with  $\text{pH}_e$ . These observations provided a method for experimental manipulation of  $\text{pH}_i$  for studies on its relevance to control of DNA synthesis. By contrast, it is known that in  $\text{HCO}_3^-$ -containing media, hepatocytes can maintain relatively constant  $\text{pH}_i$  in the presence of a range of  $\text{pH}_e$ . As shown in Figure 2, under conditions where  $\text{pH}_i$  is largely unaffected by  $\text{pH}_e$  ( $\text{HCO}_3^-$ -buffered medium),  $\text{pH}_e$  had little effect on DNA synthesis. In contrast, in HEPES-buffered medium where  $\text{pH}_i$  varies with  $\text{pH}_e$ , rates of EGF- and insulin-stimulated DNA synthesis became strongly dependent on  $\text{pH}_e$ . In this study, the  $\text{pH}_e$  (and hence  $\text{pH}_i$ ) were controlled throughout a 36-h period from after cell attachment until beginning the DNA synthesis measurements, so it is possible that  $\text{pH}_i$  is influencing one or more events at any points in the sequence from initial actions of EGF or insulin through to S-phase entry. Results are consistent with findings by Moolenaar et al. (1986) that a  $\text{pH}_i$  above 7.2 is a permissive requirement for human fibroblasts to initiate DNA synthesis.

$\text{NH}_4\text{Cl}$  exposure was found to significantly stimulate hepatocyte DNA synthesis even under conditions where  $\text{HCO}_3^-$   $\text{pH}_i$ -regulating mechanisms are operative (Fig. 4). Although a rapid alkalization effect of  $\text{NH}_4\text{Cl}$  has not been directly demonstrated in hepatocytes in  $\text{HCO}_3^-$ -buffered medium, it is likely that  $\text{NH}_4\text{Cl}$  has a similar effect to that seen in mesangial cells (Ganz et al., 1989). Brief exposure (5 h) to  $\text{NH}_4\text{Cl}$  or monensin may be approximating signaling events and provide some indication that relatively short periods of  $\text{pH}_i$  increase may be sufficient to set cell cycle events in train and that elevated  $\text{pH}_i$  may not be required for full pre-S-phase period. In fact, this notion is supported by our observation that both  $\text{NH}_4\text{Cl}$  and monensin induced hepatic *c-myc* mRNA expression measured after 2 h of exposure (Lee and Edwards, 2003). Furthermore, effects of a single addition of  $\text{NH}_4\text{Cl}$  may of limited duration since  $\text{NH}_4\text{Cl}$  is rapidly metabolized by liver (Williamson et al., 1974). Taken together, our data provided good evidence that hepatocyte DNA synthesis is dependent on  $\text{pH}_i$  and early cytoplasmic alkalization plays a role in regulating the growth of hepatocytes.

Under conditions where  $\text{HCO}_3^-$   $\text{pH}_i$ -regulating mechanisms are inoperative, insulin and TPA exhibited parallel pH dependence with maximal effect near pH 7.2 (Fig. 3). In contrast, HCH had no effect at low pH range (6.8–7.2) but strongly enhanced EGF-stimulated DNA synthesis as the pH increased from 7.2 to 7.8. These observations suggest that growth-stimulatory effects of HCH are more pH-dependent and differ in mechanisms from that of TPA and insulin.

Some difficulties were encountered in our hands when the fluorescent dye, BCECF, was used as an indicator of  $\text{pH}_i$ . This is consistent with findings from several other laboratories (Gleeson et al., 1989; Renner et al., 1989; Ochsner et al., 1990). Probenecid has been used to block efflux of fluorescent calcium indicators from other cell types (Arkhammar et al., 1989, 1990; McDonough and

Button, 1989). We, therefore, opted to pre-treat hepatocytes with probenecid prior to testing agents for effects on  $\text{pH}_i$ . Results showed probenecid could block the gradual rise in fluorescence in control incubations of hepatocytes (Fig. 6) and did not prevent  $\text{pH}_i$  recovery from an acid load (data not shown), suggesting that it does not interfere with  $\text{Na}^+/\text{H}^+$  antiport activity. The rise in  $\text{pH}_i$  stimulated by TPA in hepatocytes previously treated with probenecid, was sensitive to amiloride analogue EIA, suggesting that the effect seen was due to the activation of  $\text{Na}^+/\text{H}^+$  antiporter and that probenecid does not block the response to one-well known activation of the antiporter. Having overcome the difficulty of obtaining reproducible baseline fluorescence and established that, in our cultures, the  $\text{Na}^+/\text{H}^+$  antiport indeed plays an important role in regulating  $\text{pH}_i$  in hepatocytes (Fig. 5), the direct effect of liver tumor promoters on hepatocytes  $\text{pH}_i$  was explored. Our results consistently showed that PB and HCH caused an elevation of  $\text{pH}_i$  (0.05–0.08 pH units) over that in control cells, and that the elevation of  $\text{pH}_i$  was amiloride analogue-sensitive (Fig. 7). Our preliminary data showed that other liver tumor promoters (eg., DDT and PCN) which stimulated DNA synthesis, also induced cytoplasmic alkalization in hepatocytes (data not shown). In contrast, the estrogen groups of tumor promoters (Lee and Edwards, 2001a) had no effect on  $\text{pH}_i$  of hepatocytes (data not shown). This is consistent with the view that estrogenic liver tumor promoters act via distinct mechanisms in stimulating hepatocytes DNA synthesis (Lee and Edwards, 2001a).

The exact molecular events underlying the activation of  $\text{Na}^+/\text{H}^+$  antiporter by the various liver tumor promoters (PB, HCH, DDT, PCN) are currently not known. These xenobiotics of diverse structures could directly or indirectly affect activity of the antiporter. Other signaling mechanisms, involving mediators such as protein kinase C (Gillies et al., 1991; Ho et al., 1989; Lowe et al., 1990; Tanaka et al., 1994), cyclic AMP (Moule and McGivan, 1990), intracellular  $\text{Ca}^{2+}$  (Little et al., 1988; Alvarez et al., 1989; Hendey et al., 1989) and tyrosine kinase (Moule and McGivan, 1990; Tanaka et al., 1994) have been shown to increase  $\text{Na}^+/\text{H}^+$  antiport activity in different cell types. While the effects of TPA in liver cells presumably involve activation of protein kinase C, liver tumor promoters had no effect on protein kinase C activity and do not elevate cyclic AMP and  $\text{Ca}^{2+}$  levels in hepatocytes (Yusof and Edwards, results not shown), thereby, ruling out the possible role on these signaling mechanisms as mediators. Further studies will be required to determine whether tyrosine kinase activation or any other mechanisms mediate the action of liver tumor promoters in activating the  $\text{Na}^+/\text{H}^+$  antiporter in hepatocytes. Future studies will also include examination on the effect of liver tumor promoters on individual hepatocytes rather than the entire population of isolated hepatocytes as described in this study.

In conclusion, this study has provided for the first time evidence that hepatocyte DNA synthesis is dependent on  $\text{pH}_i$  and that elevation in  $\text{pH}_i$  is at least permissive in the stimulation of DNA synthesis. Some liver tumor promoters were found to elevate  $\text{pH}_i$  of isolated hepatocytes within minutes via activation of  $\text{Na}^+/\text{H}^+$

antiport. Taken together, results from this study suggest that activation of  $\text{Na}^+/\text{H}^+$  antiport may be one mechanism by which some liver tumor promoters stimulate growth in hepatocytes.

### ACKNOWLEDGMENT

Chow H. Lee was a recipient of Flinders University Research Scholarship.

### LITERATURE CITED

- Alvarez J, Garcia-Sancho J, Mollinedo F, Sanchez A. 1989. Intracellular  $\text{Ca}^{2+}$  potentiates  $\text{Na}^+/\text{H}^+$  exchange and cell differentiation induced by phorbol ester in U937 cells. *Eur J Biochem* 183: 709–714.
- Anwer MS, Nolan K. 1988. Characterization of  $\text{H}^+$  efflux pathways in rat hepatocytes. *Hepatology* 8:728–734.
- Arias IM, Forgac M. 1984. The sinusoidal domain of the plasma membrane of rat hepatocytes contains amiloride-sensitive  $\text{Na}^+/\text{H}^+$  antiport. *J Biol Chem* 259:5406–5408.
- Arkhammar P, Nilsson T, Berggren PO. 1989. Glucose-stimulated efflux of Fura-2 in pancreatic  $\beta$ -cells is prevented by probenecid. *Biochem Biophys Res Commun* 159:223–228.
- Arkhammar P, Nilsson T, Berggren PO. 1990. Glucose-stimulated efflux of indo-1 from pancreatic  $\beta$ -cells is reduced by probenecid. *FEBS Letts* 273:182–184.
- Burton KA. 1956. A study of the conditions and mechanism of the diphenylamine reaction for the colorimetric estimation of DNA. *Biochem J* 62:315–322.
- Church JG, Mills GB, Buick RN. 1989. Activation of the  $\text{Na}^+/\text{H}^+$  antiport is not required for epidermal growth factor-dependent gene expression, growth inhibition, or proliferation in human breast cancer cells. *Biochem J* 257:151–157.
- Di Sario A, Baroni GS, Bendia E, Ridolfi F, Saccomanno S, Ugili L, Trozzi L, Marziani M, Jezequel AM, Macarri G, Benedetti A. 2001. Intracellular pH regulation and  $\text{Na}^+/\text{H}^+$  exchange activity in human stellate cells: Effect of platelet-derived growth factor, insulin-like growth factor an insulin. *J Hepatol* 34:378–385.
- Di Virgilio F, Steinberg TH, Swanson JA. 1988. Fura-2 secretion and sequestration in macrophages. *J Immunol* 140:915–920.
- Frelin C, Vigne P, Ladoux A, Lazdunski M. 1988. The regulation of the intracellular pH in cells from vertebrates. *Eur J Biochem* 174: 3–14.
- Ganz MB, Boyarsky G, Sterzel RB, Boron WF. 1989. Arginine vasopressin enhances  $\text{pH}_i$  regulation in the presence of  $\text{HCO}_3^-$  by stimulating three acid-base transport systems. *Nature* 337:648–651.
- Garcia-Canero R, Trilla C, Perez de Diego J, Diaz-Gill JJ, Cobo JM. 1999.  $\text{Na}^+/\text{H}^+$  exchange inhibition induces intracellular acidosis and differentially impairs cell growth and viability of human and rat hepatocarcinoma cells. *Toxicol Letts* 106:215–228.
- Gillies RJ, Martinez R, Sneider JM, Hujer PB. 1991. Sphingosine inhibits phorbol 12-myristate 13-acetate-, but not serum-induced, activation of  $\text{Na}^+/\text{H}^+$  exchange in mammalian cells. *J Cell Physiol* 139:125–130.
- Gleeson D, Smith ND, Boyer JL. 1989. Bicarbonate-dependent and -independent intracellular cellular pH regulatory mechanisms in rat hepatocytes. *J Clin Invest* 84:312–321.
- Goodrich AL, Suchy FJ. 1990.  $\text{Na}^+/\text{H}^+$  exchange in basolateral plasma vesicles from neonatal rat liver. *Am J Physiol* 259:G334–G339.
- Grinstein S, Rotin D, Mason MJ. 1989.  $\text{Na}^+/\text{H}^+$  exchange and growth factor-induced cytosolic pH changes. Role in cellular proliferation. *Biochim Biophys Acta* 988:73–97.
- Hendey B, Mamrack MD, Putnam RW. 1989. Thrombin induces a calcium transient that mediates an activation of the  $\text{Na}^+/\text{H}^+$  exchanger in human fibroblasts. *J Biol Chem* 264:19540–19547.
- Ho AK, Chik CL, Weller JL, Cragoe EJ, Jr., Klein DC. 1989. Evidence of  $\beta$ 1-adrenergic protein kinase C  $\text{Na}^+/\text{H}^+$  antiporter-dependent increase in pinealocyte intracellular pH. *J Biol Chem* 264:12983–12988.
- Isom LL, Cragoe EJ, Jr., Limbird LE. 1987.  $\alpha_2$ -adrenergic receptors accelerate  $\text{Na}^+/\text{H}^+$  exchange in neuroblastoma  $\times$  Glioma cells. *J Biol Chem* 262:6750–6757.
- Johnson JD, Epel D, Paul M. 1976. Intracellular pH and activation of sea urchin eggs after fertilization. *Nature* 262:661–664.
- Kaneko A, Hayashi N, Tanaka Y, Horimoto M, Ito T, Sasaki Y, Fusamoto H, Kamada T. 1995. Activation of  $\text{Na}^+/\text{H}^+$  exchanger by hepatocyte-growth factor in hepatocytes. *Hepatology* 2:629–636.
- Koch KS, Leffert HL. 1979. Increased sodium ion influx is necessary to initiate rat hepatocyte proliferation. *Cell* 18:153–163.
- Koch KS, Leffert H. 1980. Growth control of differentiated adult rat hepatocytes in primary culture. In: Borek C, Williams GM, editors. *Differentiation and carcinogenesis in liver cell cultures*. New York: New York Acad Sci p 111–127.
- Kruijer W, Skelly H, Botter F, van der Putten H, Barber JR, Verma IM, Leffert HL. 1986. Proto-oncogenes expression in regenerating liver is stimulated in cultures of primary adult rat hepatocytes. *J Biol Chem* 261:7929–7933.
- Lee CH, Edwards AM. 2001a. Stimulation of DNA synthesis and *c-fos* mRNA expression in primary rat hepatocytes by estrogens. *Carcinogenesis* 22:1473–1481.
- Lee CH, Edwards AM. 2001b. Stimulation of DNA synthesis by tumor promoters in primary rat hepatocytes is not mediated by arachidonic acid metabolites. *J Cell Physiol* 187:336–344.
- Lee CH, Edwards AM. 2003. Differential of *c-fas* and *c-myc* proto-oncogenes by estrogens, xenobiotics, and other growth-stimulatory agents in primary rat hepatocytes. *Arch Toxicol* (in press).
- Leffert HL, Koch KS, Lu X-P, Brenner DA, Karin M, Skelly HF, Rippe RA, Fey G, Chojkier M. 1990. Cellular and molecular biology of hepatocyte growth, regeneration, and gene expression. In: Nishizuka Y, editor. *The biology and medicine of signal transduction*. New York: Raven Press. p 325–358.
- Little PJ, Weissberg PL, Cragoe EJ, Jr., Bobik A. 1988. Dependence of  $\text{Na}^+/\text{H}^+$  antiport activation in cultured rat aortic smooth muscle on calmodulin, calcium, and ATP. *J Biol Chem* 263:16700–16786.
- Lowe JHN, Huang C-L, Ives HE. 1990. Sphingosine differentially inhibits activation of the  $\text{Na}^+/\text{H}^+$  exchanger by phorbol esters and growth factors. *J Biol Chem* 265:7188–7194.
- Madhus IH. 1988. Regulation of intracellular pH in eukaryotic cells. *Biochem J* 250:1–8.
- McDonough PM, Button DC. 1989. Measurement of cytoplasmic  $\text{Ca}^{2+}$  concentration in cell suspensions: Correction for extracellular Fura-2 through use of  $\text{Mn}^{2+}$  and probenecid. *Cell Calcium* 10: 171–180.
- Michalopoulos GK, Pitot HC. 1975. Primary culture of parenchymal liver cells on collagen membrane. *Exp Cell Res* 94:70–78.
- Mills GB, Cragoe EJ, Jr., Gelfand EW, Grinstein S. 1985. Interleukin-2 induces a rapid increase in intracellular pH through activation of a  $\text{Na}^+/\text{H}^+$  antiport. *J Biol Chem* 260:12500–12507.
- Moolenaar WH, Tsien RY, Van der Saag PT, de Laat SW. 1983.  $\text{Na}^+/\text{H}^+$  exchange and cytoplasmic pH in the action of growth factors in human fibroblasts. *Nature* 304:645–648.
- Moolenaar WH, Tertoolen LGJ, de Laat SW. 1984. The regulation of cytoplasmic pH in human fibroblasts. *J Biol Chem* 259:7563–7569.
- Moolenaar WH, Defize LHK, de Laat SW. 1986. Ionic signalling by growth factor receptors. *J Exp Biol* 124:359–373.
- Moseley RH, Barrett C, Boyer JL. 1986a.  $\text{Na}^+/\text{H}^+$  exchange activity is enhanced in liver plasma membrane vesicles derived from partially hepatectomized rats. *Gastroenterology* 90:1749.
- Moule SK, McGivan JD. 1990. Epidermal growth factor and cyclic AMP stimulate  $\text{Na}^+/\text{H}^+$  exchange in isolated rat hepatocytes. *Eur J Biochem* 187:677–682.
- Ochsner M, Creba J, Walker J, Bentley P, Muakkassah-Kelly SF. 1990. Nafenopin, a hypolipidemic and non-genotoxic hepatocarcinogen increases intracellular calcium and transiently decreases intracellular pH in hepatocytes without generation of inositol phosphates. *Biochem Pharmacol* 40:2247–2257.
- Oliver IT, Edwards AM, Pitot HC. 1978. Hormonal regulation of phosphoenolpyruvate carboxykinase in primary cultures of adult rat liver parenchymal cells. *Eur J Biochem* 87:221–227.
- Petronijevic T, Edwards AM. 1993. Role of  $\text{Ca}^{2+}$  in stimulation of DNA synthesis by epidermal growth factor and tumor promoters in cultured rat hepatocytes. *J Cell Physiol* 155:44–53.
- Petzinger E, Follman W, Acker H, Hentchel J, Zierold K, Kinne RKH. 1988. Primary liver cell cultures grown on gas permeable membrane as source for the collection of primary bile. *In Vitro Cell Develop Biol* 24:491–499.
- Pollock AS. 1984. Intracellular pH of hepatocytes in primary monolayer culture. *Am J Physiol* 246:F738–F744.
- Pressman BC. 1976. Biological applications of ionophores. *Ann Rev Biochem* 45:501–530.
- Quinn DA, Dahlberg CGW, Bonventre JP, Scheid CR, Honeyman T, Joseph PM, Thompson BT, Hales CA. 1996. The role of  $\text{Na}^+/\text{H}^+$



- exchange and growth-factors in pulmonary-artery smooth-muscle cell-proliferation. *Am J Resp Cell Mol Biol* 14:139–145.
- Renner EL, Lake JR, Persico M, Scharschmidt BF. 1989. Na<sup>+</sup>-H<sup>+</sup> exchange activity in rat hepatocytes: Role in regulation of intracellular pH. *Am J Physiol* 256:G44–G52.
- Reshkin SJ, Bellizzi A, Caldeira S, Albarani V, Malanchi I, Poignee M, AlunniFabbioni M, Casavola V, Tommasino M. 2000. Na<sup>+</sup>/H<sup>+</sup> exchanger-dependent intracellular alkalinization is an early event in malignant transformation and plays an essential role in the development of subsequent transformation-associated phenotypes. *FASEB J* 14:2185–2197.
- Simchowicz L, Cragoe EJ, Jr. 1986. Inhibition of chemotactic factor-activated Na<sup>+</sup>/H<sup>+</sup> exchange in human neutrophils by analogues of amiloride: Structure-activity relationships in the amiloride series. *Mol Pharmacol* 30:112–120.
- Smith ND, Boyer JL. 1986. Phorbol esters stimulate Na-H exchange in isolated rat hepatocytes. *Hepatology* 6:1170.
- Soltoff SP, Cantley LC. 1988. Mitogens and ion fluxes. *Ann Rev Physiol* 50:207–223.
- Strazzabosco M, Poci C, Spirli C, Zsembery A, Granato A, Massimino ML, Crepaldi G. 1995. Intracellular pH regulation in HepG2 cells—effects of epidermal growth factor, transforming growth factor alpha, and insulin-like growth factor-II on Na<sup>+</sup>/H<sup>+</sup> exchange activity. *Hepatology* 22:588–597.
- Tanaka Y, Hayashi N, Kaneko A, Ito T, Horimoto M, Sasaki Y, Kasahara A, Fusamoto H, Kamada T. 1994. Characterization of signaling pathways to Na<sup>+</sup>/H<sup>+</sup> exchanger activation with epidermal growth factor in hepatocytes. *Hepatology* 20:966–974.
- Thomas JA, Buchsbaum RN, Zimnick A, Raeker C. 1979. Intracellular pH measurements in Ehrlich tumor cells utilizing spectroscopic probes generated in situ. *Biochemistry* 18:2210–2218.
- Whiting MJ, Edwards AM. 1979. Measurements of cholic acid synthesis and secretion by isolated rat hepatocytes. *J Lipid Res* 20:914–918.
- Williamson JR, Meyer AJ, Ohkawa K. 1974. Interrelations between anion transport, ureogenesis and gluconeogenesis in rat liver cells. In: Lingquist F, Tygstrup N, editors. *Regulation of hepatic metabolism*. New York: Academic Press. p 45–479.
- Yusof YAM, Edwards AM. 1990. Stimulation of DNA synthesis in primary rat hepatocyte cultures by liver tumor promoters: Interaction with other growth factors. *Carcinogenesis* 11:761–770.