



Role of mineralocorticoid receptor in insulin resistance

Rajesh Garg and Gail K. Adler

Purpose of review

Recent data suggest that mineralocorticoid receptor activation can affect insulin resistance independent of its effects on blood pressure. This review discusses new evidence linking mineralocorticoid receptor to insulin resistance and the underlying mechanisms of these effects.

Recent findings

Observational studies have shown mineralocorticoid activity to be associated with insulin resistance irrespective of race, blood pressure or body weight. Increased mineralocorticoid activity may be the common link between obesity, hypertension, dyslipidemia and insulin resistance, features that make up the metabolic syndrome. Treatment of primary aldosteronism is associated with a decrease in insulin resistance and provides one of the most convincing evidences in favor of the contribution of mineralocorticoid receptor to insulin resistance. Dietary salt restriction, which increases aldosterone levels, is also associated with an increase in insulin resistance. Potential mechanisms by which mineralocorticoid receptor may contribute to insulin resistance include a decreased transcription of the insulin receptor gene, increased degradation of insulin receptor substrates, interference with insulin signaling mechanisms, decreased adiponectin production and increased oxidative stress and inflammation. Advantages of mineralocorticoid receptor antagonists on insulin resistance have been demonstrated in animal models.

Summary

There may be a benefit of mineralocorticoid receptor antagonists in human insulin resistance states, but more clinical research is needed to explore these possibilities.

Keywords

aldosterone, insulin resistance, mineralocorticoid receptor, obesity

INTRODUCTION

Traditionally, mineralocorticoid effects have been linked to fluid and electrolyte balance with blockade of the mineralocorticoid receptor reducing blood pressure (BP) in individuals with hypertension. Recent studies have shown mineralocorticoid receptor antagonists to improve clinical outcomes in acute myocardial infarction with heart failure, chronic heart failure and diabetic nephropathy [1–5]. These effects of mineralocorticoid receptor blockade are not dependent on BP regulation, but involve beneficial effects of mineralocorticoid receptor blockade on vascular function/injury, inflammation, cardiac fibrosis and renal damage. Accumulating data also suggest a key role of the mineralocorticoid receptor in metabolic regulation. This article reviews scientific evidence linking mineralocorticoids to glucose metabolism and insulin resistance with special emphasis on the underlying mechanisms of these effects.

OBSERVATIONAL STUDIES LINKING MINERALOCORTICIDS AND INSULIN RESISTANCE

Observational studies have linked mineralocorticoid receptor to insulin resistance in many different conditions including obesity, hypertension, metabolic syndrome, primary aldosteronism and also in normal healthy individuals.

Division of Endocrinology, Diabetes and Hypertension, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA

Correspondence to Gail K. Adler, MD, PhD, Associate Professor of Medicine, Division of Endocrinology, Diabetes and Hypertension, 221 Longwood Avenue, Boston, MA 02115, USA. Tel: +1 617 732 5666; fax: +1 617 732 5764; e-mail: gadler@partners.org

Curr Opin Endocrinol Diabetes Obes 2012, 19:168–175

DOI:10.1097/MED.0b013e3283533955

KEY POINTS

- High aldosterone levels are present in human insulin resistant states and correlate with measures of insulin resistance.
- Mineralocorticoid receptor activity may interfere with insulin signaling, directly or via an increase in oxidative stress and inflammation.
- Advantages of mineralocorticoid receptor antagonists have been demonstrated in animal models but remain to be proven in human insulin resistance states.

OBESITY, MINERALOCORTICOIDS AND INSULIN RESISTANCE

The relationship between mineralocorticoids and insulin resistance has been reported for almost 2 decades beginning with the observation that obese individuals often have high BP, activated renin angiotensin aldosterone system (RAAS) and insulin resistance [6]. Urinary aldosterone excretion and angiotensin II-stimulated aldosterone levels are higher in overweight individuals and aldosterone production correlates with insulin resistance suggesting a potential role for aldosterone in the pathophysiology of obesity-mediated insulin resistance [7]. Weight loss by low calorie diet leads to an improvement in insulin sensitivity as well as a decrease in BP [8], and these changes are associated with reversal of the RAAS overactivity [9]. Similarly, weight loss after gastric bypass surgery lowers RAAS activity and improves insulin resistance [10], whereas weight loss after gastric banding has been demonstrated to reduce aldosterone levels along with a reduction in BP and insulin resistance [11].

The mechanisms of increased mineralocorticoid activity in obesity are not established. According to one hypothesis, fatty acids from visceral adipocytes may induce hepatic formation of an adrenal secretagogue that leads to increased aldosterone production [12]. Alternatively, there may be a direct effect of lipids on aldosterone as a lipid infusion was shown to increase aldosterone along with an increase in insulin resistance [13[■]]. Finally, adipocytes may produce factors that directly activate mineralocorticoid receptor [14[■]].

HYPERTENSION, MINERALOCORTICOIDS AND INSULIN RESISTANCE

Patients with essential hypertension have higher aldosterone levels than normotensive patients and aldosterone may be one of the mediators of increased insulin resistance and dyslipidemia in these patients as suggested by a positive correlation

between plasma aldosterone and measures of insulin resistance [15,16]. Thus, aldosterone may contribute to not only the maintenance of high BP but also insulin resistance and increased cardiovascular risk in patients with essential hypertension [17,18].

MINERALOCORTICOIDS AND INSULIN RESISTANCE IN HEALTHY INDIVIDUALS

Aldosterone production is also a predictor of insulin sensitivity in normal healthy individuals. This was demonstrated in a well controlled physiological study wherein angiotensin II-stimulated aldosterone predicted 8% of the variance in insulin sensitivity, which is almost one-third of that predicted by age, BMI and diastolic BP together (23%) [19[■]]. In a recent study involving 483 African-Americans without any cardiovascular or renal disease, aldosterone levels were predictive of homeostatic model assessment (HOMA)-insulin resistance independent of all other variables [20]. In another recent study of 1088 individuals plasma aldosterone levels predicted insulin resistance [21[■]]. Further, when 564 of these individuals were followed for 10 years, insulin resistance developed more often in individuals with high baseline plasma aldosterone than in those with low baseline plasma aldosterone [21[■]].

MINERALOCORTICOIDS AND THE METABOLIC SYNDROME

Thus, increased aldosterone is observed in obesity and in hypertension. In both conditions, as well as in healthy individuals, aldosterone is associated with insulin resistance. On the basis of these observations and on cellular and animal studies described below, the hypothesis has been proposed that aldosterone is a pathological link between obesity, insulin resistance, hypertension and dyslipidemia, features that make up the metabolic syndrome (MetS) [22,23]. This concept is supported by data from the Framingham Offspring Study that showed high circulating aldosterone levels to be associated with the development of MetS, in a predominately white population [24]. Further, African-American hypertensive men with MetS had high plasma aldosterone that was predictive of HOMA-insulin resistance [25,26]. These data suggest that aldosterone has a role in the causation of insulin resistance irrespective of race, body weight and BP.

GENE POLYMORPHISMS, MINERALOCORTICOIDS AND INSULIN RESISTANCE

Gene polymorphism studies provide another way of exploring the relationship between aldosterone and

insulin resistance. Genetic variations in aldosterone synthase gene (CYP11B2) that are associated with increased aldosterone are predictive of high plasma glucose levels and insulin resistance [27,28]. These gene polymorphisms are also associated with increased risk of type 2 diabetes, hypertension and the MetS [29]. Similarly, an angiotensinogen gene polymorphism that is associated with increased RAAS activity is associated with high fetal total glycated hemoglobin, a surrogate of fetal insulin resistance at birth [30]. Another relation between insulin activity and mineralocorticoids is shown by the hepatic nuclear factor (HNF)1A gene mutation that causes type 3 maturity-onset diabetes of the young, the most common single gene mutation causing type 2 diabetes mellitus. HNF1A-null mice show an overactive adrenal gland with increased aldosterone levels [31].

PRIMARY ALDOSTERONISM AND INSULIN RESISTANCE

Studies on patients with primary aldosteronism provide the most compelling evidence about the role of aldosterone in insulin resistance in humans. Patients with primary aldosteronism have increased insulin resistance and impaired glucose metabolism [32] and serum aldosterone levels correlate with insulin resistance [33]. Removal of aldosterone excess by surgery or by medical therapy improves insulin sensitivity, implicating aldosterone as a main determinant of the alteration in insulin sensitivity [33–35]. Low adiponectin levels have also been described in primary aldosteronism and seem to be the result of aldosterone excess because therapeutic interventions to correct the aldosterone effect reverse these changes [36]. Also, patients with primary aldosteronism in the presence of specific adiponectin gene polymorphisms are more prone to develop MetS [37]. Recently, nonalcoholic fatty liver disease (NAFLD), a known complication of insulin resistance, was found to be more frequent in patients with primary aldosteronism, further indicating a role for aldosterone in inducing insulin resistance [38]. A small study showed that treatment with spironolactone decreases insulin levels and improves insulin sensitivity in NAFLD [39].

LOW SALT DIET AND INSULIN RESISTANCE

Additional evidence in favor of aldosterone as a mediator of insulin resistance comes from studies involving manipulations of dietary sodium intake. It is well known that as individuals transition from a liberal to a low salt diet there is activation of RAAS

activity. Recent studies demonstrated an increase in insulin resistance in individuals on a low salt diet as compared with a liberal salt diet [40,41]. Further, salt sensitivity of BP, due to insufficient suppression of aldosterone production on a liberal salt diet is strongly associated with insulin resistance, which again suggests a role for aldosterone in insulin resistance [42].

Fasting insulin, triglycerides, inflammatory markers and the insulin response to oral glucose, all rise significantly after dietary salt restriction raising the possibility of an effect of RAAS activation on these metabolic parameters [43–45]. Adiponectin levels are lower on lower salt diet as compared with high salt diet [46,47]. In this context, recent studies showing increased mortality in association with low salt intake [48–51] present an interesting argument against the prevailing views about the benefits of low salt diet [52].

POTENTIAL MECHANISMS OF MINERALOCORTICOID INDUCED INSULIN RESISTANCE

A number of in-vitro and in-vivo studies have explored the mechanisms of mineralocorticoid-mediated insulin resistance. These mechanisms are summarized in Fig. 1.

MINERALOCORTICOIDS AND INSULIN RECEPTOR LEVELS

Treatment of human promonocytic cells with aldosterone leads to decreases in glucose uptake and reductions in insulin receptor mRNA levels due to a decrease in insulin receptor gene transcription [53,54]. Addition of a mineralocorticoid receptor antagonist prevents these changes in insulin receptor mRNA levels [53,54]. Further, incubation of 3T3-L1 adipocytes with aldosterone leads to increased degradation of insulin receptor substrate (IRS) 1 and IRS2 [55]. These data suggest an effect of mineralocorticoid receptor on gene transcription and stability of factors involved in the insulin signaling pathways.

MINERALOCORTICOIDS AND OXIDATIVE STRESS

Another suggested mechanism underlying the interaction between mineralocorticoid receptor and insulin signaling may involve reactive oxygen species (ROS). Aldosterone inhibits expression of uncoupling protein-1 (UCP-1) [56] and decreases in UCP-1 are associated with increased production of ROS. In 3T3-L1 adipocytes, activation of




FIGURE 1. Potential mechanisms of mineralocorticoid receptor induced insulin resistance. GLUT, glucose transported; IRS, insulin receptor substrate; MR, mineralocorticoid receptor; ROS, reactive oxygen species.

mineralocorticoid receptor by aldosterone increases intracellular ROS levels [57] and these increases in ROS contribute to the degradation of IRS protein [55,58]. Obesity is a state of oxidative stress due to increased ROS generation by circulating leukocytes as well as adipocytes [59]. Treatment of obese ob/ob and db/db mice with the mineralocorticoid antagonist, eplerenone reduces insulin resistance along with suppression in adipose tissue of macrophage infiltration and ROS production [57]. These results suggest that mineralocorticoid receptor-mediated insulin resistance is the result at least in part of activation of oxidative stress.

MINERALOCORTICOIDS AND INFLAMMATION

Inflammatory pathways are known to affect insulin signaling and induce insulin resistance. It is well established from in-vivo studies, that activation of mineralocorticoid receptor increases inflammation, and mineralocorticoid receptor blockade reduces inflammation, in the vasculature, heart, kidney and adipose tissue [22,60]. There appear to be direct effects of mineralocorticoid receptor on inflammatory molecules, as incubation of 3T3-L1 preadipocytes and adipocytes with aldosterone increases mRNA expression of multiple inflammatory factors including monocyte chemoattractant protein-1 (MCP-1), tumor necrosis factor (TNF) α and plasminogen activator inhibitor [61]. By analogy to the effects of TNF- α on inflammation and insulin resistance, some investigators have suggested that induction of inflammation may be the underlying mechanism linking aldosterone to insulin resistance [62]. Consistent with this hypothesis, treatment of obese mice with eplerenone reduces expression

of proinflammatory and prothrombotic factors in adipose tissue and increases expression of adiponectin in heart and adipose tissue [61]. These effects of mineralocorticoid receptor blockade on adiponectin and adipokine gene expression may influence insulin resistance.

The relationship between mineralocorticoid receptor, inflammation, oxidative stress and insulin resistance has been studied in the deoxycorticosterone acetate (DOCA) and salt hypertensive rat model. These animals have mineralocorticoid receptor activation resulting in increased oxidative stress, inflammation and insulin resistance; treatment with hemin reduces oxidative stress and inflammation and improves insulin sensitivity [63]. Further, treatment of Zucker diabetic-fatty rat with hemoxygenase-1 reduces aldosterone, reduces markers of oxidative stress and inflammation (e.g. nuclear factor- κ B, AP-1, and AP-2), increases AMP-activated protein kinase and increases glucose transported (GLUT4) expression [64].

MINERALOCORTICOIDS AND OTHER INSULIN SIGNALING PATHWAYS

The transgenic Ren2 rat manifests increased tissue RAAS with elevated serum aldosterone and is both hypertensive and insulin resistant with reduced muscle IRS-1 tyrosine phosphorylation, Akt phosphorylation/activation, and GLUT4 expression [65]. These findings suggest that aldosterone contributes to insulin resistance in the transgenic Ren2 rats by interfering with insulin signaling pathways [65]. Similarly, rats injected with aldosterone daily for 15 days showed decreased transcription of GLUT2 and GLUT4 genes and decreased translocation of GLUT4 to the plasma membrane in liver and

skeletal muscle [66]. In DOCA implanted hypertensive rats, GLUT4 levels were depressed several-fold in aortae and carotid arteries compared with sham-treated normotensive rats and uptake of the glucose analog, 2-deoxyglucose (2-DOG), was reduced by 53% [67].

Another mechanism that may involve overlap between mineralocorticoid receptor and insulin receptor signaling is related to RAS-related C3 botulinum substrate 1 (RAC1) pathways. RAC1 is able to directly activate mineralocorticoid receptor even in the absence of mineralocorticoids or glucocorticoids. Overexpression of circulating peptide coupling factor 6 (CF6) in mice increases renal RAC-1 leading to activation of renal mineralocorticoid receptor [68]. These mice also have defects in insulin signaling mechanisms in the skeletal muscle and the liver resulting in a decrease of plasma membrane-bound GLUT4 leading to insulin resistance [68].

MINERALOCORTICOIDS AND ADIPOCYTE DIFFERENTIATION

Aldosterone and mineralocorticoid receptor may also be involved in adipocyte biology at the more basic level. Differentiation of 3T3-L1 preadipocytes into adipocytes is dependent on the presence of mineralocorticoid receptor. Using siRNA technology to reduce mineralocorticoid receptor and glucocorticoid receptor levels and pharmacological inhibition of mineralocorticoid receptor and glucocorticoid receptor, investigators demonstrated that this differentiation is dependent on activation of the mineralocorticoid receptor by either glucocorticoids or mineralocorticoids [69]. Further, the aldosterone effect on adipose maturation is accompanied by induction of PPAR- γ expression [69]. These results suggest that mineralocorticoid receptor may be an important proadipogenic transcription factor that may mediate both aldosterone and glucocorticoid effects on adipose tissue development [69]. These studies provide further mechanistic support for the concept that mineralocorticoid receptor may be involved in the pathophysiology of obesity and MetS.

MINERALOCORTICOIDS AND INSULIN SECRETION

There is also a school of thought that mineralocorticoids impair insulin secretion, an effect that has been attributed to aldosterone induced hypokalemia. Recently, aldosterone was shown to suppress insulin secretion via activation of ROS [70]. This effect of aldosterone was independent of

potassium and angiotensin II and was not mediated through mineralocorticoid receptor [70]. This effect in addition to aldosterone mediated increases in insulin resistance will further impair glucose metabolism.

MINERALOCORTICOID RECEPTOR ANTAGONISTS IN ANIMAL MODELS OF INSULIN RESISTANCE AND METABOLIC SYNDROME

Mineralocorticoid receptor antagonists reverse many of the changes associated with insulin resistance and metabolic syndrome. In a mouse model of diet-induced diabetes and NAFLD, the mineralocorticoid receptor blocker, spironolactone treatment improved glucose tolerance and hepatic mRNA expression of proinflammatory cytokines (TNF α , IL-6, and MCP-1) without any effect on calorie intake or body weight [71[■]]. In a rat model of RAAS activation, low-dose spironolactone improved IRS-1/Akt phosphorylation in renal cortical tissue and reduced albuminuria [72]. Mineralocorticoid receptor blockade reduces renal injury in rodent models of type 1 and 2 diabetes [73] and reduces aortic superoxide production and improves endothelium-dependent vasorelaxation in rodent models of type 1 diabetes [74]. Treatment with mineralocorticoid receptor antagonist also improved muscle insulin signaling parameters and systemic insulin sensitivity in association with reductions in NADPH oxidase activity and ROS production in transgenic Ren2 rats [65]. These results indicate that inhibition of the mineralocorticoid effect might be a beneficial therapeutic approach for multiple phenotypes of insulin resistance.

THERAPEUTIC CONSIDERATIONS

Metabolic dysregulation is common in conditions like diabetes, obesity, polycystic ovarian syndrome (PCOS), hypertension, and heart failure and when present contributes to the pathophysiology of these disorders. Given our increasing understanding of the relevance of aldosterone to the pathophysiology of hypertension, vascular and target organ damage, mineralocorticoid receptor blockade is being used with increasing frequency in the treatment of patients with hypertension and heart failure. With new data implicating the mineralocorticoid receptor in adipocyte biology and glucose metabolism, it may be worth considering the use of mineralocorticoid receptor antagonists in treatment of disorders with metabolic dysregulation.

High aldosterone levels along with high androgen levels are present in women with PCOS [75] and

treatment of PCOS with metformin improves insulin sensitivity along with a reduction in aldosterone and androgen levels [76]. Chlorthalidone, which tends to activate the RAAS, causes an increase in insulin resistance in hypertensive patients that is avoided by addition of spironolactone [77]. Treatment with spironolactone was recently demonstrated to decrease insulin levels in NAFLD [39^o]. Multiple studies have suggested that interruption of the RAAS with angiotensin inhibition can improve insulin sensitivity and potentially reduce the incidence of diabetes [78]. As aldosterone escape is a known phenomenon in patients treated with an angiotensin converting enzyme inhibitor or angiotensin receptor blocker, there may be a role for the use of mineralocorticoid receptor antagonists in addition to these drugs. The largest clinical trials with the use of mineralocorticoid receptor antagonists have been conducted in heart failure [1,2,4], but we were unable to find any literature related to insulin resistance in these trials.

CONCLUSION

Thus, available data suggest a role of mineralocorticoid receptor in the pathophysiology of insulin resistance. Although clinical data show a reduction in insulin resistance after treatment of primary aldosteronism, there is very little available data on the use of mineralocorticoid receptor antagonists for prevention or amelioration of insulin resistance in other clinical conditions. Given the increasing evidence for inappropriately elevated aldosterone levels in many common disorders associated with insulin resistance, for example, obesity and hypertension, this is an important area of future research that could lead to new therapies to prevent complications of insulin resistance like diabetes and cardiovascular disease.

Acknowledgements

None.

Conflicts of interest

There are no conflicts of interest.

REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

Additional references related to this topic can also be found in the Current World Literature section in this issue (p. 235).

1. Pitt B, Zannad F, Remme WJ, *et al.* The effect of spironolactone on morbidity and mortality in patients with severe heart failure. Randomized Aldosterone Evaluation Study Investigators. *N Engl J Med* 1999; 341:709–717.
2. Pitt B, Remme W, Zannad F, *et al.* Eplerenone, a selective aldosterone blocker, in patients with left ventricular dysfunction after myocardial infarction. *N Engl J Med* 2003; 348:1309–1321.
3. Mehdi UF, Adams-Huet B, Raskin P, *et al.* Addition of angiotensin receptor blockade or mineralocorticoid antagonism to maximal angiotensin-converting enzyme inhibition in diabetic nephropathy. *J Am Soc Nephrol* 2009; 20:2641–2650.
4. Zannad F, McMurray JJ, Krum H, *et al.* Eplerenone in patients with systolic heart failure and mild symptoms. *N Engl J Med* 2011; 364:11–21.
5. Epstein M, Williams GH, Weinberger M, *et al.* Selective aldosterone blockade with eplerenone reduces albuminuria in patients with type 2 diabetes. *Clin J Am Soc Nephrol* 2006; 1:940–951.
6. Licata G, Scaglione R, Ganguzza A, *et al.* Central obesity and hypertension. Relationship between fasting serum insulin, plasma renin activity, and diastolic blood pressure in young obese subjects. *Am J Hypertens* 1994; 7:314–320.
7. Bentley-Lewis R, Adler GK, Perlstein T, *et al.* Body mass index predicts aldosterone production in normotensive adults on a high-salt diet. *J Clin Endocrinol Metab* 2007; 92:4472–4475.
8. Ikeda T, Gomi T, Hirawa N, *et al.* Improvement of insulin sensitivity contributes to blood pressure reduction after weight loss in hypertensive subjects with obesity. *Hypertension* 1996; 27:1180–1186.
9. Fogari R, Zoppi A, Corradi L, *et al.* Effect of body weight loss and normalization on blood pressure in overweight nonobese patients with stage 1 hypertension. *Hypertens Res* 2010; 33:236–242.
10. Ruano M, Silvestre V, Castro R, *et al.* Morbid obesity, hypertensive disease and the renin-angiotensin-aldosterone axis. *Obes Surg* 2005; 15:670–676.
11. Dall'Asta C, Vedani P, Manunta P, *et al.* Effect of weight loss through laparoscopic gastric banding on blood pressure, plasma renin activity and aldosterone levels in morbid obesity. *Nutr Metab Cardiovasc Dis* 2009; 19:110–114.
12. Goodfriend TL, Egan BM, Kelley DE. Plasma aldosterone, plasma lipoproteins, obesity and insulin resistance in humans. *Prostaglandins Leukot Essent Fatty Acids* 1999; 60:401–405.
13. Florian JP, Pawelczyk JA. Sympathetic and haemodynamic responses to lipids ■ in healthy human ageing. *Exp Physiol* 2010; 95:486–497.
14. Nguyen Dinh Cat A, Briones AM, Callera GE, *et al.* Adipocyte-derived factors ■ regulate vascular smooth muscle cells through mineralocorticoid and glucocorticoid receptors. *Hypertension* 2011; 58:479–488.
- In this study, cultured mouse vascular smooth muscle cells (VSMCs) were exposed to adipocyte-conditioned medium (ACM) from differentiated 3T3-L1 adipocytes. ACM-induced phosphorylation of stress-activated protein kinase/c-Jun N-terminal kinase, p38MAPK, and extracellular signal-regulated kinase 1/2 and increased expression of proinflammatory and proliferative markers in VSMCs. Eplerenone (mineralocorticoid receptor antagonist), mifepristone (glucocorticoid receptor antagonist) and candesartan (AT1R antagonist) inhibited these effects. These findings suggest that adipocyte-derived factors regulate VSMC function through specific MAPKs linked to mineralocorticoid receptor, glucocorticoid receptor, and AT1R.
15. Lind L, Reneland R, Andersson PE, *et al.* Insulin resistance in essential hypertension is related to plasma renin activity. *J Hum Hypertens* 1998; 12:379–382.
16. Goodfriend TL, Egan B, Stepniakowski K, Ball DL. Relationships among plasma aldosterone, high-density lipoprotein cholesterol, and insulin in humans. *Hypertension* 1995; 25:30–36.
17. Colussi G, Catena C, Lapenna R, *et al.* Insulin resistance and hyperinsulinemia are related to plasma aldosterone levels in hypertensive patients. *Diabetes Care* 2007; 30:2349–2354.
18. Lind L, Lithell H, Wide L, Ljunghall S. Metabolic cardiovascular risk factors and the renin-aldosterone system in essential hypertension. *J Hum Hypertens* 1992; 6:27–29.
19. Garg R, Hurwitz S, Williams GH, *et al.* Aldosterone production and insulin ■ resistance in healthy adults. *J Clin Endocrinol Metab* 2010; 95:1986–1990.
- Significant negative correlations were present between insulin sensitivity index (ISI) measured by Matsuda index and age, BMI, diastolic BP and angiotensin II-stimulated aldosterone level. On multivariate regression analysis, stimulated aldosterone level was an independent predictor of ISI after adjusting for age, BMI and diastolic BP. Stimulated aldosterone level predicted 8% of the variance in ISI while age, BMI, and diastolic BP together predicted 23% of the variance in ISI.
20. Huan Y, Deloach S, Keith SW, *et al.* Aldosterone and aldosterone: renin ratio associations with insulin resistance and blood pressure in African Americans. *J Am Soc Hypertens* 2012; 6:56–65.
21. Kumagai E, Adachi H, Jacobs DR Jr, *et al.* Plasma aldosterone levels and ■ development of insulin resistance: prospective study in a general population. *Hypertension* 2011; 58:1043–1048.
- Plasma aldosterone levels and HOMA were measured in 1088 nondiabetic patients and 564 patients were followed prospectively to predict incident of insulin resistance. A significant cross-sectional relationship between plasma aldosterone and HOMA was present at baseline. In the prospective analysis, a significantly higher relative risk for the development of insulin resistance was observed in the highest tertile versus lowest tertile of plasma aldosterone, after adjustment for confounding factors. This 10-year prospective study demonstrated that plasma aldosterone levels predict the development of insulin resistance in the general population.

22. Tirosh A, Garg R, Adler GK. Mineralocorticoid receptor antagonists and the metabolic syndrome. *Curr Hypertens Rep* 2010; 12:252–257.
 23. Sowers JR, Whaley-Connell A, Epstein M. Narrative review: the emerging clinical implications of the role of aldosterone in the metabolic syndrome and resistant hypertension. *Ann Intern Med* 2009; 150:776–783.
 24. Ingelsson E, Pencina MJ, Toffler GH, *et al.* Multimarker approach to evaluate the incidence of the metabolic syndrome and longitudinal changes in metabolic risk factors: the Framingham Offspring Study. *Circulation* 2007; 116:984–992.
 25. Kidambi S, Kotchen JM, Grim CE, *et al.* Association of adrenal steroids with hypertension and the metabolic syndrome in blacks. *Hypertension* 2007; 49:704–711.
 26. Kidambi S, Kotchen JM, Krishnaswami S, *et al.* Hypertension, insulin resistance, and aldosterone: sex-specific relationships. *J Clin Hypertens (Greenwich)* 2009; 11:130–137.
 27. Ranade K, Wu KD, Risch N, *et al.* Genetic variation in aldosterone synthase predicts plasma glucose levels. *Proc Natl Acad Sci U S A* 2001; 98:13219–13224.
 28. Li XM, Ling Y, Lu DR, *et al.* Association of the aldosterone synthase gene-344T>C polymorphism with essential hypertension and glucose homeostasis: a case-control study in a Han Chinese population. *Clin Exp Pharmacol Physiol* 2011; 38:598–604.
 29. Bellili NM, Foucan L, Fumeron F, *et al.* Associations of the-344 T>C and the 3097 G>A polymorphisms of CYP11B2 gene with hypertension, type 2 diabetes, and metabolic syndrome in a French population. *Am J Hypertens* 2010; 23:660–667.
 30. Schlemm L, Haumann HM, Ziegner M, *et al.* New evidence for the fetal insulin hypothesis: fetal angiotensinogen M235T polymorphism is associated with birth weight and elevated fetal total glycated hemoglobin at birth. *J Hypertens* 2010; 28:732–739.
 31. Bonzo JA, Patterson AD, Krausz KW, Gonzalez FJ. Metabolomics identifies novel Hnf1alpha-dependent physiological pathways in vivo. *Mol Endocrinol* 2010; 24:2343–2355.
 32. Mosso LM, Carvajal CA, Maiz A, *et al.* A possible association between primary aldosteronism and a lower beta-cell function. *J Hypertens* 2007; 25:2125–2130.
 33. Giacchetti G, Ronconi V, Turchi F, *et al.* Aldosterone as a key mediator of the cardiometabolic syndrome in primary aldosteronism: an observational study. *J Hypertens* 2007; 25:177–186.
 34. Catena C, Lapenna R, Baroselli S, *et al.* Insulin sensitivity in patients with primary aldosteronism: a follow-up study. *J Clin Endocrinol Metab* 2006; 91:3457–3463.
 35. Sindelka G, Widimsky J, Haas T, *et al.* Insulin action in primary hyperaldosteronism before and after surgical or pharmacological treatment. *Exp Clin Endocrinol Diabetes* 2000; 108:21–25.
 36. Fallo F, Della Mea P, Sonino N, *et al.* Adiponectin and insulin sensitivity in primary aldosteronism. *Am J Hypertens* 2007; 20:855–861.
 37. Ronconi V, Turchi F, Rilli S, *et al.* Metabolic syndrome in primary aldosteronism and essential hypertension: relationship to adiponectin gene variants. *Nutr Metab Cardiovasc Dis* 2010; 20:93–100.
 38. Fallo F, Dalla Pozza A, Tecchio M, *et al.* Nonalcoholic fatty liver disease in primary aldosteronism: a pilot study. *Am J Hypertens* 2010; 23:2–5.
 39. Polyzos SA, Kountouras J, Zafeiriadou E, Patsiaoura K, Katsiki E, Deretzi G, *et al.* Effect of spironolactone and vitamin E on serum metabolic parameters and insulin resistance in patients with nonalcoholic fatty liver disease. *J Renin Angiotensin Aldosterone Syst* 2011; 12:498–503.
- This is a small randomized controlled trial of 8 weeks treatment with vitamin E versus spironolactone and vitamin E in biopsy-proven NAFLD patients. Insulin and HOMA decreased significantly in the spironolactone and vitamin E group as compared with vitamin E alone group.
40. Garg R, Williams GH, Hurwitz S, *et al.* Low-salt diet increases insulin resistance in healthy subjects. *Metabolism* 2011; 60:965–968.
- Healthy individuals were studied after 7 days of low salt diet (urine sodium <20 mmol/d) and 7 days of high salt diet (urine sodium >150 mmol/d). Mean HOMA was significantly higher on low salt compared with high salt diet.
41. Townsend RR, Kapoor S, McFadden CB. Salt intake and insulin sensitivity in healthy human volunteers. *Clin Sci (Lond)* 2007; 113:141–148.
 42. Yatabe MS, Yatabe J, Yoneda M, *et al.* Salt sensitivity is associated with insulin resistance, sympathetic overactivity, and decreased suppression of circulating renin activity in lean patients with essential hypertension. *Am J Clin Nutr* 2010; 92:77–82.
 43. Egan BM, Stepienkowski K, Goodfriend TL. Renin and aldosterone are higher and the hyperinsulinemic effect of salt restriction greater in subjects with risk factors clustering. *Am J Hypertens* 1994; 7 (10 Pt 1):886–893.
 44. Nakandakare ER, Charf AM, Santos FC, *et al.* Dietary salt restriction increases plasma lipoprotein and inflammatory marker concentrations in hypertensive patients. *Atherosclerosis* 2008; 200:410–416.
 45. Srikumar N, Brown NJ, Hopkins PN, *et al.* PAI-1 in human hypertension: relation to hypertensive groups. *Am J Hypertens* 2002; 15:683–690.
 46. Vaidya A, Forman JP, Underwood PC, *et al.* The influence of body mass index and renin-angiotensin-aldosterone system activity on the relationship between 25-hydroxyvitamin D and adiponectin in Caucasian men. *Eur J Endocrinol* 2011; 164:995–1002.
 47. Lely AT, Krikken JA, Bakker SJ, *et al.* Low dietary sodium and exogenous angiotensin II infusion decrease plasma adiponectin concentrations in healthy men. *J Clin Endocrinol Metab* 2007; 92:1821–1826.
 48. Ekinci EI, Clarke S, Thomas MC, *et al.* Dietary salt intake and mortality in patients with type 2 diabetes. *Diabetes Care* 2011; 34:703–709.
 49. Stolarz-Skrzypek K, Kuznetsova T, Thijs L, *et al.* Fatal and nonfatal outcomes, incidence of hypertension, and blood pressure changes in relation to urinary sodium excretion. *JAMA* 2011; 305:1777–1785.
 50. Thomas MC, Moran J, Forsblom C, *et al.* The Association Between Dietary Sodium Intake, ESRD, and All-Cause Mortality in Patients With Type 1 Diabetes. *Diabetes Care* 2011; 34:861–866.
 51. Taylor RS, Ashton KE, Moxham T, *et al.* Reduced dietary salt for the prevention of cardiovascular disease: a meta-analysis of randomized controlled trials (Cochrane review). *Am J Hypertens* 2011; 24:843–853.
 52. Appel LJ, Frohlich ED, Hall JE, *et al.* The importance of population-wide sodium reduction as a means to prevent cardiovascular disease and stroke: a call to action from the American Heart Association. *Circulation* 2011; 123:1138–1143.
 53. Campion J, Maestro B, Molero S, *et al.* Aldosterone impairs insulin responsiveness in U-937 human promonocytic cells via the downregulation of its own receptor. *Cell Biochem Funct* 2002; 20:237–245.
 54. Campion J, Maestro B, Mata F, *et al.* Inhibition by aldosterone of insulin receptor mRNA levels and insulin binding in U-937 human promonocytic cells. *J Steroid Biochem Mol Biol* 1999; 70:211–218.
 55. Wada T, Ohshima S, Fujisawa E, *et al.* Aldosterone inhibits insulin-induced glucose uptake by degradation of insulin receptor substrate (IRS) 1 and IRS2 via a reactive oxygen species-mediated pathway in 3T3-L1 adipocytes. *Endocrinology* 2009; 150:1662–1669.
 56. Kraus D, Jager J, Meier B, *et al.* Aldosterone inhibits uncoupling protein-1, induces insulin resistance, and stimulates proinflammatory adipokines in adipocytes. *Horm Metab Res* 2005; 37:455–459.
 57. Hirata A, Maeda N, Hiuge A, *et al.* Blockade of mineralocorticoid receptor reverses adipocyte dysfunction and insulin resistance in obese mice. *Cardiovasc Res* 2009; 84:164–172.
 58. Hitomi H, Kiyomoto H, Nishiyama A, *et al.* Aldosterone suppresses insulin signaling via the downregulation of insulin receptor substrate-1 in vascular smooth muscle cells. *Hypertension* 2007; 50:750–755.
 59. Dandona P, Mohanty P, Ghanim H, *et al.* The suppressive effect of dietary restriction and weight loss in the obese on the generation of reactive oxygen species by leukocytes, lipid peroxidation, and protein carbonylation. *J Clin Endocrinol Metab* 2001; 86:355–362.
 60. Gilbert KC, Brown NJ. Aldosterone and inflammation. *Curr Opin Endocrinol Diabetes Obes* 2010; 17:199–204.
 61. Guo C, Ricchiuti V, Lian BQ, *et al.* Mineralocorticoid receptor blockade reverses obesity-related changes in expression of adiponectin, peroxisome proliferator-activated receptor-gamma, and proinflammatory adipokines. *Circulation* 2008; 117:2253–2261.
 62. Li P, Zhang XN, Pan CM, *et al.* Aldosterone perturbs adiponectin and PAI-1 expression and secretion in 3T3-L1 adipocytes. *Horm Metab Res* 2011; 43:464–469.
 63. Ndisang JF, Jadhav A. The heme oxygenase system attenuates pancreatic lesions and improves insulin sensitivity and glucose metabolism in deoxycorticosterone acetate hypertension. *Am J Physiol Regul Integr Comp Physiol* 2010; 298:R211–R223.
- DOCA and salt hypertensive rat, a model of primary aldosteronism, was treated with hemin. Hemin therapy lowered BP and increased insulin sensitivity and adiponectin with significant reduction of glycemia. The enhanced insulin sensitization in hemin-treated animals was accompanied by a reduction in urinary and muscle oxidative and inflammatory markers and an increase in antioxidants. These results suggest that impaired insulin signaling in response to aldosteronism is mediated via oxidative and inflammatory mechanisms.
64. Ndisang JF, Lane N, Jadhav A. The heme oxygenase system abates hyperglycemia in Zucker diabetic fatty rats by potentiating insulin-sensitizing pathways. *Endocrinology* 2009; 150:2098–2108.
 65. Lastra G, Whaley-Connell A, Manrique C, *et al.* Low-dose spironolactone reduces reactive oxygen species generation and improves insulin-stimulated glucose transport in skeletal muscle in the TG(mRen2)27 rat. *Am J Physiol Endocrinol Metab* 2008; 295:E110–E116.
 66. Selvaraj J, Muthusamy T, Srinivasan C, Balasubramanian K. Impact of excess aldosterone on glucose homeostasis in adult male rat. *Clin Chim Acta* 2009; 407:51–57.
 67. Atkins KB, Johns D, Watts S, *et al.* Decreased vascular glucose transporter expression and glucose uptake in DOCA-salt hypertension. *J Hypertens* 2001; 19:1581–1587.
 68. Osanai T, Tanaka M, Magota K, *et al.* Coupling factor 6-induced activation of ecto-F(1)F(o) complex induces insulin resistance, mild glucose intolerance and elevated blood pressure in mice. *Diabetologia* 2012; 55:520–529.
 69. Caprio M, Feve B, Claes A, *et al.* Pivotal role of the mineralocorticoid receptor in corticosteroid-induced adipogenesis. *FASEB J* 2007; 21:2185–2194.
 70. Luther JM, Luo P, Kreger MT, *et al.* Aldosterone decreases glucose-stimulated insulin secretion in vivo in mice and in murine islets. *Diabetologia* 2011; 54:2152–2163.

71. Wada T, Kenmochi H, Miyashita Y, *et al.* Spironolactone improves glucose and lipid metabolism by ameliorating hepatic steatosis and inflammation and suppressing enhanced gluconeogenesis induced by high-fat and high-fructose diet. *Endocrinology* 2010; 151:2040–2049.

Effect of spironolactone on glucose and lipid metabolism was examined in a mouse model of diet-induced diabetes and nonalcoholic fatty liver disease. Treatment with spironolactone did not affect the calorie intake or body weight but it decreased BP, serum triglyceride, free fatty acids, leptin, and total cholesterol levels. Elevation of blood glucose during glucose and insulin tolerance tests was significantly lowered by spironolactone. Hepatic mRNA expression of proinflammatory cytokines (TNF α , IL-6, and MCP-1) and gluconeogenic genes was significantly suppressed by spironolactone. These results indicate that inhibition of mineralocorticoid receptor might be a beneficial therapeutic approach for diet-induced phenotypes of metabolic syndrome and fatty liver.

72. Whaley-Connell A, Habibi J, Wei Y, *et al.* Mineralocorticoid receptor antagonism attenuates glomerular filtration barrier remodeling in the transgenic Ren2 rat. *Am J Physiol Renal Physiol* 2009; 296:F1013–F1022.

73. Guo C, Martinez-Vasquez D, Mendez GP, *et al.* Mineralocorticoid receptor antagonist reduces renal injury in rodent models of types 1 and 2 diabetes mellitus. *Endocrinology* 2006; 147:5363–5373.
74. Schafer A, Vogt C, Fraccarollo D, *et al.* Eplerenone improves vascular function and reduces platelet activation in diabetic rats. *J Physiol Pharmacol* 2010; 61:45–52.
75. Mejia-Montilla J, Reyna-Villasamil E, Torres-Cepeda D, *et al.* Plasma renin and aldosterone levels in obese and nonobese women with polycystic ovary syndrome. *Endocrinol Nutr* 2012; 59:21–27.
76. Diamanti-Kandarakis E, Economou FN, Livadas S, *et al.* Hyperreninemia characterizing women with polycystic ovary syndrome improves after metformin therapy. *Kidney Blood Press Res* 2009; 32:24–31.
77. Menon DV, Arbique D, Wang Z, *et al.* Differential effects of chlorthalidone versus spironolactone on muscle sympathetic nerve activity in hypertensive patients. *J Clin Endocrinol Metab* 2009; 94:1361–1366.
78. McMurray JJ, Holman RR, Haffner SM, *et al.* Effect of valsartan on the incidence of diabetes and cardiovascular events. *N Engl J Med* 2010; 362:1477–1490.