

Journal of the American College of Nutrition

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uacn20>

Effects of Dairy Products on Intracellular Calcium and Blood Pressure in Adults with Essential Hypertension

Kirsten F. Hilpert PhD^a, Sheila G. West PhD^b, Deborah M. Bagshaw BS^a, Valerie Fishell BS^a, Linda Barnhart DO^b, Michael Lefevre PhD^d, Marlene M. Most PhD^d, Michael B. Zemel PhD^e, Mosuk Chow PhD^c, Alan L. Hinderliter MD^f & Penny M. Kris-Etherton PhD^a

^a Departments of Nutritional Sciences (K.F.H., D.M.B., V.F., P.M.K.-E.)

^b Biobehavioral Health (S.G.W., L.B.)

^c Statistics (M.C.), The Pennsylvania State University, University Park, Pennsylvania

^d Division of Functional Foods Research, Pennington Biomedical Research Center, Baton Rouge, Louisiana (M.L., M.M.M.)

^e Department of Nutrition, University of Tennessee, Knoxville, Tennessee (M.B.Z.)

^f Department of Medicine, Division of Cardiology, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina (A.L.H.)

Published online: 14 Jun 2013.

To cite this article: Kirsten F. Hilpert PhD, Sheila G. West PhD, Deborah M. Bagshaw BS, Valerie Fishell BS, Linda Barnhart DO, Michael Lefevre PhD, Marlene M. Most PhD, Michael B. Zemel PhD, Mosuk Chow PhD, Alan L. Hinderliter MD & Penny M. Kris-Etherton PhD (2009) Effects of Dairy Products on Intracellular Calcium and Blood Pressure in Adults with Essential Hypertension, *Journal of the American College of Nutrition*, 28:2, 142-149, DOI: [10.1080/07315724.2009.10719765](https://doi.org/10.1080/07315724.2009.10719765)

To link to this article: <http://dx.doi.org/10.1080/07315724.2009.10719765>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Original Research

Effects of Dairy Products on Intracellular Calcium and Blood Pressure in Adults with Essential Hypertension

Kirsten F. Hilpert, PhD, Sheila G. West, PhD, Deborah M. Bagshaw, BS, Valerie Fishell, BS, Linda Barnhart, DO, Michael Lefevre, PhD, Marlene M. Most, PhD, Michael B. Zemel, PhD, Mosuk Chow, PhD, Alan L. Hinderliter, MD, and Penny M. Kris-Etherton, PhD

Departments of Nutritional Sciences (K.F.H., D.M.B., V.F., P.M.K.-E.), Biobehavioral Health (S.G.W., L.B.), and Statistics (M.C.), The Pennsylvania State University, University Park, Pennsylvania; the Division of Functional Foods Research, Pennington Biomedical Research Center, Baton Rouge, Louisiana (M.L., M.M.M.); the Department of Nutrition, University of Tennessee, Knoxville, Tennessee (M.B.Z.); and the Department of Medicine, Division of Cardiology, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina (A.L.H.)

Key words: dairy, milk, calcium, intracellular calcium, intracellular magnesium, vitamin D, hypertension

Background: Consumption of dairy foods has been associated with lower blood pressure in certain populations.

Objective: This study examined the effects of dairy foods on blood pressure (BP) and intracellular calcium ($(Ca)_i$) and the dependence of BP changes on changes in $(Ca)_i$.

Design: Twenty-three stage 1 hypertensive adults were fed the following 3 experimental diets (5 wk each) in a randomized cross-over design study; a dairy-rich, high fruits and vegetables diet (D-F&V; 30% fat, 7% saturated fat (SFA), 3.4 servings/d dairy), a high fruits and vegetables diet (F&V; 30% fat, 7% SFA, 0.4 servings/d dairy), and an average Western diet (control; 36% fat, 15% SFA, 0.4 servings/d dairy). Systolic (SBP) and diastolic (DBP) BP, calcium regulatory hormones, and erythrocyte $(Ca)_i$ were determined.

Results: SBP and DBP were significantly reduced by ~ 2 mm Hg following both D-F&V and F&V diets vs. the control ($P < 0.05$). The D-F&V diet significantly lowered 1,25-dihydroxyvitaminD compared with the F&V and control diets ($P < 0.01$). Serum calcium, parathyroid hormone, calcitonin, and renin activity were unchanged. The D-F&V diet lowered $(Ca)_i$ vs. the other two diets ($P < 0.01$), and this change correlated with the fall in DBP ($r = 0.52$, $P < 0.05$). Subjects who responded to the D-F&V diet by significantly reducing $(Ca)_i$ exhibited significantly greater net decreases in DBP on the D-F&V vs. the F&V (-2.8 ± 1.0 mm Hg) and control diets (-5.4 ± 1.0 mm Hg; diet \times group interaction, $P < 0.02$).

Conclusion: Consumption of dairy foods beneficially affects $(Ca)_i$, resulting in improved BP in a subgroup defined by $(Ca)_i$ response.

INTRODUCTION

The Dietary Approaches to Stop Hypertension (DASH) study [1] showed that a diet high in fruits and vegetables significantly reduced systolic (SBP) and diastolic (DBP) blood pressure by 2.8 and 1.1 mm Hg, respectively, compared with a typical Western diet. The DASH diet, which emphasized fruits,

vegetables and low-fat dairy products and lower saturated fat, elicited even greater reductions in blood pressure (BP) of 5.5 mm Hg SBP and 3.0 mm Hg DBP (5.5/3.0 mm Hg). The BP reductions on the DASH diet were similar to those achieved by medication, and the diet controlled hypertension in 70% of subjects with hypertension [2]. Although it suggested a unique hypotensive effect of dairy foods, the DASH study was not

Address correspondence to: SG West, 315 Health and Human Development Building East, Department of Biobehavioral Health, University Park, PA 16802. E-mail: sgw2@psu.edu

Supported by a grant from the National Dairy Council, Rosemont, IL (to PMK-E, SGW, ML, MMM, ALH); supported in part by the General Clinical Research Center, The Pennsylvania State University (NIH grant M01RR10732).

Disclaimer: This study was funded by the dairy industry through a research grant to the investigators (to P.M.K.-E., S.G.W., M.L., M.M.M., A.L.H.). The funding group had no involvement in data analysis and preparation of the manuscript.

Journal of the American College of Nutrition, Vol. 28, No. 2, 142–149 (2009)

Published by the American College of Nutrition

specifically designed to identify principal nutrients/foods that accounted for its BP effects. Furthermore, the biological mechanism(s) for this effect are not well understood.

Epidemiologic studies have shown that consumption of dairy products is associated with reductions in BP [3–6] and lower risks for stroke [7,8], type 2 diabetes [9–11], and metabolic syndrome [12,13]. The present study was designed to assess the contribution of dairy foods to the hypotensive effect of a DASH-type diet and to examine changes in intracellular calcium $[(Ca)_i]$ as a potential mediator of this effect. $(Ca)_i$, a potent mediator of cellular response, is key in the regulation of smooth muscle function, peripheral vascular tone, BP and volume homeostasis [14]. Resnick and others [15,16] have proposed that hypertension and associated disorders result from elevated $(Ca)_i$ concentrations and reciprocally suppressed intracellular magnesium levels. Hypertension is associated with significantly higher intracellular calcium and sodium concentrations [17–20] and lower intracellular magnesium levels [21–23]. A direct correlation between BP and $(Ca)_i$ levels has been shown in a variety of cell types [17,19,24]. Several clinical studies indicate that dietary calcium normalizes intracellular ion levels, via the suppression of calcium-regulating hormones [25–28]. However, not all studies show a benefit of dietary calcium on BP [28–30]. Therefore, this study was conducted to clarify the effect of dairy foods as part of a DASH-type diet on BP, intracellular ions, and calcium-regulating hormones in individuals with stage 1 hypertension.

METHODS

Subjects

Participants had untreated stage 1 hypertension (SBP = 140–159 and/or DBP = 90–99 mm Hg), based on the mean of 2 measurements collected during 3 screening visits. All participants were Caucasian and included 16 men and 7 women

22–70 years of age, with normal or moderately elevated cholesterol (LDL cholesterol < 4.91 mmol/L, HDL-cholesterol in the 15–95th percentile of NHANES III, and triacylglycerols < 3.95 mmol/L). Exclusion criteria included: body mass index > 35 kg/m²; smoking; history of congestive heart failure, peripheral artery disease, myocardial infarction, stroke, angina, intermittent claudication, kidney stones, or renal disease; allergies to food, latex, or nitroglycerin; presence of any disease known to affect calcium balance; and use of nutritional supplements, hormone replacement therapy, or any drug known to alter calcium balance, BP, or cholesterol metabolism. Table 1 shows characteristics of the subjects at study entry.

Study Design

A randomized, three-period, crossover controlled feeding study was conducted at two sites (Pennsylvania State University and Pennington Biomedical Research Center). Twenty-three subjects were studied, 17 at Penn State and 6 at Pennington. Twenty completed all diet periods and 3 completed two diet periods. The protocol was approved by the Biomedical Committee of the Institutional Review Board at both sites and written informed consent was obtained.

Endpoints were assessed at the end of each 5-week diet period, and a 2-week compliance break separated diets. Food was prepared in research kitchens according to study protocol. On weekdays, subjects ate one meal in the feeding center and body weight was recorded. Other meals and snacks were packaged for takeout. Diet compliance was monitored according to procedures used in our research centers. Subjects were instructed to maintain their usual lifestyle/exercise habits, to consume all foods provided, to abstain from consuming other foods, and to limit beverages containing caffeine (< 5 servings/d) and alcohol (< 2 servings/wk). Weight was maintained by adjusting calories.

Table 1. Subject Characteristics at Study Entry Stratified by Intracellular Calcium Response^{1,2}

	All subjects (n = 23)	$(Ca)_i$ Change group (n = 8)	$(Ca)_i$ Stable group (n = 10)
Age (years)	45.3 ± 2.0	49.7 ± 3.4	44.2 ± 3.0
Weight (kg) ³	89.3 ± 4.1	82.1 ± 6.0	95.0 ± 5.3
Percent Male	70	75	80
Body mass index (kg/m ²)	28.8 ± 0.9	26.7 ± 1.5	30.0 ± 1.4
Systolic blood pressure (mm Hg)	140.5 ± 1.9	143.5 ± 3.0	139.4 ± 2.7
Diastolic blood pressure (mm Hg)	91.1 ± 1.3	92.4 ± 2.2	89.3 ± 1.9
Total cholesterol (mmol/L)	5.2 ± 0.2	5.1 ± 0.3	5.3 ± 0.2
LDL cholesterol (mmol/L)	3.3 ± 0.2	3.3 ± 0.2	3.4 ± 0.2
HDL cholesterol (mmol/L)	1.2 ± 0.1	1.2 ± 0.1	1.3 ± 0.1
Triacylglycerols (mmol/L)	1.4 ± 0.2	1.3 ± 0.3	1.4 ± 0.2

¹ All values are LS means ± SE. $(Ca)_i$ Change Group, decrease in RBC calcium $\geq 3.4 \mu\text{mol/L}$ and $(Ca)_i$ Stable Group, decrease in RBC calcium < $3.4 \mu\text{mol/L}$ (Change in RBC calcium calculated by D-F&V diet - mean F&V and AWD diets). The Change and Stable groups, n = 18; five samples were lost during the study.

² No significant differences between groups at study entry (PROC MIXED model and Tukey's least-squares-means test).

³ No significant differences between men and women, except women weighed significantly less than men, $P < 0.003$. (PROC MIXED model and Tukey's least-squares-means test).

Experimental Diets

Three experimental diets were designed to evaluate the unique role of dairy foods in BP regulation (Table 2). Diets included an average Western diet (AWD) as a control, a diet high in fruits and vegetables (F&V), and a similar diet containing 3.4 servings/d of dairy products (D-F&V). The F&V and D-F&V diets were matched for macronutrients, cholesterol, and fiber, and included the same fruits and vegetables in similar proportions. The F&V and AWD diets contained 0.4 servings/d of dairy products. The D-F&V diet included low-fat and non-fat milk and yogurt and full-fat cheese. The AWD diet included full-fat dairy products, while the F&V diet included lower fat dairy products. Sodium was matched across all 3 diets, and calcium was matched on the F&V and AWD diets and did not exceed 600 mg/d at the highest kilocalorie level.

Diets were formulated using menus from the DASH Study [1] and were modified to meet the experimental criteria using Nutritionist V (N-Squared Computing, San Bruno, CA.). A 6-day menu cycle with 8 kilocalorie levels (1800–3900 kcal) was developed for each diet. Nutrient composition of diets prepared at both study sites was determined by chemical analyses. Protein content was analyzed with a Perkin Elmer nitrogen analyzer. Total fat content was extracted with chloroform and methanol and fatty acid content was measured by gas chromatography [31]. Minerals (Ca, Na, Mg, K, and P) were determined by inductively coupled plasma-mass spectrometry.

Blood Pressure

During the last week of each diet period, participants made four visits for BP measurements. SBP and DBP were measured in a controlled environment using a calibrated mercury manometer, stethoscope, and appropriately sized cuffs, following the

JNC-VII guidelines (17). Participants were instructed not to exercise or ingest food/caffeine 30 minutes before the visit. After a 5 minute rest period, three stethoscopic measurements were taken, separated by 1 minute. Participants were seated with feet flat on the floor and arm held at heart level. SBP was recorded at appearance of Korotkoff sounds (phase I) and DBP was recorded at disappearance (phase V). Means of all endpoint measures were used for analyses. Technicians were certified in the use of a standard protocol and were blind to treatment assignment.

Biochemical Assays

At study entry, serum total and HDL cholesterol and triacylglycerols were measured by enzymatic assays with commercially available kits. HDL cholesterol was determined after precipitation of apolipoprotein B-containing lipoproteins with dextran sulfate and magnesium. LDL cholesterol was calculated using the Friedewald equation.

Fasting blood samples and 24-hour urine samples were collected and stored at -80°C until analysis. Serum intact parathyroid hormone (iPTH) was measured by immunochemiluminometric assay (Nichols Institute Diagnostics, San Clemente, CA). Radioimmunoassays were used to determine serum 1,25-dihydroxyvitaminD (Immundiagnostic System Inc., Fountain Hills, AZ), serum calcitonin (Diagnostic Systems Laboratories, Inc., Webster, TX), and plasma renin activity. Serum and urine calcium were measured by colorimetric assay. Urine sodium and potassium were measured with the ion-sensitive electrode method.

Heparinized blood was collected in chilled tubes for determination of total intracellular calcium, magnesium, sodium, and potassium in red blood cell (RBC), according to the method

Table 2. Assayed Composition of Experiment Diets (Based on 2100 kcal per Day)¹

Nutrient	AWD	F&V	D-F&V	D-F&V minus F&V	D-F&V minus AWD
Carbohydrate (%en)	46	52	52	None	6
Protein (%en)	18	18	18	None	None
Fat (%en)	36	30	30	None	−6
Saturated (%en)	15	7	7	None	−8
Monounsaturated (%en)	14	14	14	None	None
Polyunsaturated (%en)	7	9	9	None	2
Cholesterol (mg)	300	175	175	None	−125
Fiber (g)	10.5	27	27	None	16.5
Sodium (mg)	3500	3500	3500	None	None
Potassium (mg)	1750	3900	4600	700	2850
Calcium (mg)	400	400	1200	800	800
Phosphorus (mg)	1150	1300	1700	400	550
Magnesium (mg)	190	370	420	50	230
Fruit (serving/day)	1.6	5.2	5.2	None	3.6
Vegetable (serving/day)	2.0	4.4	4.4	None	2.4
Dairy (serving/day)	0.4	0.4	3.4	3	3

¹ Nutrient values were calculated using Nutritionist V software (N-Squared Computing, San Bruno, CA) and subsequently measured by chemical analyses at both study sites. AWD, average Western diet; F&V, fruits and vegetables diet; D-F&V, dairy-rich fruits and vegetables diet; %en, percent of energy. To convert values to millimoles, multiply by 0.043498 for sodium, 0.025577 for potassium, 0.024951 for calcium, 0.032285 for phosphorus, and 0.041144 for magnesium.

described by Zemel et al. [26] using inductively coupled plasma mass spectrometry (Finnigan Element High Resolution). Erythrocytes were chosen as a cell model to represent global changes in intracellular calcium status and due to the ease with which they can be isolated.

Statistical Analysis

The specific aims of the study were to test the effects of the D-F&V diet compared with the control diet; the effects of the F&V diet compared with the control diet; and the effects of the D-F&V diet compared with the F&V diet. The key outcome variables were change in SBP and change in DBP, defined as the difference between screening and end-of-diet levels. Statistical analyses were performed with SAS (version 9.1; SAS Institute Inc, Cary, NC). Results are expressed as least squares means \pm SE. Data for serum triacylglycerols were logarithmically transformed (base 10) before conducting analyses because of skewed distributions. The mixed procedure (PROC MIXED) was used to test for effects of diet, order (the sequence of the 3 diets given to each subject), diet period and their interactive effects on each outcome variable. All models were adjusted for study site and age; models for BP change were also adjusted for screening values. When significant diet effects were detected, a Tukey's least-significant difference test was performed to determine the source of significant main effects or interactions ($P \leq 0.05$).

We also tested whether treatment effects differed according to magnitude of $(Ca)_i$ response in a subset of participants ($n = 18$). To determine diet-related changes in $(Ca)_i$, levels of RBC calcium on the F&V and control diets, which were not significantly different (12.1 ± 1.1 and $12.6 \pm 1.0 \mu\text{mol/L}$, $P = 0.9$), were pooled and subtracted from the D-F&V diet. The mean decrease in $(Ca)_i$ was $-4.9 \mu\text{mol/L}$, with a natural break between -3.3 and $-6.1 \mu\text{mol/L}$ (Fig. 1). Subjects who responded to the D-F&V diet by decreasing $(Ca)_i \geq 3.4 \mu\text{mol/L}$ were defined as the $(Ca)_i$ Change Group ($n = 8$). Others were labeled the $(Ca)_i$ Stable Group ($n = 10$) and calcium status was added to the models. Within-subject correlation analyses were used to test associations between RBC calcium and BP.

After a small reduction in weight after the first period ($-0.85 \pm 0.35 \text{ kg}$, $P < 0.03$), weight remained stable for the duration of the study. Changes in weight were equal in magnitude among diets ($P = 1.0$) and groups ($P = 0.4$). Adjustment for weight had no effect on the results, and unadjusted results are presented.

RESULTS

Diet Effects on BP Change, Calcium Hormones, RBC Ions, and 24-Hour Urine Electrolytes

All three experimental diets significantly decreased SBP and DBP from screening (SBP: -9.9 ± 1.5 , -12.3 ± 1.5 ,




Fig. 1. Distribution of changes in RBC calcium on the dairy-rich diet. The $(Ca)_i$ Change Group decreased RBC calcium $\geq 3.4 \mu\text{mol/L}$ (black bars), whereas the $(Ca)_i$ Stable Group decreased it $< 3.3 \mu\text{mol/L}$ (white bars). Change in RBC calcium was calculated by subtracting the D-F&V diet from the mean of F&V and AWD diets (no significant difference between F&V and AWD diets). $(Ca)_i$ Change group had a significantly greater reduction in RBC calcium following the D-F&V diet than the $(Ca)_i$ Stable group ($-7.5 \pm 1.1 \mu\text{mol/L}$ vs. $-2.6 \pm 0.7 \mu\text{mol/L}$, $P < 0.01$). PROC MIXED model and Tukey's least-squares-means test.

$-12.0 \pm 1.5 \text{ mm Hg}$; DBP: -5.3 ± 1.2 , -7.2 ± 1.2 , $-7.0 \pm 1.2 \text{ mm Hg}$ for AWD, F&V, and D-F&V diets, respectively, $P < 0.01$ for all). The reductions in SBP and DBP following the D-F&V and F&V diets were significantly greater compared with AWD (Table 3). The changes in SBP or DBP did not differ between the two intervention diets. For SBP, we observed a significant period effect. SBP was significantly lower during period 3 vs. period 1 ($P = 0.02$), suggesting that SBP continued to decrease throughout the study.

While serum calcium remained constant, we observed the expected lower concentration of 1,25-dihydroxyvitaminD on the D-F&V diet vs. the F&V and control diets (Table 4). Measures of iPTH, calcitonin, and renin activity did not differ significantly between diets.

The concentration of RBC calcium was significantly lower and the concentration of RBC magnesium was significantly higher after the D-F&V diet vs. both the F&V and control diets (Table 4). Consequently, the ratio of RBC calcium:magnesium was significantly lower on the D-F&V diet. RBC potassium and RBC sodium did not differ significantly between diets.

There were significant main effects of diet for urinary excretion of calcium and potassium (Table 4). Urine calcium was greatest on the D-F&V diet, and was significantly different from the F&V diet. Urine potassium was significantly elevated on the D-F&V diet compared with the F&V diet and the AWD diet. Urine sodium did not differ across diets.

Group Effects on BP Response by Diet ($n = 18$)

By design, the $(Ca)_i$ Change group had a significantly greater reduction in RBC calcium following the D-F&V diet than the $(Ca)_i$ Stable group ($-7.5 \pm 1.1 \mu\text{mol/L}$ vs. $-2.6 \pm$

Table 3. Mean Changes in Blood Pressure according to Diet in All Subjects and Intracellular Calcium Response¹

	Effect of diet P-value	Interaction of diet and group P-value	Change in F&V minus change in AWD		Change in D-F&V minus change in AWD		Change in D-F&V minus change in F&V	
			mmHg	Tukey P	mm Hg	Tukey P	mm Hg	Tukey P
SBP								
All subjects ²	0.015	-	-2.4 ± 0.9	<0.02	-2.0 ± 0.9	<0.05	0.3 ± 0.9	<0.9
(Ca) _i Group	-	0.05						
Stable			-2.7 ± 1.4	<0.2	-1.5 ± 1.4	<0.5	1.2 ± 1.4	<0.6
Change			-2.1 ± 1.5	<0.3	-4.2 ± 1.5	<0.02	-2.1 ± 1.5	<0.3
DBP								
All subjects	0.005	-	-1.8 ± 0.6	<0.01	-1.7 ± 0.6	<0.02	0.1 ± 0.6	<1.0
(Ca) _i Group	-	<0.0001						
Stable			-1.9 ± 1.0	<0.2	-0.4 ± 1.0	<0.9	1.5 ± 1.0	<0.3
Change			-2.6 ± 1.0	<0.02	-5.4 ± 1.0	<0.01	-2.8 ± 1.0	<0.02

¹ Data are least squares means ± SE, n = 23. Mean values have been adjusted for study site, age, and screening levels (refer to Table 1). The Ca Stable group, n = 10 and the Ca Change group, n = 8. AWD, average Western diet; F&V, fruits and vegetables diet; D-F&V, dairy-rich fruits and vegetables diet; SBP, systolic blood pressure; DBP, diastolic blood pressure

² Period effect, P = 0.03. PROC MIXED model and Tukey's least-squares-means test.

Table 4. Effect of Three Experimental Diets on Calcium Related Variables, Red Blood Cell Ions, and Urinary Electrolytes¹

Variable	AWD	F&V	D-F&V	P-value
Serum Calcium (mmol/L)	2.33 ± 0.03	2.35 ± 0.03	2.35 ± 0.03	NS
1,25(OH) ₂ D(pmol/L)	112.2 ± 7.0 ^a	104.0 ± 7.0 ^a	87.4 ± 7.0 ^b	0.0004
iPTH (pg/mL)	44.7 ± 4.5	44.8 ± 4.4	42.8 ± 4.5	NS
Calcitonin (pg/mL)	9.6 ± 0.7	9.6 ± 0.7	9.4 ± 0.7	NS
Renin activity (ng/mL/hr)	0.6 ± 0.2	0.8 ± 0.2	0.9 ± 0.2	NS
Red Blood Cell ²				
Calcium (μmol/L)	12.6 ± 1.0 ^a	12.1 ± 1.1 ^a	6.9 ± 1.1 ^b	0.0001
Magnesium (mmol/L)	2.3 ± 0.1 ^a	2.3 ± 0.1 ^a	2.7 ± 0.1 ^b	0.0010
Sodium (mmol/L)	6.6 ± 0.6	8.0 ± 0.6	6.2 ± 0.6	0.0518
Potassium (mmol/L)	86.1 ± 2.6	89.0 ± 2.6	88.0 ± 2.8	NS
Calcium:magnesium	5.6 ± 0.5 ^a	5.5 ± 0.5 ^a	2.9 ± 0.5 ^b	0.0001
24-hour Urine ³				
Calcium (mg)	107.2 ± 17.3 ^{a,b}	87.8 ± 16.0 ^b	126.5 ± 16.1 ^a	0.0214
Sodium (mg)	3605 ± 322	3570 ± 280	3871 ± 281	NS
Potassium (mg)	1983 ± 265 ^a	2510 ± 237 ^a	3334 ± 238 ^b	0.0001

¹ All values are least squares means ± SE; n = 23. AWD, average Western diet; F&V, fruits and vegetables diet; D-F&V, dairy-rich fruits and vegetables diet; 1,25(OH)₂VD, 1,25-dihydroxyvitaminD; iPTH, intact parathyroid hormone; NS, nonsignificant. Values in a row with different superscript letters are significantly different, Tukey P ≤ 0.02 (PROC MIXED model and Tukey's least-squares-means test).

² n = 18.

³ To convert values to millimoles per day, multiply by 0.024951 for calcium, 0.043498 for sodium, and 0.025577 for potassium.

0.7 μmol/L, P < 0.01). The groups did not differ significantly in study entry characteristics (Table 1).

There was a significant diet by group interaction for change in SBP (Table 3). In the (Ca)_i Change group, the D-F&V diet elicited a significantly greater reduction in SBP compared with the AWD, while the F&V diet did not differ from the other diets.

There was a significant diet by group interaction for DBP (Table 3). In the (Ca)_i Change group, the D-F&V diet elicited significantly greater decreases in DBP compared with the F&V and AWD diets. The magnitude of the change in DBP on the F&V diet also was significantly greater than the control. This pattern in BP response was not evident in the (Ca)_i Stable group.

Within-subject correlations between RBC calcium and BP support the subgroup analyses. The reduction in RBC calcium associated with the D-F&V diet correlated with the decrease in DBP (r = 0.52, P < 0.05) and 1,25-dihydroxyvitaminD (r = 0.59, P < 0.03). Participants who exhibited the greatest decreases in RBC calcium also experienced the greatest reductions in 1,25-dihydroxyvitaminD and DBP. Changes in SBP were not significantly correlated with changes in RBC calcium.

DISCUSSION

In the present study, we observed a modest additional reduction in SBP and DBP (about 2 mm Hg) after the F&V and

D-F&V diets versus the control diet. The addition of 3 servings of dairy to a diet high in fruits and vegetables, however, had no effect on mean blood pressure changes, except in the subgroup characterized by appreciable decreases in intracellular calcium. The D-F&V diet significantly reduced $(Ca)_i$ levels and significantly increased intracellular magnesium levels, and reductions in $(Ca)_i$ were correlated to improved BP. Thus, a hypotensive effect of dairy products was evident only in subjects most responsive to changes in intracellular ion levels.

Our findings agree with previous studies showing that a calcium-rich diet favorably affects (i.e., decreases) intracellular ion levels. A clinical feeding trial [26] in 11 salt-sensitive hypertensive African Americans showed that a low calcium (8.9 mmol/d), high sodium (174 mmol/d) diet significantly increased PTH, which was associated with significant increases in RBC calcium (5.4 ± 0.7 to $11.1 \pm 3.7 \mu\text{mol/L}$) and RBC sodium (9.6 ± 0.3 to $11.4 \pm 0.4 \text{ mmol/L}$) and significant decreases in RBC magnesium (2.2 ± 0.1 to $1.8 \pm 0.1 \text{ mmol/L}$). The addition of calcium (to equal 23.4 mmol/d) to the high sodium diet reversed these effects [26] and was associated with lower SBP and DBP [25,32]. Calcium supplementation (50 mmol/d for 16 wk) in normotensive men also has been shown to reduce intracellular calcium and sodium, 1,25-dihydroxyvitaminD, and SBP [27,33]. In the present study, we observed a similar reduction in RBC calcium, RBC sodium and 1,25-dihydroxyvitaminD, and an increase in RBC magnesium during the dairy-rich diet. Collectively, these results suggest that intracellular ion levels are responsive to diet manipulation and that consumption of dairy foods produces a favorable intracellular environment (lower ratio of intracellular calcium to magnesium). It is important to note that while we focused on calcium and its role in decreasing blood pressure in the current study, it is possible that changes in other micronutrients contributed to the response. The D-F&V diet was higher in calcium, potassium, magnesium, and phosphorous, and RBC levels of calcium were lower and magnesium was higher. Thus, it may be that the blood pressure response noted was the result of changes in micronutrients beyond calcium or due to other components of dairy products.

In agreement with several studies [26,28–30], we observed significantly greater decreases in 1,25-dihydroxyvitaminD on the dairy-supplemented diet vs. the F&V and control diets, which contained lower levels of dietary calcium. The present study found a direct relationship between the changes in 1,25-dihydroxyvitaminD and RBC calcium. 1,25-dihydroxyvitaminD has been shown to stimulate rapid calcium influx through increasing calcium-channel currents in a variety of cells including vascular smooth muscle cells [34–36]. Thus, we propose that the calcium-rich D-F&V diet favorably affected (decreased) $(Ca)_i$ levels via suppression of 1,25-dihydroxyvitaminD.

Our reduction in blood pressure was notably less than that reported in all subjects (4.2/5.4) by the original DASH study [1]. However, in Caucasian subjects the DASH diet produced a

less robust blood pressure reduction of 3.3/2.4 mm Hg, which is comparable to the results of our small sample of 23 Caucasians [37]. There are other possible reasons why our BP changes were smaller in magnitude than in the original DASH study. It is likely that the beneficial effects of our intervention diets were masked by parallel reductions in blood pressure during the control diet. Compared to screening levels, both the D-F&V and the F&V diets produced similar drops in blood pressure (-12 mm Hg SBP and -7 mm Hg DBP), however, unlike the original DASH study, our control diet also significantly reduced blood pressure (-10 mm Hg SBP and -5 mm Hg DBP). This effect of our control diet was unexpected; nonetheless, this response partially accounts for the smaller net decreases in BP observed in our intervention diets. We speculate that our control diet was healthier than the participants habitual diet, a finding we note repeatedly in our studies. In addition, inclusion of a run-in period in our study (as was done in the DASH study) may have prevented the BP decrease on the control diet.

Another reason the present study did not elicit the same BP-lowering effect as the DASH trial may relate to the lack of diversity of our small sample. Like the hypertensive subgroup in the DASH study [2], our subjects were overweight, middle-aged adults with stage 1 hypertension. In the DASH trial, 60% of the subjects were African American, the group showing the most robust dietary response [37]. Therefore, our study of 23 Caucasian hypertensive adults may have underestimated the magnitude of BP change that is possible with dairy foods, as African Americans are particularly vulnerable to the complications of untreated hypertension and are more sensitive to changes in calcium and potassium intake [38].

In the DASH trial, there was a greater contrast in the nutrient composition of the experimental diets than in the present study. Our control diet and the two intervention diets were matched for sodium, protein, and cholesterol content, whereas sodium was the only nutrient held constant across the three diets of the DASH study. In fact, the DASH study had greater differences in protein, cholesterol, fiber, and magnesium content between the control and DASH diet than our control vs. the D-F&V diet. Furthermore, the present study was specifically designed to isolate the contribution of 3 servings of dairy foods by comparing the D-F&V diet to a similar diet that was low in dairy. Therefore, the two intervention diets (D-F&V and the F&V) were carefully matched to eliminate the macronutrient differences present in the DASH trial, while allowing calcium, magnesium, potassium, and phosphorus to increase with the increased consumption of dairy foods (refer to Table 2). In the DASH trial, the DASH diet provided less total fat (27% vs 37% of total kcal), less saturated fat (6% vs. 16% of total kcal), less cholesterol (150 vs. 300 mg/day), and more protein (18% vs. 15% of total kcal) than its fruit and vegetable counterpart. Plus, the amount and type of fruits and vegetables were identical across our two intervention diets, and this differs from the original DASH diet design. Thus, it is possible that the

magnitude of the blood pressure reduction in the present study was related to the magnitude of the difference in specific nutrient intakes between the control and interventions diets, and we did not demonstrate such large differences as in the DASH study.

Although epidemiologic studies of the relationship between dietary calcium and blood pressure reveal that those with a low intake of calcium (<300–600 mg/d) exhibit higher blood pressures [5,39], intervention studies using calcium supplementation have shown inconsistent results. It is important to note that this lack of effect of dietary calcium may be influenced by the heterogeneity of the hypertensive population. Clinical studies showing a hypotensive effect of dietary calcium in salt-sensitive [26] and/or low renin hypertensive [40] subjects lend support to the theory that calcium and/or dairy foods may play a key role in blood pressure regulation in specific patient groups (i.e. those thought to be in a state of calcium deficiency). In our study, we investigated intracellular calcium changes and observed that those who had the greatest drop in intracellular calcium also had the greatest BP-lowering response to the D-F&V diet. The (Ca)_i Change group exhibited a significantly greater reduction in BP on the D-F&V vs. control diet (–4.2/–5.4 mm Hg) and F&V diet (–2.1/–2.8 mm Hg; nonsignificant for SBP). Although addition of 3 servings of dairy foods to a diet high in fruits and vegetables did not elicit further reductions in BP in the sample as a whole, subgroup analysis revealed a unique role for dairy foods in the treatment of hypertension. Future studies should examine whether the (Ca)_i response is dose dependent and determine the basis for this phenotypic response in order to facilitate *a priori* discrimination between responders and non-responders to this dietary intervention.

In conclusion, we observed a modest reduction in SBP and DBP of about 2 mm Hg during both the F&V and D-F&V diets compared to the control diet. This magnitude of reduction has been estimated to reduce the prevalence of hypertension, the risk of coronary heart disease, and the risk of stroke in the population by 17%, 6%, and 15%, respectively [41]. The addition of 3 servings of dairy products to a diet already high in fruits and vegetables conferred significant reductions in (Ca)_i and significant increases in intracellular magnesium levels. Furthermore, the hypotensive effect of dairy products was evident only in subjects most responsive to changes in intracellular calcium, which lends support to other studies that have identified specific populations that are particularly responsive to calcium-rich diets. Our findings suggest that the BP responses to increasing dietary calcium are associated with attenuation of the widely reported elevation in intracellular calcium in hypertensive subjects.

ACKNOWLEDGMENTS

We appreciate technical support from Katrina La Rue Hurston and John Kittleson. K.F.H., S.G.W., M.L., M.M.M.,

A.L.H., D.M.B., V.F., L.B., and P.M.K.-E. were responsible for the study design, menu development, endpoint assays, data analyses and interpretation, and writing of the manuscript. M.C. consulted on study design and data analyses. M.B.Z. provided valuable expertise in the determination of intracellular cations, data analyses and interpretation, and writing of the manuscript. None of the authors had a conflict of interest in the study design and conduct, as well as interpretation and reporting of study results.

REFERENCES

1. Appel LJ, Moore TJ, Obarzanek E, Vollmer WM, Svetkey LP, Sacks FM, Bray GA, Vogt TM, Cutler JA, Windhauser MM, Lin PH, Karanja N: A clinical trial of the effects of dietary patterns on blood pressure. DASH Collaborative Research Group. *N Engl J Med* 336:1117–1124, 1997.
2. Conlin PR, Chow D, Miller ER, 3rd, Svetkey LP, Lin PH, Harsha DW, Moore TJ, Sacks FM, Appel LJ: The effect of dietary patterns on blood pressure control in hypertensive patients: results from the Dietary Approaches to Stop Hypertension (DASH) trial. *Am J Hypertens* 13:949–955, 2000.
3. Garcia-Palmeri MR, Costas R, Jr., Cruz-Vidal M, Sorlie PD, Tillotson J, Havlik RJ: Milk consumption, calcium intake, and decreased hypertension in Puerto Rico. Puerto Rico Heart Health Program study. *Hypertension* 6:322–328, 1984.
4. Trevisan M, Krogh V, Farinero E, Panico S, Mancini M: Calcium-rich foods and blood pressure: findings from the Italian National Research Council Study (the Nine Communities Study). *Am J Epidemiol* 127:1155–1163, 1988.
5. McCarron DA, Morris CD, Henry HJ, Stanton JL: Blood pressure and nutrient intake in the United States. *Science* 224:1392–1398, 1984.
6. Ruidavets JB, Bongard V, Simon C, Dallongeville J, Ducimetiere P, Arveiler D, Amouyel P, Bingham A, Ferrieres J: Independent contribution of dairy products and calcium intake to blood pressure variations at a population level. *J Hypertens* 24:671–681, 2006.
7. Abbott RD, Curb JD, Rodriguez BL, Sharp DS, Burchfiel CM, Yano K: Effect of dietary calcium and milk consumption on risk of thromboembolic stroke in older middle-aged men. The Honolulu Heart Program. *Stroke* 27:813–818, 1996.
8. Elwood PC, Strain JJ, Robson PJ, Fehily AM, Hughes J, Pickering J, Ness A: Milk consumption, stroke, and heart attack risk: evidence from the Caerphilly cohort of older men. *J Epidemiol Community Health* 59:502–505, 2005.
9. Choi HK, Willett WC, Stampfer MJ, Rimm E, Hu FB: Dairy consumption and risk of type 2 diabetes mellitus in men: a prospective study. *Arch Intern Med* 165:997–1003, 2005.
10. Pittas AG, Dawson-Hughes B, Li T, Van Dam RM, Willett WC, Manson JE, Hu FB: Vitamin D and calcium intake in relation to type 2 diabetes in women. *Diabetes Care* 29:650–656, 2006.
11. Liu S, Choi HK, Ford E, Song Y, Klevak A, Buring JE, Manson JE: A prospective study of dairy intake and the risk of type 2 diabetes in women. *Diabetes Care* 29:1579–1584, 2006.
12. Liu S, Song Y, Ford ES, Manson JE, Buring JE, Ridker PM:

- Dietary calcium, vitamin D, and the prevalence of metabolic syndrome in middle-aged and older U.S. women. *Diabetes Care* 28: 2926–2932, 2005.
13. Azadbakht L, Mirmiran P, Esmailzadeh A, Azizi F: Dairy consumption is inversely associated with the prevalence of the metabolic syndrome in Tehranian adults. *Am J Clin Nutr* 82:523–530, 2005.
14. Walsh MP: Regulation of vascular smooth muscle tone. *Can J Physiol Pharmacol* 72:919–936, 1994.
15. Barbagallo M, Resnick LM, Dominguez LJ, Licata G: Diabetes mellitus, hypertension and ageing: the ionic hypothesis of ageing and cardiovascular-metabolic diseases. *Diabetes Metab* 23:281–294, 1997.
16. Resnick L: The cellular ionic basis of hypertension and allied clinical conditions. *Prog Cardiovasc Dis* 42:1–22, 1999.
17. Erne P, Bolli P, Burgisser E, Buhler FR: Correlation of platelet calcium with blood pressure. Effect of antihypertensive therapy. *N Engl J Med* 310:1084–108, 1984.
18. Hiraga H, Oshima T, Yoshimura M, Matsuura H, Kajiyama G: Abnormal platelet Ca^{2+} handling accompanied by increased cytosolic free Mg^{2+} in essential hypertension. *Am J Physiol* 275: R574–R579, 1998.
19. Brickman AS, Nyby MD, von Hungen K, Eggena P, Tuck ML: Calcitropic hormones, platelet calcium, and blood pressure in essential hypertension. *Hypertension* 16:515–522, 1990.
20. Hvarfner A, Larsson R, Morlin C, Rastad J, Wide L, Akerstrom G, Ljunghall S: Cytosolic free calcium in platelets: relationships to blood pressure and indices of systemic calcium metabolism. *J Hypertens* 6:71–77, 1988.
21. Resnick LM, Gupta RK, Gruenspan H, Alderman MH, Laragh JH: Hypertension and peripheral insulin resistance. Possible mediating role of intracellular free magnesium. *Am J Hypertens* 3:373–379, 1990.
22. Touyz RM, Milne FJ, Reinach SG: Platelet and erythrocyte Mg^{2+} , Ca^{2+} , Na^{+} , K^{+} and cell membrane adenosine triphosphatase activity in essential hypertension in blacks. *J Hypertens* 10:571–578, 1992.
23. Touyz RM, Milne FJ, Reinach SG: Intracellular Mg^{2+} , Ca^{2+} , Na^{2+} and K^{+} in platelets and erythrocytes of essential hypertension patients: relation to blood pressure. *Clin Exp Hypertens A* 14:1189–1209, 1992.
24. Pritchard K, Raine AE, Ashley CC, Castell LM, Somers V, Osborn C, Ledingham JG, Conway J: Correlation of blood pressure in normotensive and hypertensive individuals with platelet but not lymphocyte intracellular free calcium concentrations. *Clin Sci (Lond)* 76:631–635, 1989.
25. Zemel MB, Gualdoni SM, Walsh M: Effects of sodium and calcium on calcium metabolism and blood pressure regulation in hypertensive black adults. *J Hypertens* 4:S364–S366, 1986.
26. Zemel MB, Kraniak J, Standley PR, Sowers JR: Erythrocyte cation metabolism in salt-sensitive hypertensive blacks as affected by dietary sodium and calcium. *Am J Hypertens* 1:386–392, 1988.
27. Petrov V, Lijnen P: Modification of intracellular calcium and plasma renin by dietary calcium in men. *Am J Hypertens* 12:1217–1224, 1999.
28. Sanchez M, de la Sierra A, Coca A, Poch E, Giner V, Urbano-Marquez A: Oral calcium supplementation reduces intraplatelet free calcium concentration and insulin resistance in essential hypertensive patients. *Hypertension* 29:531–536, 1997.
29. Levey WA, Manore MM, Vaughan LA, Carroll SS, VanHalderen L, Felicetta J: Blood pressure responses of white men with hypertension to two low-sodium metabolic diets with different levels of dietary calcium. *J Am Diet Assoc* 95:1280–1287, 1995.
30. Kynast-Gales SA, Massey LK: Effects of dietary calcium from dairy products on ambulatory blood pressure in hypertensive men. *J Am Diet Assoc* 92:1497–1501, 1992.
31. American Oil Chemists Society: “Fatty Acid Composition by GLC (Marine Oils).” Champaign, IL: AOCS Press, 1993.
32. Zemel MB, Gualdoni SM, Sowers JR: Sodium excretion and plasma renin activity in normotensive and hypertensive black adults as affected by dietary calcium and sodium. *J Hypertens* 4:S343–S345, 1986.
33. Lijnen P, Petrov V: Dietary calcium, blood pressure and cell membrane cation transport systems in males. *J Hypertens* 13:875–882, 1995.
34. Tornquist K, Tashjian AH, Jr: Dual actions of 1,25-dihydroxycholecalciferol on intracellular Ca^{2+} in GH4C1 cells: evidence for effects on voltage-operated Ca^{2+} channels and $\text{Na}^{+}/\text{Ca}^{2+}$ exchange. *Endocrinology* 124:2765–2776, 1989.
35. Fleet JC: Vitamin D receptors: not just in the nucleus anymore. *Nutr Rev* 57:60–62, 1999.
36. Shan J, Resnick LM, Lewanczuk RZ, Karpinski E, Li B, Pang PK: 1,25-dihydroxyvitamin D as a cardiovascular hormone. Effects on calcium current and cytosolic free calcium in vascular smooth muscle cells. *Am J Hypertens* 6:983–988, 1993.
37. Svetkey LP, Simons-Morton D, Vollmer WM, Appel LJ, Conlin PR, Ryan DH, Ard J, Kennedy BM: Effects of dietary patterns on blood pressure: subgroup analysis of the Dietary Approaches to Stop Hypertension (DASH) randomized clinical trial. *Arch Intern Med* 159:285–293, 1999.
38. West SG, Light KC, Hinderliter AL, Stanwyck CL, Bragdon EE, Brownley KA: Potassium supplementation induces beneficial cardiovascular changes during rest and stress in salt sensitive individuals. *Health Psychol* 18:229–240, 1999.
39. Harlan WR, Harlan LC: An epidemiological perspective on dietary electrolytes and hypertension. *J Hypertens Suppl* 4:S334–339, 1986.
40. Resnick LM, Laragh JH, Sealey JE, Alderman MH: Divalent cations in essential hypertension. Relations between serum ionized calcium, magnesium, and plasma renin activity. *N Engl J Med* 309:888–891, 1983.
41. Cook NR, Cohen J, Hebert PR, Taylor JO, Hennekens CH: Implications of small reductions in diastolic blood pressure for primary prevention. *Arch Intern Med* 155:701–709, 1995.

Received October 6, 2008; revision accepted January 5, 2009.