

# Estrogen, Progesterone, and Vascular Reactivity: Potential Cellular Mechanisms\*

MARGUERITTE MABRY WHITE, STACY ZAMUDIO, TROY STEVENS,  
ROBERT TYLER, JOANN LINDENFELD, KIMBERLY LESLIE, AND  
LORNA G. MOORE

*Department of Obstetrics and Gynecology (M.M.W., K.L.), Cardiovascular Pulmonary Research Laboratory (M.M.W., T.S., R.T., J.L., L.G.M.), and Department of Anesthesia (S.Z.), University of Colorado Health Sciences Center, Denver, Colorado 80262; Department of Medicine (J.L.), Division of Cardiology, University of Colorado Health Sciences Center, Denver, Colorado 80262; Department of Anthropology (L.G.M.), University of Colorado at Denver, Denver, Colorado 80217; Women's Health Research Center (M.M.W., S.Z., J.L., K.L., L.G.M.), University of Colorado Health Sciences Center, Denver, Colorado 80262*

- I. Introduction
- II. Clinical Correlations
  - A. Protective effects of female hormones
    - 1. Cardiovascular disease
    - 2. Pregnancy
  - B. Speculations about involvement of female hormones in vascular disease
  - C. Summary
- III. Potential Mechanisms of Hormone-Mediated Effects
  - A. Receptor-mediated actions
  - B. Effects of estrogen and progesterone on vascular function
    - 1. Cardiovascular disease
    - 2. Pregnancy
  - C. Steroid-mediated effects on endothelial and vascular smooth muscle cell function
    - 1. NO
    - 2. ET-1
    - 3. Eicosanoids
    - 4. Adrenergic responsiveness
    - 5.  $\text{Ca}^{++}$  homeostasis
- IV. Summary, Conclusions, and Directions for Future Research

## I. Introduction

FEMALE hormones are broadly recognized as affecting susceptibility to vascular disease, yet comparatively little is known about the cellular mechanisms by which they exert their effects. While clinical and epidemiological data support a protective effect of estrogen on cardiovascular function, a female predominance in other vascular disorders

suggests that sex steroids influence the pathogenesis of vascular disease as well. Most of the literature on cellular effects of estrogen and progesterone has emerged from studies in the breast, uterus, and brain. Whether similar mechanisms occur in the vasculature and influence susceptibility to vascular disease is largely unknown.

The purpose of this review is to provide a framework for identifying the mechanisms by which female sex steroids influence vascular function. A diverse literature encompassing clinical observations, whole animal, isolated vessel, and cell culture studies has been incorporated in the hopes of providing a common basis for considering the influences of the female sex steroids on the vascular system. In view of the importance of the vascular endothelium and underlying smooth muscle in vascular control, emphasis is placed on the mechanisms by which estrogen and progesterone may influence these cell types and the interactions between them. Specifically, the effects of the female hormones on the production of vasoactive endothelial products, adrenergic sensitivity, and  $\text{Ca}^{++}$  homeostasis will be reviewed. Data from nonvascular tissue relevant to potential mechanisms of steroid effects in the vasculature are also presented. It is hoped that such a review will prompt further studies that will provide a resolution of the present paradoxical association between female hormones and protection from some vascular diseases (e.g. coronary artery disease) and susceptibility to others (e.g. migraines, Raynaud's phenomena, primary pulmonary hypertension).

## II. Clinical Correlations

### A. Protective effects of female hormones

Several studies point to a role for female sex steroids in favoring the balance of vasodilators to vasoconstrictors. Serum levels of nitrate and nitrite, stable metabolites of the endothelium-derived vasodilator nitric oxide (NO), increase during the follicular phase of the menstrual cycle in conjunction with rising  $17\beta$ -estradiol levels and decrease during the postovulatory, high progesterone phase (1). Lower en-

Address reprint requests to: Margueritte Mabry White, M.D., Division of Maternal Fetal Medicine, Department of Obstetrics and Gynecology (Box B-198) & Cardiovascular Pulmonary Research Laboratory, University of Colorado Health Sciences Center, 4200 East 9th Avenue, Denver, Colorado 80262

\*This review was supported by NIH HL07171, HL14985 and a Maternal Fetal Medicine fellowship. We would like to express our appreciation to Dr. Caleb Awoniyi for his thoughtful comments and advice.

dothelin (ET-1) levels, an endothelium-derived vasoconstrictor, have been reported in premenopausal women compared to age-matched men, and a reduction in ET-1 levels in male transsexuals after estrogen treatment has also been observed (2). Circulating norepinephrine levels are also reduced in the follicular compared with the luteal phase of the menstrual cycle (3). These are consistent with the possibility that estradiol enhances vasodilator and/or inhibits vasoconstrictor activity whereas progesterone antagonizes estrogen's effects.

**1. Cardiovascular disease.** The presence of female hormones is associated with a decreased incidence of cardiovascular disease. Premenopausal women lag behind men in the development of coronary artery disease by approximately 10 yr (4), but the difference between men and women narrows after 50 yr of age (5). Additionally, women with systemic hypertension have a lower morbidity and mortality when compared with hypertensive men (6, 7). The observation that hypertensive women matched with men for age, race, and mean arterial pressure have a lower blood pressure response to isometric exercise (8) suggests that alterations in vascular reactivity may be involved. It is well documented that women receiving estrogen replacement therapy have a 40–50% reduction in the risk of coronary artery disease (9–12). While a substantial portion of this effect has been attributed to estrogen-mediated increases in high-density lipoprotein-cholesterol and decreases in low-density lipoprotein-cholesterol (5, 13, 14), the alterations in lipid profile account for only about 50% of the cardiovascular benefit observed in estrogen-treated women, suggesting that other mechanisms may be involved (15).

Estrogen therapy appears beneficial for women suffering from cardiac syndrome X, a recently recognized disorder marked by severe angina in the presence of normal coronary arteries. This condition is associated with ovarian insufficiency (16) and a diminished endothelium-dependent hyperemic response (17). In one of the few studies to examine chronic estrogen administration, 2 months of 17 $\beta$ -estradiol treatment restored the hyperemic response to control values and markedly improved cardiac symptoms (16).

**2. Pregnancy.** Pregnancy is associated with widespread vascular changes that are important for ensuring fetal as well as maternal well-being. There is a fall in blood pressure and systemic vascular resistance, rise in cardiac output, and reduction in the pressor responses to several vasoconstrictors in humans, baboons, sheep, guinea pigs, and rats (18–22). The fall in vascular resistance is due to vasodilation in both the uterine and nonuterine circulations (23) and is accompanied by alterations in levels of vasoactive substances as well as changes in the vasoconstrictor and vasodilator responses of isolated vessel preparations. The ratios of several vasodilator relative to vasoconstrictor substances change with pregnancy. In normal human pregnancy, a rise in the circulating levels of prostacyclin relative to thromboxane occurs with advancing gestation (24). In rats, pregnancy is associated with increases in the circulating levels and urinary excretion of nitrate and cGMP, the second messenger involved in prompting NO-induced vascular smooth muscle relaxation (Fig. 1) (25). In a study on the effect of intravenous




FIG. 1. Biosynthesis of NO. Shown here are potential mechanisms by which estrogen may modulate endothelium-dependent vasodilation through 1) effects on number and/or sensitivity of cholinergic receptors, 2) modification of  $Ca^{++}$  influx, and 3) induction of NO synthase. Flow, shear stress, or receptor activation by acetylcholine or bradykinin results in influx of  $Ca^{++}$ . Increased intracellular  $Ca^{++}$  stimulates the constitutive NO synthase. The NO formed from L-arginine diffuses to smooth muscle cells, stimulating guanylate cyclase and resulting in increased synthesis of cGMP from GTP. Nitro-L-arginine methyl ester (L-NAME) is a competitive inhibitor of NO synthase and causes vasoconstriction *in vitro*.

infusion of glyceryl dinitrate (GTN), an NO donor *in vivo*, and prostacyclin on uterine artery blood flow during human pregnancy, GTN administered during the first trimester decreased uterine vascular resistance while prostacyclin had no effect (26). The GTN-associated increase in uterine artery blood flow occurred without significant alterations in heart rate or systemic blood pressure and mimicked the normal physiological increase observed during the second trimester, suggesting a potential role for NO in uterine artery vasodilation. The effect of pregnancy on the vasoconstrictor, ET-1, is unclear; increases, decreases, as well as no differences between pregnant and nonpregnant women have been reported (2, 27, 28).

#### B. Speculations about involvement of female hormones in vascular disease

Migraine headaches, primary pulmonary hypertension, preeclampsia, and Raynaud's phenomenon are a diverse group of vascular disorders. A range of etiological mechanisms have been implicated in their pathogenesis, including alterations in central nervous system neurons and pain-modulating pathways, immunological function, and vascular remodeling (29–31). However, a number of features are common to these diseases. They all demonstrate a marked female preponderance (31–34), or, in the case of preeclampsia, occur exclusively in women. There is an overlap in their occurrence such that the presence of one disease increases the risk of developing another of these conditions (35, 36). Women with severe migraines are at an increased risk of developing preeclampsia (37). Migraine or Raynaud's phenomenon predisposes women to variant angina, a condition characterized by

transient coronary arterial vasospasm (38). The frequency of Raynaud's is greater among women with primary pulmonary hypertension than in the general population in most (31, 39–43) but not all studies (44).

Alterations in vasoactive substances and  $\text{Ca}^{++}$  homeostasis are present in many if not all of these disorders. Increased thromboxane relative to prostacyclin has been described in patients with preeclampsia, primary pulmonary hypertension, and Raynaud's phenomenon (45–47). In preeclampsia, a disease of pregnancy characterized by systemic hypertension, edema, proteinuria, and coagulation abnormalities (48), the increased ratio of vasodilator to vasoconstrictor prostaglandins of normal pregnancy is reversed in proportion to the severity of disease (49, 50). Further, a decrease in the circulating levels of the NO metabolite, nitrite, has also been noted in preeclamptic patients (51). Pregnant rats treated chronically with the NO inhibitor, nitro-L-arginine methyl ester, developed hypertension and proteinuria, and delivered growth-retarded pups (52). Increased levels of ET-1 and reduced expression of NO synthase have been reported in the lungs of patients with primary pulmonary hypertension (53, 54). A role for alterations in  $\text{Ca}^{++}$  homeostasis is suggested by the therapeutic benefits of  $\text{Ca}^{++}$  channel blockers in Raynaud's phenomenon, primary pulmonary hypertension, and migraines (29, 55, 56).  $\text{Ca}^{++}$  supplementation has also been associated with a decreased incidence of preeclampsia (57). When compared with normotensive controls, platelets from preeclamptic women have increased intracellular  $\text{Ca}^{++}$  concentration (58, 59). Further, the observation of a greater contractile response to the  $\text{Ca}^{++}$  channel agonist BayK8644 in inferior epigastric arteries from preeclamptic women suggests that voltage-gated  $\text{Ca}^{++}$  channel activity may be enhanced in comparison to normal women (60).

Despite the clear female preponderance of these disorders, the mechanisms by which female sex steroids influence their occurrence remains speculative. Studies reviewed below and in *Section II.C* suggest that such mechanisms may involve alterations in sex hormone levels and/or ratios and hormonal stimulation of vasoconstrictor pathways.

Surprisingly little is known about the role of sex steroids, their ratios, or the levels of their metabolites in the development of preeclampsia. Preeclampsia is characterized by an increase in systemic vascular resistance, vascular reactivity, and alterations in pelvic blood flow distribution that precede the onset of hypertension, suggesting a failure of the normal vascular adjustments to pregnancy (61–64). In addition to alterations in endothelial cell vasoactive substances, evidence of sympathetic activation has also been reported in preeclamptic compared with normal pregnant women (65, 66). Older studies suggesting an association between preeclampsia and low serum  $17\beta$ -estradiol levels were limited by the small number of subjects, the absence of rigorous diagnostic criteria for preeclampsia, and the likelihood that the low estradiol levels were due to other obstetrical conditions (67, 68). We have recently described a decrease in serum estradiol and increase in serum progesterone before the onset of hypertension in preeclamptic women residing at high altitude (3100 m), where the incidence of preeclampsia is markedly increased in comparison with low altitude (1600 m) (69).

Migraine headaches tend to cluster just before or during menstruation in a majority of affected women and exclusively with the menses in a substantial proportion (29, 32–34). In addition to the alterations in estradiol and progesterone levels, both of which have been shown to be higher in women with migraines than in asymptomatic controls (32, 70, 71), fluctuations in hormone levels or, in particular, estrogen withdrawal may be important (72, 73). In a small descriptive study, headaches began immediately before menses, and their onset was delayed by estrogen treatment (73). Additionally, in two controlled trials, percutaneous estradiol gel reduced the frequency of migraine headaches (74, 75). Further, the vast majority (86%) of menstrual migraines improve with pregnancy, a time when steroid hormone levels rise and their fluctuations are eliminated (74, 75). Thus, while estrogen withdrawal appears to be associated with the development of menstrual migraines, whether these effects are mediated through alterations in vascular function is unknown.

A link between hormonal fluctuations and symptoms of Raynaud's phenomenon is suggested by the observation that whereas normal women decrease their digital blood flow response to cold during the follicular compared with the luteal phase of the menstrual cycle, women with Raynaud's show no such variation (76). Further studies are needed to characterize the nature of these hormonal alterations or how they may be related to the altered vasoreactivity seen in Raynaud's phenomenon.

Estrogen and progesterone may also influence vasoconstriction depending on the vascular bed and the site of action (see *Section II.C*). Such effects may contribute to the gender-related differences in certain vascular diseases. Studies in isolated rat lungs, for instance, indicate that estradiol enhances vasoconstriction to the thromboxane mimetic U46619 which may, in turn, explain the increased vasoconstrictor response noted in the female compared with male rat lung (77). This observation may be relevant for understanding the association between primary pulmonary hypertension and female hormones. It is well known that primary pulmonary hypertension occurs twice as often in women as men (31). Further, it is associated with significant mortality during pregnancy (78, 79). In 23 women that presented with primary pulmonary hypertension before becoming pregnant, half died while pregnant, with a majority of the deaths occurring during delivery (79). Most authors attribute maternal demise to the collapse of a compromised maternal circulation further stressed by the hemodynamic changes of labor and delivery, but pregnancy-associated increases in hormones and/or their decline at the time of delivery may also be important.

### C. Summary

Female hormones are linked both to protection and susceptibility to vascular disease. There are several possibilities by which this apparent contradiction might be resolved. One possibility concerns the levels of female hormones. While fluctuations in hormone levels appeared to be associated with the onset of menstrual migraines and primary pulmonary hypertension in some studies, there is not sufficient information to judge whether there are differences in circulating levels of estrogen and progesterone, their fluctuations,

or their ratios in women with vasospastic conditions compared with well matched controls. A second possibility concerns the separate *vs.* combined effects of the hormones involved. It is well recognized that estrogen and progesterone can act in ways that are antagonistic to each other. For example, in isolated dog coronary artery rings, progesterone attenuates estrogen-induced stimulation of the endothelium-dependent responses (80). These antagonistic effects may, in turn, be receptor-mediated. In reproductive tissues such as the uterus and breast, estrogen has been shown to up-regulate progesterone receptors while progesterone down-regulates its own receptor (81–86). It has also been demonstrated that in the chick oviduct and rodent uterus, progesterone prevents the continuation of an estrogen effect (87–89), possibly by interfering with the replenishment of cytosolic estrogen receptors (90–92). A third possibility concerns the site of hormonal action. Since the same action, namely an increase in intracellular  $\text{Ca}^{++}$  promotes vasodilation or vasoconstriction depending on whether it occurs in the endothelial or vascular smooth muscle cell, respectively, one possibility for this apparent paradox is that female hormones have endothelial *vs.* vascular smooth muscle cell sites of action in different disease states.

### III. Potential Mechanisms of Hormone-Mediated Effects

#### A. Receptor-mediated actions

Estrogen and progesterone are lipophilic molecules whose receptor-mediated modes of action have been well studied in the breast, uterus, and brain (86, 93–95). Expression of estrogen and progesterone receptors in the vasculature has been controversial, perhaps reflecting intervessel and/or species variability as well as limitations of the different techniques employed.

Using radioactive ligand binding, estrogen and progesterone receptors have been demonstrated in cytosolic fractions of the canine aorta and inferior vena cava (96). Immunocytochemistry has been used to demonstrate the presence of estrogen receptors in guinea pig uterine artery vascular smooth muscle and endothelial cells (97). While estrogen and progesterone receptors have been found in rabbit and human uterine artery vascular smooth cells (98), these studies were not able to demonstrate receptors in rabbit or guinea pig aortas or rabbit renal, femoral, hepatic, or pulmonary arteries using the same technique.

More recently, studies have investigated both the presence and functional status of steroid receptors in vascular tissue. Using Northern analysis and a monoclonal antiestrogen receptor antibody that recognizes the DNA-binding domain of the estrogen receptor, Orimo *et al.* (99) demonstrated estrogen receptor mRNA and protein, respectively, in cultured rat aortic vascular smooth muscle cells. The authors also showed that a concentration of  $17\beta$ -estradiol at the high end of the physiological range (10 nM) up-regulated the expression of estrogen-dependent *c-fos* mRNA, suggesting estrogen stimulation of gene transcription in vascular smooth muscle. Using the ribonuclease protection assay and immunofluorescence techniques, others have demonstrated low levels of

estrogen receptor mRNA and protein in human vascular smooth muscle cells derived from the mammary artery and saphenous vein (100). The low level of estrogen receptor found in these cells may be related to the use of cell cultures. The concentration of estrogen receptor decreases markedly over time in cultured rabbit uterine myocytes (101). Functional integrity of the receptor was demonstrated by transfection of vascular smooth muscle cells with a reporter plasmid containing an estrogen response element driving the expression of the luciferase gene (ERE-Luc) or the control plasmid lacking the estrogen response element (Tk-Luc) to  $17\beta$ -estradiol. Vascular smooth muscle cells containing the luciferase gene, but not the control cells, had a dose-dependent increase in activity of the ERE-Luc plasmid after exposure to  $17\beta$ -estradiol (100).

Some studies suggest that receptor levels within the vasculature are altered by changes in the hormonal milieu. Spiral arteries of the human uterus demonstrate intense staining of both estrogen and progesterone receptors during the first trimester of pregnancy (102). While immunocytochemistry revealed no estrogen receptor in guinea pig heart, aorta, and carotid specimens, receptors were demonstrated in the uterine artery (97). In addition, a midpregnancy fall in estrogen receptor levels compared with nonpregnant or term animals was demonstrated, which may have been due to a rise in serum progesterone relative to estradiol levels (97). The recent observation that estrogen receptors are diminished in coronary arteries of premenopausal women with atherosclerosis compared with normal premenopausal controls suggests that receptor levels may also be altered in disease states (103).

#### B. Effects of estrogen and progesterone on vascular function

1. *Cardiovascular disease.* The protective effect of female hormones in coronary artery disease may be due to the effects of estradiol. Several studies have reported that estradiol administration increases endothelium-dependent vasodilation in response to acetylcholine stimulation in the coronary circulation (Fig. 1). Using quantitative coronary angiography in monkeys fed a high cholesterol diet, Williams *et al.* (104) demonstrated that infusion of the endothelium-dependent vasodilator acetylcholine constricted the coronary arteries of ovariectomized animals but dilated those of an estrogen-treated group. Although the estrogen-treated group had reduced plaque formation, the altered vasoreactivity did not correlate with the degree of plaque present. These findings have been supported by studies in postmenopausal women with evidence of endothelial cell dysfunction as indicated by a vasoconstrictor response to acetylcholine. Administration of physiological levels of  $17\beta$ -estradiol and the synthetic steroid, ethinyl estradiol, by intraarterial or intracoronary infusion, potentiated vasodilation to acetylcholine in large and microvascular coronary arteries as well as in the forearm circulation (105–107). Estradiol also increased the vasodilator response to sodium nitroprusside, an endothelium-independent vasodilator, in the forearm but not the coronary circulation (105), suggesting that estradiol influences vascular smooth muscle sensitivity in some vascular beds. At supra-physiological doses, estradiol or progesterone caused relax-

ation of precontracted, endothelium-denuded rabbit isolated coronary artery rings, also supporting the involvement of endothelial-independent actions (108, 109). The extent to which baseline circulatory parameters are affected is unclear; the synthetic steroid, ethinyl estradiol, but not  $17\beta$ -estradiol, increased baseline coronary blood flow and decreased resistance (106, 107).

2. *Pregnancy.* A vasodilatory effect of estradiol in the sheep uterine circulation has long been recognized. Acute or chronic administration of  $17\beta$ -estradiol to oophorectomized ewes reduces systemic and uterine vascular resistance, markedly increases uterine blood flow, and decreases the pressor response to angiotensin II (110–113). However, the vasodilatory effects of estradiol administration are not consistent across species or study preparations. Whereas acute intravenous administration of  $17\beta$ -estradiol decreased the systemic pressor response to angiotensin II in oophorectomized rats (114), 2–3 week treatment with subcutaneous pellets or intraperitoneal injection did not have similar effects (115–117). Even within a species, the effects of chronic estradiol administration differ in the uterine and nonuterine circulations. Systemic vasodilation and refractoriness to angiotensin II in sheep were maintained throughout (and in the case of angiotensin II, even beyond the infusion period), but uterine values returned to baseline after 7 days. Given the prolonged nature of the hormone stimulation of pregnancy and the consistency among species in pregnancy's vasodilatory effects, these discrepancies suggest that estradiol alone cannot account for pregnancy-associated vasodilation. The possible involvement of progesterone is supported by studies in which progesterone alone or in combination with estrogen decreased pressor response to angiotensin II in whole animals or isolated tail arteries of rats (115, 118). However, other studies in the rat using more physiological levels of hormone showed no effect (114, 116, 117).

$\text{Ca}^{++}$  channels may play a role in the physiological changes observed during pregnancy. Voltage-gated  $\text{Ca}^{++}$  channels are activated by depolarization of the plasma membrane when extracellular  $[\text{K}^+]$  is increased (Fig. 6). In isolated porcine uterine arteries, pregnancy and estradiol-treatment decreased the contractile response to potassium chloride. That this decrease was associated with a reduction in  $\text{Ca}^{++}$  uptake into uterine vascular smooth cells (119–121) suggests that pregnancy and estradiol may reduce intracellular  $\text{Ca}^{++}$  via an effect on voltage-gated  $\text{Ca}^{++}$  channels.

### C. Steroid-mediated effects on endothelial and vascular smooth muscle cell function

Included among the mechanisms that modulate vascular tone and reactivity are alterations in endothelial vasodilator or vasoconstrictor activity and vascular smooth muscle cell function. Reviewed below are several lines of evidence suggesting that estrogen and progesterone influence the production or activity of the endothelial-derived substances, NO, ET-1, and eicosanoids. There are a number of other endothelial-derived factors that represent potential sites of influence that are beyond the scope of this review, including endothelial-derived hyperpolarizing factor, angiotensin II,

and platelet-activating factor (Table 1). In addition, since this review is primarily concerned with vascular reactivity, factors important in angiogenesis such as vascular endothelial growth factor (VEGF) (122) are not discussed.

1. *NO.* Endothelial-derived NO mediates vascular smooth muscle relaxation by stimulating soluble guanylate cyclase (Fig. 1). NO is produced by the endothelium under basal conditions as well as upon stimulation by agonists such as acetylcholine and bradykinin that act via endothelial surface membrane receptors (Fig. 1 and Table 1), some of which are linked to G protein activation (123, 124). Vascular smooth muscle, neural, and other cell types can, under certain circumstances, also produce NO (125). Studies from a broad range of experimental preparations suggest that physiological alterations in female hormones modulate endothelial cell function through effects on basal as well as stimulated NO production and/or activity.

Female gender has been associated with increased basal NO release. Aortic rings isolated from female compared with male rabbits and precontracted with phenylephrine showed greater relaxation to the inhibitor of NO inactivation, superoxide dismutase, and greater constriction after inhibition of NO synthesis by nitro-methyl-L-arginine (126). Oophorectomy abolished this gender difference, suggesting that physiological levels of female hormones stimulate basal NO activity. The absence of gender differences in relaxation to acetylcholine indicated that stimulated NO activity was not affected.

Pregnancy has been associated with increased stimulated release of NO. Guinea pig uterine artery and rat aortic rings demonstrate increased relaxation to acetylcholine during pregnancy (127–129). These effects of pregnancy are similar to those observed with acute and chronic estradiol treatment, suggesting that the pregnancy-associated changes are

TABLE 1. Endothelial targets for actions of estrogen and progesterone

I. Endothelial Derived Products	
Vasodilators	Vasoconstrictors
PGE <sub>2</sub>	Angiotensin II
PGI <sub>2</sub>	Endothelin 1
Nitric oxide	Arachadonic acid
Histamine	PGH <sub>2</sub>
PGH <sub>2</sub>	Thromboxane A <sub>2</sub>
Platelet activating factor	Histamine
Superoxide anion	Superoxide anion
Leukotrienes	Platelet activating factor
EDHF	Leukotrienes
II. Endothelial Membrane Receptors	
Acetylcholine	
Bradykinin	
ATP/ADP	
Serotonin	
Thrombin	
$\alpha_2$ -Adrenergic	
Endothelin B (ET <sub>B</sub> )	
Histamine	
Thromboxane A <sub>2</sub>	
Angiotensin II	
Vasopressin	
Substance P	
Vascular endothelial growth factor (VEGF)	

mediated, at least in part, by estradiol. In oophorectomized ewes, 80% of the rise in uterine blood flow with acute estradiol infusion was blocked by the NO synthesis inhibitor, nitro-L-arginine methyl ester (130). The possibility that estradiol increased NO synthesis was supported by the observation that the inhibition of the uterine blood flow response could be reversed with excess administration of L-arginine (Fig. 1). Chronic treatment with 17 $\beta$ -estradiol also increased relaxation to acetylcholine in isolated rabbit femoral artery rings and aortas from spontaneously hypertensive rats (131, 132). Not all studies support that estradiol-induced vasodilation is mediated through NO; estradiol relaxed precontracted rabbit isolated coronary artery rings, and this relaxation was reversed by NO inhibition in some but not other studies (133, 134).

Estradiol may stimulate endothelium-dependent relaxation through both receptor-mediated and non-receptor-mediated mechanisms. Chronic treatment with 17 $\beta$ -estradiol increased relaxation to acetylcholine in isolated rabbit femoral artery rings and aortas from spontaneously hypertensive rats (131, 132). Increased relaxation to acetylcholine but not to non-receptor-mediated substances such as the  $\text{Ca}^{++}$  ionophore A23187 has been observed in rabbit femoral artery rings, suggesting that chronic estrogen treatment may increase the number or sensitivity of muscarinic receptors (132). However, increased relaxation to A23187 in rabbit isolated aortic rings indicated that non-receptor-mediated mechanisms of NO release may also be present in some species or vessels (135).

The stimulatory effects of pregnancy and estradiol treatment on NO activity may take place by increasing endothelial constitutive NO synthase (NOS III) production or activity. Pregnancy has been associated with a 4-fold increase in  $\text{Ca}^{++}$  dependent NOS III activity in the guinea pig uterine artery as measured by conversion of radiolabeled L-arginine to citrulline (136). Using semiquantitative polymerase chain reaction, increased relaxation to acetylcholine in the pregnant rat aorta correlated with a 2-fold increase in NOS III mRNA in pregnant compared with nonpregnant rings (129). Five-day intraperitoneal injection of 17 $\beta$ -estradiol to gonadally intact female guinea pigs also increased NOS III activity in heart, kidney, esophagus, and skeletal muscle, while progesterone had no effect (136). Additionally, 17 $\beta$ -estradiol increased NO production and protein levels of NOS III in cultured human aortic endothelial cells and up-regulated NOS III gene expression in the aorta of gonadectomized rats (129, 137). Of interest, the effect of estradiol on NOS III was only 50% of that seen with pregnancy, suggesting that other factors during pregnancy also influence expression of this gene. Finally, the presence of two left-half palindromic sites of an estrogen receptor-binding element on the human endothelial NOS gene supports a potential receptor-mediated effect of estrogen on gene expression (138).

Thus, data in several species suggest that estradiol modulates endothelium-dependent relaxation in a number of vascular beds through increased NO activity which, in turn, may be due to up-regulation of NOS III gene expression. However, the stimulation observed during pregnancy is not fully reproduced by estradiol treatment and the effects of estradiol treatment are not as generalized among vascular

beds as those observed during pregnancy. Thus, other mechanisms may also be involved, such as increased sensitivity or affinity of endothelial receptors or effects on intermediary steps involved in receptor-mediated NO release, such as modulation of G-protein activation.

2. *ET-1*. In subnanomolar concentrations, ET-1 is a potent endothelial-derived vasoconstrictor, but at lower concentrations, it can also induce transient vasodilation (139). Two classes of ET-1 receptors, ET-1 A ( $\text{ET}_A$ ) and ET-1 B ( $\text{ET}_B$ ), both of which appear to be G-protein linked, have been identified in vascular beds. Activation of  $\text{ET}_A$  in vascular smooth muscle cells causes constriction which, in turn, is mediated by  $\text{Ca}^{++}$  channel activation and activation of phospholipase C (Figs. 2 and 3).  $\text{ET}_B$  activation on endothelial cells may also




FIG. 2. Simplified schema of receptor-mediated mechanism of smooth muscle contraction. Estrogen may directly inhibit voltage-gated  $\text{Ca}^{++}$  channels in vascular smooth muscle cells, thereby decreasing  $\text{Ca}^{++}$  influx. Progesterone may increase  $\text{Ca}^{++}$  entry through activation of these channels. Female sex steroids may also influence the sensitivity of contractile proteins such as myosin light chain kinase (MLCK\*) and calmodulin ( $\text{CaM}^*$ ) to  $\text{Ca}^{++}$ . Binding of agonist to a receptor linked to a G protein (G) results in activation of phospholipase C (PLC). PLC hydrolyzes a membrane phospholipid phosphatidyl-inositol-4,5-bisphosphate ( $\text{PIP}_2$ ) into two second messengers: diacylglycerol (DAG) and inositol-1,4,5-trisphosphate ( $\text{InSP}_{1,4,5}$ ).  $\text{InSP}_{1,4,5}$  diffuses across the cytoplasm triggering the release of intracellular  $\text{Ca}^{++}$  from internal stores. Increased cytoplasmic  $\text{Ca}^{++}$  promotes activation of the  $\text{Ca}^{++}$  binding protein calmodulin ( $\text{CaM}^*$ ), which in turn activates myosin light chain kinase (MLCK\*). Muscle contraction is initiated with myosin phosphorylation. DAG activates protein kinase C (PKC), which in turn phosphorylates a number of proteins resulting in smooth muscle contraction. Agonists may also activate  $\text{Ca}^{++}$  channels through effects of second messengers such as  $\text{InSP}_{1,4,5}$  and PKC.




FIG. 3. Biosynthesis of ET-1. ET-1 is synthesized from prepro-ET-1. The mechanism(s) by which estrogen decreases ET-1 levels is unknown but may involve effects on gene transcription of important enzymes such as endothelin-converting enzyme (ECE).

stimulate NO and prostacyclin release. In cultured bovine and porcine endothelial cells, ET-1 release can be stimulated by a number of agonists including thrombin, arginine vasopressin, angiotensin II, and A23187 (140, 141). In the porcine isolated aorta, increased NO resulting from the inhibition of the NO scavenger, superoxide dismutase, or the nonhydrolyzable analog of cGMP, 8-bromo-cGMP, reduced thrombin-stimulated ET-1 production (142). Thus, a dynamic balance exists, involving feedback relations between NO and ET-1 such that ET-1 stimulates NO which, in turn, reduces ET-1 production. ET-1 may also stimulate prostacyclin release. In cultured human umbilical vein endothelial cells, high doses of ET-1 ( $10^{-9}$  M) stimulated prostacyclin production as measured by increased 6-keto-prostaglandin  $F_{1\alpha}$  formation (143).

Evidence for steroid-mediated effects on ET-1 is limited. Doses of  $17\beta$ -estradiol 1000-fold greater than the normal physiological range (3–30  $\mu$ M) decreased the contractile response to ET-1 in isolated rabbit coronary artery rings (134). However, the decrease was similar in rings from male and nonpregnant female animals, and there were no gender differences in the magnitude of the ET-1-induced contraction. Further, nitro-L-arginine methyl ester did not alter the response to  $17\beta$ -estradiol, suggesting that estradiol acted via an endothelium-independent mechanism. Consistent with this possibility, similar doses of  $17\beta$ -estradiol (10–30  $\mu$ M) decreased the contractile response to ET-1 in endothelium-denuded coronary arteries from male and female animals. The authors hypothesized that estradiol affected vascular smooth muscle cell  $Ca^{++}$  channels since estradiol reduced the contractile response to BAY K8644, a specific agonist of voltage-dependent  $Ca^{++}$  channels (134). While  $17\beta$ -estradiol ( $10^{-8}$  M) potentiated the ET-1 increase in prostacyclin production in human umbilical vein endothelial cells, the mechanism remains unknown (143).

The effects of pregnancy and physiological alterations in estradiol or progesterone on ET-1 have not been fully explored. Among the possibilities requiring further investigation are direct effects of pregnancy and female hormones on vascular smooth muscle cell response to ET-1 stimulation and indirect effects on NO activity which, in turn, exert a feedback inhibition on ET-1 production.

**3. Eicosanoids.** Endothelial cells metabolize arachadonic acid to produce eicosanoids through cyclooxygenase, peroxidase,

and lipoxygenase enzymatic pathways (Fig. 4). In general, vasodilation is mediated by prostacyclin and prostaglandin  $E_2$  while thromboxane  $A_2$ , prostaglandin  $F_{2\alpha}$ , and some leukotrienes are vasoconstrictors (144). Vascular smooth muscle cell relaxation results from formation of cAMP and activation of cAMP-dependent protein kinase (Fig. 5). Thromboxane  $A_2$  is thought to antagonize the effects of prostacyclin through inositol-1,4,5-triphosphate ( $InsP_{1,4,5}$ )-mediated release of intracellular  $Ca^{++}$  (Fig. 2).

Conflicting evidence exists regarding the effects of physiological alterations in estrogen or progesterone on eicosanoid production in the vasculature. Two- to three-week treatment with intraperitoneal injections of estradiol increased prostacyclin release in isolated aortic rings of male and intact female rats as measured by elevated levels of the stable metabolite 6-keto-prostaglandin  $F_{1\alpha}$  (145) (whether blood estradiol levels obtained were within the physiological range is unknown). Physiological concentrations of  $17\beta$ -estradiol have been shown to increase prostacyclin production and biosynthetic activity in cultured rat and rabbit aortic smooth muscle cells (145, 146). Estradiol also appears to stimulate the production of contractile prostanoids, as demonstrated by the observation that indomethacin reduced the estradiol-induced increase in contractile response to norepinephrine in isolated rabbit aortic rings (135). However, cyclooxygenase inhibition did not reverse the vasodilator effects of pregnancy in whole animals (21, 22), or in isolated




FIG. 4. Endothelial cells produce eicosanoids from arachadonic acid through the cyclooxygenase, lipoxygenase, and peroxidase pathways. Primary end products include thromboxane  $A_2$  (TXA<sub>2</sub>), prostaglandin  $E_2$  (PGE<sub>2</sub>), and prostacyclin (PGI<sub>2</sub>) from the cyclooxygenase pathway and leukotrienes from the lipoxygenase pathway. Hydroxyperoxyeicosatetraenoic acid (II-HPETE) and hydroxyeicosatetraenoic acid (HETE) are intermediate products in the cyclooxygenase and peroxidase pathways. Precise mechanisms by which estradiol increases PGI<sub>2</sub> production remain unclear.






FIG. 5. Simplified schema of cAMP pathway of smooth muscle relaxation. Estrogen has been shown to modify expression of  $\beta$ -receptors in a number of tissues and may stimulate prostacyclin production in endothelial cells. Key proteins in the pathway of vascular smooth muscle cell relaxation include hormone receptors (Rec), a stimulatory protein (Gs), catalytic adenylyl cyclase (AC), and phosphodiesterase (PDE) that hydrolyzes cAMP. cAMP stimulates cAMP-dependent protein kinases (PK-A, PK-G) that are involved in the phosphorylation of myosin light chain kinase (MLCK) triggering relaxation.

rabbit femoral or coronary artery rings (108, 132), suggesting that estradiol-mediated vasodilation is not dependent on cyclooxygenase products (Fig. 4). Since most studies look at single blockers and do not examine interactions between the cyclooxygenase and lipooxygenase pathways, the possibility that estradiol or pregnancy acts on more than one eicosanoid pathway to affect the balance of vasodilator to vasoconstrictor eicosanoids has not been investigated. The effects of female hormones on endothelial cells may also differ from those seen in vascular smooth muscle cells; supraphysiological doses of estradiol did not alter prostacyclin or thromboxane B2 production in cultured human umbilical vein endothelial cells (143).

**4. Adrenergic responsiveness.** Most (21, 128, 147) but not all (148) studies suggest that pregnancy decreases contractile response to adrenergic stimulation in a broad range of species and study preparations. However, the effects of estrogen and progesterone on adrenergic responsiveness vary among species and vascular beds. Femoral artery rings from oophorectomized female rabbits treated chronically with physiological concentrations of 17 $\beta$ -estradiol decreased their contractile response to norepinephrine (149) while mesenteric arteries from female and male rats administered higher dosages of estrogen demonstrated increased adrenergic sensitivity (150). Few studies have examined the effect of chronic progesterone treatment on adrenergic responses. In the perfused tail artery of female rats, 3-day treatment with supraphysiological levels of progesterone decreased the pressor response to norepinephrine while estradiol treatment had no effect (118). The results from these studies may reflect not only interspecies and vessel differences but also effects of varying dosages of hormones as well as the use of gonadally

intact *vs.* castrated preparations. Additionally, the hormones may be exerting their effects at alternate sites; endothelial  $\alpha_2$ -receptor stimulation causes NO release and vasodilation, while vascular smooth muscle  $\alpha_1$ - and  $\alpha_2$ -receptor activation produces vasoconstriction (151, 152).

Ovarian steroids may influence adrenergic responsiveness via alterations in adrenergic receptor affinity. Both estrogen and progesterone appear to increase  $\alpha_1$ -receptor affinity. The mesenteric arteries of estrogen-treated male and female rats showed an increased contractile response to adrenergic stimulation that was associated with increased  $\alpha_1$ -adrenergic receptor affinity (150). In the gilt uterine artery, decreased uterine blood flow during the luteal (high progesterone) phase was associated with increased  $\alpha_1$ -receptor binding affinity (153).

An additional potential mechanism of steroid-mediated effects involves modification of catecholamine levels via alterations in uptake, storage, and efflux from nerve endings (147, 154). Sensitivity to norepinephrine was increased in pregnant compared with nonpregnant rat mesenteric arteries in the presence of cocaine, an inhibitor of neuronal reuptake, suggesting that pregnancy may be associated with increased neuronal reuptake and/or deactivation of norepinephrine (147). Studies in canine saphenous veins suggest that estradiol may also stimulate norepinephrine release from nerve endings (154). Less is known concerning the effects of female hormones on  $\beta$ -receptors. While  $\beta_1$  and  $\beta_2$  have been identified in cultured bovine pulmonary artery endothelial cells (155), and ovarian hormones have been shown to modify the number and/or affinity of  $\beta$ -adrenergic receptors in the myometrium, brain, lung, and heart (156, 157), little is known about their effects in the vasculature.

Thus, while pregnancy and ovarian steroids appear to modulate adrenergic responses in a number of vascular beds, their specific effects on endothelial *vs.* vascular smooth muscle cells await study.

**5.  $Ca^{++}$  homeostasis.**  $Ca^{++}$  homeostasis is central to both endothelial and vascular smooth muscle cell function; endothelial release of NO and prostacyclin is  $Ca^{++}$  dependent (158), and increased intracellular  $Ca^{++}$  prompts vascular smooth muscle contraction (159). The regulation of cytosolic  $Ca^{++}$  in both cell types is a function of extracellular  $Ca^{++}$  influx, intracellular  $Ca^{++}$  release or reuptake, and intracellular  $Ca^{++}$  efflux (Fig. 6). Ion channels play a key role in both cell types. In the vascular smooth muscle, the predominant current is carried by voltage-dependent  $Ca^{++}$  channels that are activated by depolarization of the plasmalemma (160). Voltage-gated channels have not been identified in large vessel endothelium although they appear to exist in microvascular endothelial cells (161).  $Ca^{++}$  entry also occurs through leak channels, stretch-activated channels, and receptor-operated channels (158) (Fig. 6).

Little is known about the effects of estrogen and progesterone on ion channels in vascular endothelial and smooth muscle cells. Studies in other cell types suggest that estrogen and progesterone influence the expression and activity of ion channels through genomic and nongenomic mechanisms. In myometrial smooth muscle cells, estrogen and progesterone act on ion channels through the classic mechanism of gene






FIG. 6. Mechanism of calcium homeostasis. Ovarian steroids may inhibit voltage-gated  $\text{Ca}^{++}$  channels at the cellular membrane or influence synthesis of  $\text{Ca}^{++}$ ,  $\text{K}^{+}$ , and  $\text{Na}^{+}$  channels via gene transcription.  $\text{Ca}^{++}$  entry may occur through leak (L), stretch-activated (S), voltage-dependent (V), or receptor-activated (R) channels.  $\text{Ca}^{++}$  may be extruded from cell by  $\text{Ca}^{++}\text{-H}^{+}$  ATPase and  $\text{Na}^{+}\text{-Ca}^{++}$  exchange. Elevated intracellular  $\text{Ca}^{++}$  may be sequestered into the mitochondrion (MIT) or the endoplasmic reticulum (ER) via an ATP- $\text{Ca}^{++}$  pump. Other factors that influence  $\text{Ca}^{++}$  homeostasis by regulating membrane potential include  $\text{K}^{+}$  and voltage-gated  $\text{Na}^{+}$  channels as well as the  $\text{Na}^{+}\text{-K}^{+}\text{-Cl}^{-}$  antiporter, the  $\text{Na}^{+}\text{-H}^{+}$  exchanger, and the  $\text{Na}^{+}\text{-K}^{+}$  pump.  $\text{K}^{+}$  channels identified in vascular smooth muscle cells include receptor-operated, voltage-dependent, and independent channels.

transcription. Three-day progesterone treatment increases the number and/or density of  $\text{Ca}^{++}$  channels and increases  $\text{Ca}^{++}$  current in cultured myometrial cells (162). Activation of the progesterone receptor is likely involved, since progesterone injection did not increase  $\text{Ca}^{++}$  current in the presence of the progesterone antagonist RU486. Chronic estrogen treatment has also been shown to increase the number of nitrendipine binding sites in myometrial smooth muscle cell membranes (nitrendipine is a dihydropyridine  $\text{Ca}^{++}$  channel blocker) (163), suggesting an estrogen-induced increase in the number of voltage-gated  $\text{Ca}^{++}$  channels.

Although estrogen and progesterone receptors have been demonstrated in a number of vascular beds, it is unclear whether the effects of these hormones are exerted through genomic or nongenomic mechanisms and, if genomic, the specific gene products involved. The immediate effects of estrogen and progesterone on  $\text{Ca}^{++}$  homeostasis in vascular as well as nonvascular tissue support the existence of nongenomic influences. Acute administration of progesterone and estradiol alone caused relaxation in endothelium-denuded rabbit coronary artery rings (108, 109) precontracted with Bay K8644, a voltage-gated  $\text{Ca}^{++}$  channel agonist. Incubation with  $17\beta$ -estradiol shifted the  $\text{Ca}^{++}$ -dependent concentration curve to the right in a high  $\text{K}^{+}$  depolarization medium (108). These observations suggest that the female hormones may alter vascular smooth muscle  $\text{Ca}^{++}$  influx through effects on voltage-gated  $\text{Ca}^{++}$  channels (Fig. 6). In support of this possibility, using the whole cell patch-clamp technique,  $17\beta$ -estradiol ( $10\text{ }\mu\text{M}$ ) was shown to have an inhibitory effect on voltage-dependent  $\text{Ca}^{++}$  channels in the A7r5 vascular smooth muscle cell culture line (164).

In nonvascular tissue, progesterone inhibited  $\text{Ca}^{++}$ -induced contraction in uterine smooth muscle cells within 1–3 min of  $\text{Ca}^{++}$  addition, and the inhibitory effect was immediately reversed when the hormone was washed out (165). In guinea pig myocytes,  $17\beta$ -estradiol decreased intracellular  $\text{Ca}^{++}$  within 1 min of administration and inhibited  $\text{Ca}^{++}$  current in a dose-dependent fashion (166), suggesting an immediate, negative inotropic effect in these cells. These acute effects of ovarian hormones may occur at the cell membrane. Human sperm cells have been shown to have progesterone receptors in the cell membrane (167). In these cells, acute progesterone administration stimulates rapid  $\text{Ca}^{++}$  influx (168, 169). The progesterone receptor has been hypothesized to interact with  $\text{Ca}^{++}$  channels directly or via a cell membrane G protein. The membrane receptor appeared to be different from the classic intranuclear receptor because the progesterone receptor blockers RU38486 and ZK98,299 were ineffective in inhibiting the progesterone-induced  $\text{Ca}^{++}$  influx.  $\text{K}^{+}$  and  $\text{Na}^{+}$  channels also play a role in altering intracellular  $\text{Ca}^{++}$  levels in nonvascular tissues. In isolated rat myometrial smooth muscle cells, hormonal status of the animal influenced the expression of  $\text{K}^{+}$  channels (170). Rat myometrial cells have also been shown to express putative  $\text{K}^{+}$  channel mRNA after chronic estrogen treatment of the animal and to undergo an increase in fast  $\text{Na}^{+}$  channels density during gestation (171, 172). Whether similar changes occur in vascular smooth muscle or endothelial cells is unknown.

Results from the above studies should be interpreted with caution since concentrations of hormone used are either not reported or in some studies involve supraphysiological levels ranging from  $10\text{ }\mu\text{M}$  to  $30\text{ }\mu\text{M}$  of estradiol and  $0.3$  to  $30\text{ }\mu\text{M}$  of progesterone. While effects of these steroid concentrations may well be nonspecific, the hormone levels to which the cell is actually exposed may exceed reported blood levels (169). Clearly further studies are needed on genomic and nongenomic effects of estradiol or progesterone on  $\text{Ca}^{++}$  homeostasis in vascular smooth muscle and endothelial cells.

#### IV. Summary, Conclusions, and Directions for Future Research

Gender differences have long been described in the occurrence of several vascular disorders. Female hormones exert protective effects against some disease (e.g. coronary artery disease), and restoration of hormone levels improves symptoms in some disorders such as menstrual migraines and cardiac syndrome X. In other circumstances, alterations in female hormones may be linked to the onset of symptoms.

The protective effects of female sex steroids may be related to their vasodilatory properties. Studies suggest that estradiol enhances vasodilation by stimulating endothelial NO and prostacyclin activity and by attenuating vascular smooth muscle cell responses to vasoconstrictors such as ET-1. Estradiol may also have an inhibitory effect on voltage-gated  $\text{Ca}^{++}$  channels.

The influence of female hormones on the onset of symptoms or disease remains speculative and may include alterations in hormone levels and/or ratios. In certain vascular

beds, however, estrogen and progesterone may also stimulate vasoconstrictor pathways by increasing  $\alpha_1$ -adrenergic activity and/or catecholamine release.

The cellular mechanisms underlying these hormonal effects remain to be determined. Studies primarily conducted in nonvascular cell types support the likelihood that estrogen and progesterone can influence  $\text{Ca}^{++}$  homeostasis through genomic as well as nongenomic mechanisms of action. Research is needed to better define the effects of estrogen and progesterone in different vascular beds, the cellular mechanisms through which these effects are mediated, whether receptor-mediated induction of enzymes and proteins and/or immediate effects at the cell membrane are involved, and how the normal mechanisms are altered under conditions of vascular disease.

In addition, studies are needed to determine the clinical relevance of many of these observations. For instance, does chronic estrogen replacement therapy enhance endothelial cell vasodilator/vasoconstrictor function? The effects of progesterone also require study since, thus far, the primary focus has been on the effects of estrogen alone. The cardiovascular effects of long-term use of progesterone-only contraceptives, such as Norplant, remain to be determined. Finally, studies on the effects of both hormones combined are important as rarely, if ever, is the level of only one altered *in vivo*.

## References

- Rosselli M, Imthurn B, Macas E, Keller PJ, Dubey RK 1994 Circulating nitrite/nitrate levels increase with follicular development: indirect evidence for estradiol mediated NO release. *Biochem Biophys Res Commun* 202:1543–1552
- Polderman KH, Steouwer CDA, van Kamp GP, Dekker GA, Verheugt FWA, Gooren LJG 1993 Influence of sex hormones on plasma endothelin levels. *Ann Intern Med* 118:429–432
- Goldstein DS, Levinson P, Keiser HR 1983 Plasma and urinary catecholamines during the human ovulatory cycle. *Am J Obstet Gynecol* 146:824–829
- Becker RC 1992 Coronary artery disease in women: risks and prevention. *Coronary Acute Care* 3:2–10
- Bush TL 1990 The epidemiology of cardiovascular disease in postmenopausal women. *Ann NY Acad Sci* 592:263–271
- Anastos K, Charney P, Charon RA, Cohen E, Jones CY, Marte C, Swiderski DM, Wheat ME, Williams S 1991 Hypertension in women: what is really known? The Women's Caucus, Working Group on Women's Health of the Society of General Internal Medicine. *Ann Intern Med* 115:287–293
- Sokolow M, Perloff D 1961 The prognosis of essential hypertension treated conservatively. *Circulation* 23:697–713
- Messerli FH, Garavaglia GE, Schmieder RE, Sundgaard-Riise K, Nunez BD, Amodeo C 1987 Disparate cardiovascular findings in men and women with essential hypertension. *Ann Intern Med* 107:158–161
- Henderson BE, Ross RK, Paganini-Hill A, Mack TM 1986 Estrogen use and cardiovascular disease. *Am J Obstet Gynecol* 154:1181–1186
- Barrett-Connor E, Wingard DL, Criqui MH 1989 Postmenopausal estrogen use and heart disease risk factors in the 1980s. *Rancho Bernardo, Calif, revisited. JAMA* 261:2095–2100
- Ross RK, Paganini-Hill A, Mack TM, Arthur M, Henderson BE 1981 Menopausal oestrogen therapy and protection from death from ischaemic heart disease. *Lancet* 1:858–860
- Stampfer MJ, Colditz GA, Willett WC, Manson JE, Rosner B, Speizer FE, Hennekens CH 1991 Postmenopausal estrogen therapy and cardiovascular disease. Ten-year follow-up from the nurses' health study. *N Engl J Med* 325:756–762
- Gruchow HW, Anderson AJ, Barboriak JJ, Sobocinski KA 1988 Postmenopausal use of estrogen and occlusion of coronary arteries. *Am Heart J* 115:954–963
- Walsh BW, Schiff I, Rosner B, Greenberg L, Ravnkar V, Sacks FM 1991 Effects of postmenopausal estrogen replacement on the concentrations and metabolism of plasma lipoproteins. *N Engl J Med* 325:1196–1204
- Bush TL, Barrett-Connor E, Cowan LD, Criqui MH, Wallace RB, Suchindran CM, Tyroler HA, Rifkind BM 1987 Cardiovascular mortality and noncontraceptive use of estrogen in women: results from the Lipid Research Clinics Program Follow-up Study. *Circulation* 75:1102–1109
- Sarrel PM, Lindsay D, Rosano GM, Poole-Wilson PA 1992 Angina and normal coronary arteries in women: gynecologic findings. *Am J Obstet Gynecol* 167:467–471
- Sax FL, Cannon RO, Hanson C, Epstein SE 1987 Impaired forearm vasodilator reserve in patients with microvascular angina. Evidence of a generalized disorder of vascular function? *N Engl J Med* 317:1366–1370
- Gant NF, Daley GL, Chand S, Whalley PJ, MacDonald PC 1973 A study of angiotensin II pressor response throughout primigravid pregnancy. *J Clin Invest* 52:2682–2689
- Phippard AF, Horvath JS, Glynn EM, Garner MG, Fletcher PJ, Duggin GG, Tiller DJ 1986 Circulatory adaptation to pregnancy—serial studies of haemodynamics, blood volume, renin and aldosterone in the baboon (*Papio hamadryas*). *J Hypertens* 4:773–779
- Naden RP, Gant Jr NF, Rosenfeld CR 1984 The pressor response to angiotensin II: the roles of peripheral and cardiac responses in pregnant and nonpregnant sheep. *Am J Obstet Gynecol* 148:450–457
- Harrison GL, Moore LG 1989 Blunted vasoreactivity in pregnant guinea pigs is not restored by meclofenamate. *Am J Obstet Gynecol* 160:258–264
- Conrad KP, Colpoys MC 1986 Evidence against the hypothesis that prostaglandins are the vasodepressor agents of pregnancy. Serial studies in chronically instrumented, conscious rats. *J Clin Invest* 77:236–245
- Curran-Everett D, Morris Jr KG, Moore LG 1991 Regional circulatory contributions to increased systemic vascular conductance of pregnancy. *Am J Physiol* 261:H1842–H1847
- Wang Y, Walsh SW, Cuo J, Zhang J 1991 Maternal levels of prostacyclin, thromboxane, vitamin E and lipid peroxides throughout normal pregnancy. *Am J Obstet Gynecol* 165:1690–1694
- Conrad KP, Joffe GM, Kruszyna H, Kruszyna R, Rochelle LG, Smith RP, Chavez JE, Mosher MD 1993 Identification of increased nitric oxide biosynthesis during pregnancy in rats. *FASEB J* 7:566–571
- Ramsay B, De Belder A, Campbell S, Moncada S, Martins JF 1994 A nitric oxide donor improves uterine artery diastolic blood flow in normal early pregnancy and in women at high risk of pre-eclampsia. *Eur J Clin Invest* 24:76–78
- Mastrogianis DS, O'Brien WF, Krammer J, Benoit R 1991 Potential role of endothelin-1 in normal and hypertensive pregnancies. *Am J Obstet Gynecol* 165:1711–1716
- Furuhashi N, Tsujie M, Kimura H, Yajima A, Nagae H, Tsunoda K, Abe K 1992 Plasma endothelin concentrations in normal pregnancy and pre-eclampsia. *Int J Feto-Matern Med* 5:1–6
- Silberstein SD, Merriam GR 1993 Sex hormones and headache. *J Pain Symptom Manage* 8:98–114
- Coffman JD 1991 Raynaud's phenomenon. An update. *Hypertension* 17:593–602
- Tancredi RG 1992 Pulmonary vascular disease: primary pulmonary hypertension. *Cardiovasc Clin* 22:113–124
- Nattero G 1982 Menstrual headache. *Adv Neurol* 33:215–226
- Lance JW 1973 *The Mechanisms and Management of Headache*, ed 2. Butterworth, London
- Greene R, Dalton K 1953 The premenstrual syndrome. *Br Med J* 1:1007–1014
- Robertson D, Oates JA 1978 Variant angina and Raynaud's phenomenon [letter]. *Lancet* 1:452
- Heupler FA 1980 Syndrome of symptomatic arterial spasm with nearly normal coronary arteriograms. *Am J Cardiol* 45:873–881

37. Rotton WN, Sachtleben MR, Friedman EA 1959 Migraine and eclampsia. *Obstet Gynecol* 14:322-330
38. Miller D, Waters DD, Warnica W, Szlachcic J, Kreeft J, Theroux P 1981 Is variant angina the coronary manifestation of a generalized vasospastic disorder? *N Engl J Med* 304:763-766
39. Celoria GC, Friedell GH, Sommers SC 1960 Raynaud's disease and primary pulmonary hypertension. *Circulation* 22:1055-1059
40. Taft EB, Mallory GK 1946 Primary pulmonary vascular sclerosis. *Arch Pathol* 42:630-634
41. Wade G, Ball J 1957 Unexplained pulmonary hypertension. *Q J Med* 26:83
42. Smith WM, Kroop IG 1957 Raynaud's disease in primary pulmonary hypertension. *JAMA* 165:1245-1248
43. Schumacher Jr EE 1957 Cor pulmonale in association with Raynaud's disease. Report of a case with sympathectomy. *Henry Ford Hosp Med J* 5:256-258
44. Fuster V, Steele PM, Edwards WD, Gersh BJ, McGoon MD, Frye RL 1984 Primary pulmonary hypertension: natural history and the importance of thrombosis. *Circulation* 70:580-587
45. Christman BW, McPherson CD, Newman JH, King GA, Bernard GR, Groves BM, Loyd JE 1992 An imbalance between the excretion of thromboxane and prostacyclin metabolites in pulmonary hypertension. *N Engl J Med* 327:70-75
46. Fitzgerald DJ, Rocki W, Murray R, Mayo G, Fitzgerald GA 1990 Thromboxane A2 synthesis in pregnancy-induced hypertension. *Lancet* 335:751-754
47. Reilly IA, Roy L, Fitzgerald GA 1986 Biosynthesis of thromboxane in patients with systemic sclerosis and Raynaud's phenomenon. *Br Med J (Clin Res Ed)* 292:1037-1039
48. Hughes EC 1972 *Obstetrics-Gynecology Terminology*. FA Davis, Philadelphia, pp 422-423
49. Walsh SW 1985 Preeclampsia: an imbalance in placental prostacyclin and thromboxane production. *Am J Obstet Gynecol* 152:335-340
50. Fitzgerald DJ, Entman SS, Mulloy K, Fitzgerald GA 1987 Decreased prostacyclin biosynthesis preceding the clinical manifestation of pregnancy-induced hypertension. *Circulation* 75:956-963
51. Seligman SP, Buyon JP, Clancy RM, Young BK, Abramson SB 1994 The role of nitric oxide in the pathogenesis of preeclampsia. *Am J Obstet Gynecol* 171:944-948
52. Baylis C, Engels K 1992 Adverse interactions between pregnancy and a new model of systemic hypertension produced by chronic blockade of endothelial derived relaxing factor (EDRF) in the rat. *Clin Exp Hypertens [B]* 11:117-129
53. Giaid A, Yanagisawa M, Langleben D, Michel RP, Levy R, Shennib H, Kimura S, Masaki T, Duguid WP, Stewart DJ 1993 Expression of endothelin-1 in the lungs of patients with pulmonary hypertension. *N Engl J Med* 328:1732-1739
54. Giaid A, Saleh D 1995 Reduced expression of endothelial nitric oxide synthase in the lungs of patients with pulmonary hypertension. *N Engl J Med* 333:214-221
55. Smith CD, McKendry RJ 1982 Controlled trial of nifedipine in the treatment of Raynaud's phenomenon. *Lancet* 2:1299-1301
56. Rich S, Kaufman E, Levy PS 1992 The effects of high doses of calcium-channel blockers on survival in primary pulmonary hypertension. *N Engl J Med* 327:76-81
57. Belizan JM, Villar J, Gonzalez L, Campodonico L, Bergel E 1991 Calcium supplementation to prevent hypertensive disorders of pregnancy. *N Engl J Med* 325:1399-1405
58. Mochizuki M, Morikawa H, Yamasaki M, Maruo T 1991 Vascular reactivity in normal and abnormal gestation. *Am J Kidney Dis* 17:139-143
59. Kilby MD, Pipkin FB, Symonds EM 1992 Calcium and platelets in normotensive and hypertensive human pregnancy. *J Hypertens* 10:997-1003
60. Ebeigbe AB, Ezimokhai M, Aloamaka CP 1987 Responses of arterial smooth muscle from normotensive and pre-eclamptic subjects to the calcium channel agonist, Bay K 8644. *Res Exp Med (Berl)* 187:461-468
61. Gant NF, Jimenez JM, Whalley PJ, Chand S, MacDonald PC 1977 A prospective study of angiotensin II pressor responsiveness in pregnancies complicated by chronic essential hypertension. *Am J Obstet Gynecol* 127:369-375
62. Nisell H, Hjemdahl P, Linde B 1985 Cardiovascular responses to circulating catecholamines in normal pregnancy and in pregnancy-induced hypertension. *Clin Physiol* 5:479-493
63. Raab W, Schroeder G, Wagner R, Gigue W 1956 Vascular reactivity and electrolytes in normal and toxemic pregnancies. *J Clin Endocrinol* 16:1196-1216
64. Zamudio S, Palmer SK, Dahms TE, Berman JC, Moore LG 1995 Alterations in uteroplacental blood flow precede the onset of hypertension in preeclampsia at high altitude. *J Appl Physiol*, in press
65. Oian P, Kjeldsen SE, Eide I, Maltau JM 1986 Increased arterial catecholamines in pre-eclampsia. *Acta Obstet Gynecol Scand* 65:613-617
66. Nisell H, Hjemdahl P, Linde B, Lunell NO 1985 Sympathoadrenal and cardiovascular reactivity in pregnancy-induced hypertension. II. Responses to tilting. *Am J Obstet Gynecol* 152:554-560
67. Tulchinsky D, Okada DM 1975 Hormones in human pregnancy. IV. Plasma progesterone. *Am J Obstet Gynecol* 121:293-299
68. Dawood MY, Ratnam SS 1974 Serial estimations of serum unconjugated estradiol-17 beta in high-risk pregnancies. *Obstet Gynecol* 44:200-207
69. Zamudio S, Leslie K, White M, Hagerman D, Moore LG 1994 Low serum estradiol and high serum progesterone concentrations characterize hypertensive pregnancies at high altitude. *J Soc Gynecol Invest* 1:197-205
70. Horth CE, Wainscott G, Neylan C, Wilkinson MI 1975 Proceedings: progesterone, oestradiol and aldosterone levels in plasma during the menstrual cycle of women suffering from migraine. *J Endocrinol* 65:24P-25P
71. Epstein MT, Hockaday JM, Hockaday TD 1975 Migraine and reproductive hormones throughout the menstrual cycle. *Lancet* 1:543-548
72. Somerville BW 1975 Estrogen-withdrawal migraine. I. Duration of exposure required and attempted prophylaxis by premenstrual estrogen administration. *Neurology* 25:239-244
73. Somerville BW 1972 The role of estradiol withdrawal in the etiology of menstrual migraine. *Neurology* 22:355-365
74. de Lignieres B, Vincens M, Mauvais-Jarvis P, Mas JL, Touboul PJ, Bousser MG 1986 Prevention of menstrual migraine by percutaneous oestradiol. *Br Med J (Clin Res Ed)* 293:1540
75. Dennerstein L, Morse C, Burrows G, Oats J, Brown J, Smith M 1988 Menstrual migraine: a double-blind trial of percutaneous estradiol. *Gynecol Endocrinol* 2:113-120
76. Terregino CA, Seibold JR 1985 Influence of the menstrual cycle on Raynaud's phenomenon and on cold tolerance in normal women. *Angiology* 36:88-95
77. Farhat MY, Ramwell PW 1992 Estradiol potentiates the vasopressor response of the isolated perfused rat lung to the thromboxane mimic U-46619. *J Pharmacol Exp Ther* 261:686-691
78. Dawkins KD, Burke CM, Billingham ME, Jamieson SW 1986 Primary pulmonary hypertension and pregnancy. *Chest* 89:383-388
79. McCaffrey RM, Dunn LJ 1964 Primary pulmonary hypertension in pregnancy. *Obstet Gynecol Surv* 19:567-591
80. Miller VM, Vanhoutte PM 1991 Progesterone and modulation of endothelium-dependent responses in canine coronary arteries. *Am J Physiol* 261:R1022-R1027
81. Vu Hai MT, Warembourg M, Milgrom E 1977 Hormonal control of progesterone receptors. *Ann NY Acad Sci* 286:199-209
82. Horwitz KB, McGuire WL 1978 Estrogen control of progesterone receptor in human breast cancer. Correlation with nuclear processing of estrogen receptor. *J Biol Chem* 253:2223-2228
83. Milgrom E, Thi L, Atger M, Baulieu EE 1973 Mechanisms regulating the concentration and the conformation of progesterone receptor(s) in the uterus. *J Biol Chem* 248:6366-6374
84. May FE, Johnson MD, Wiseman LR, Wakeling AE, Kastner P, Westley BR 1989 Regulation of progesterone receptor mRNA by oestradiol and antioestrogens in breast cancer cell lines. *J Steroid Biochem* 33:1035-1041
85. Read LD, Snider CE, Miller JS, Greene GL, Katzenellenbogen BS 1988 Ligand-modulated regulation of progesterone receptor messenger ribonucleic acid and protein in human breast cancer cell lines. *Mol Endocrinol* 2:263-271
86. Kreitmann B, Bugat R, Bayard F 1979 Estrogen and progesterin

- regulation of the progesterone receptor concentration in human endometrium. *J Clin Endocrinol Metab* 49:926–929
87. Oka T, Schimke RT 1969 Interaction of estrogen and progesterone in chick oviduct development. II. Effects of estrogen and progesterone on tubular gland cell function. *J Cell Biol* 43:123–137
  88. Means AR, O'Malley BW 1971 Protein biosynthesis on chick oviduct polyribosomes. II. Regulation by progesterone. *Biochemistry* 10:1570–1576
  89. Bronson FH, Hamilton TH 1972 A comparison of nucleic acid synthesis in the mouse oviduct and uterus: interactions between estradiol and progesterone. *Biol Reprod* 6:160–167
  90. Leung BS, Sasaki GH 1973 Prolactin and progesterone effect on specific estradiol binding in uterine and mammary tissues *in vitro*. *Biochem Biophys Res Commun* 55:1180–1187
  91. Brenner RM, Resko JA, West NB 1974 Cyclic changes in oviductal morphology and residual cytoplasmic estradiol binding capacity induced by sequential estradiol-progesterone treatment of spayed Rhesus monkeys. *Endocrinology* 95:1094–1104
  92. Hsueh AJ, Peck Jr EJ, Clark JH 1976 Control of uterine estrogen receptor levels by progesterone. *Endocrinology* 98:438–444
  93. Cato ACB, Beato M 1985 The hormonal regulation of uteroglobin gene expression. *Anticancer Res* 5:65–72
  94. Dotzlaw H, Miller T, Varvelas J, Murphy LC 1990 Epidermal growth factor gene expression in human breast cancer biopsy samples: relationship to estrogen and progesterone receptor gene expression. *Cancer Res* 50:4204–4108
  95. Pfaff DW, McEwen BS 1983 Actions of estrogens and progestins on nerve cells. *Science* 219:808–814
  96. Horwitz KB, Horwitz LD 1982 Canine vascular tissues are targets for androgens, estrogens, progestins and glucocorticoids. *J Clin Invest* 69:750–758
  97. Lieberman JR, vanVroohoven CJ, Beckmann I, Van der Kwast TH, Wallenburg HCS 1990 Uterine artery estrogen receptors in the nonpregnant and pregnant guinea pig. *Am J Obstet Gynecol* 163:1685–1688
  98. Perrot-Applanat M, Picard-Groyer MT, Lorenzo GF, Milgrom E 1988 Immunocytochemical demonstration of estrogen and progesterone receptors in muscle cells of uterine arteries in rabbits and humans. *Endocrinology* 123:1511–1519
  99. Orimo A, Inoue S, Ikegami A, Hosoi T, Akishita M, Ouchi Y, Maramatsu M, Orimo H 1993 Vascular smooth muscle cells as target for estrogen. *Biochem Biophys Res Commun* 195:730–736
  100. Karas RH, Patterson BL, Mendelsohn MD 1994 Human vascular smooth muscle cells contain function estrogen receptor. *Circulation* 89:1943–1950
  101. Sadovsky Y, Riemer RK, Roberts JM 1992 The concentration of estrogen receptors in rabbit uterine myocytes decreases in culture. *Am J Obstet Gynecol* 167:1631–1635
  102. Perrot-Applanat M, Deng M, Fernandez H, Lelaidier C, Meduri G, Bouchard P 1994 Immunohistochemical localization of estradiol and progesterone receptors in human uterus throughout pregnancy: expression in endometrial blood vessels. *J Clin Endocrinol Metab* 78:216–224
  103. Losordo DW, Kearney M, Kim BA, Jekanowski J, Isner JM 1994 Variable expression of the estrogen receptor in normal and atherosclerotic coronary arteries in premenopausal women. *Circulation* 89:1501–1510
  104. Williams JK, Adams MR, Klopfenstein HS 1990 Estrogen modulates responses of atherosclerotic coronary arteries. *Circulation* 81:1680–1687
  105. Gilligan DM, Badar DM, Panza JA, Quyyumi AA, Cannon RO 1994 Acute vascular effects of estrogen in postmenopausal women. *Circulation* 90:786–791
  106. Reis SE, Gloth ST, Blumenthal RS, Resar JR, Zacur HA, Gerstenblith G, Brinker JA 1994 Ethinyl estradiol acutely attenuates abnormal coronary vasomotor responses to acetylcholine in postmenopausal women. *Circulation* 89:52–60
  107. Gilligan DM, Quyyumi AA, Cannon RO 1994 Effects of physiological levels of estrogen on coronary vasomotor function in postmenopausal women. *Circulation* 89:2545–2551
  108. Jiang CW, Sarrel PM, Lindsay DC, Poole-Wilson PA, Collins P 1991 Endothelium-independent relaxation of rabbit coronary artery by 17 beta-oestradiol *in vitro*. *Br J Pharmacol* 104:1033–1037
  109. Jiang CW, Sarrel PM, Lindsay DC, Poole-Wilson PA, Collins P 1992 Progesterone induces endothelium-independent relaxation of rabbit coronary artery *in vitro*. *Eur J Pharmacol* 211:163–167
  110. Rosenfeld CR, Jackson GM 1984 Estrogen-induced refractoriness to the pressor effects of infused angiotensin II. *Am J Obstet Gynecol* 148:429–435
  111. Killam AP, Rosenfeld CR, Battaglia FC, Makowski EL, Meschia G 1973 Effect of estrogens on the uterine blood flow of oophorectomized ewes. *Am J Obstet Gynecol* 115:1045–1052
  112. Magness RR, Rosenfeld CR 1989 Local and systemic estradiol-17 beta: effects on uterine and systemic vasodilation. *Am J Physiol* 256:E536–E542
  113. Magness RR, Parker Jr CR, Rosenfeld CR 1993 Systemic and uterine responses to chronic infusion of estradiol-17 beta. *Am J Physiol* 265:E690–E698
  114. Tamai T, Matsuura S, Tatsumi N, Nunotani T, Sagawa N 1984 Role of sex steroid hormones in relative refractoriness to angiotensin II during pregnancy. *Am J Obstet Gynecol* 149:177–183
  115. Nakamura T, Matsui K, Ito M, Yoshimura T, Kawasaki N, Fujisaki S, Okamura H 1988 Effects of pregnancy and hormone treatments on pressor response to angiotensin II in conscious rats. *Am J Obstet Gynecol* 159:989–995
  116. Conrad KP, Mosher MD, Brinck-Johnsen T, Colpoys MC 1994 Effects of 17 beta-estradiol and progesterone on pressor responses in conscious ovariectomized rats. *Am J Physiol* 266:R1267–R1272
  117. Novak K, Kaufman S 1991 Effects of pregnancy, estradiol, and progesterone on pressor responsiveness to angiotensin II. *Am J Physiol* 261:R1164–R1170
  118. Dogterom J, De Jong W 1974 Diminished pressor response to noradrenaline of the perfused tail artery of pregnant rats. *Eur J Pharmacol* 25:267–269
  119. Farley DB, Ford SP 1992 Evidence for declining extracellular calcium uptake and protein kinase C activity in uterine arterial smooth muscle during gestation in gilts. *Biol Reprod* 46:315–321
  120. Stice SL, Ford SP, Rosazza JP, Van Orden DE 1987 Interaction of 4-hydroxylated estradiol and potential-sensitive Ca<sup>2+</sup> channels in altering uterine blood flow during the estrous cycle and early pregnancy in gilts. *Biol Reprod* 36:369–375
  121. Stice SL, Ford SP, Rosazza JP, Van Orden DE 1987 Role of 4-hydroxylated estradiol in reducing Ca<sup>2+</sup> uptake of uterine arterial smooth muscle cells through potential-sensitive channels. *Biol Reprod* 36:361–368
  122. Ferrara N, Houck K, Jakeman L, Leung DW 1992 Molecular and biological properties of the vascular endothelial growth factor family of proteins. *Endocr Rev* 13:18–32
  123. Flavahan NA, Shimokawa H, Vanhoutte PM 1989 Pertussis toxin inhibits endothelium-dependent relaxations to certain agonists in porcine coronary arteries. *J Physiol* 408:549–560
  124. Flavahan NA, Shimokawa H, Vanhoutte PM 1991 Inhibition of endothelium-dependent relaxations by phorbol myristate acetate in canine arteries: role of a pertussis toxin-sensitive G-protein. *J Pharmacol Exp Ther* 256:50–55
  125. Culotta E, Koshland Jr DE 1992 NO news is good news. *Science* 258:1862–1865
  126. Hayashi T, Fukuto JM, Ignarro LJ, Chaudhuri G 1992 Basal release of nitric oxide from aortic rings is greater in female rabbits than in male rabbits: implications for atherosclerosis. *Proc Natl Acad Sci USA* 89:11259–11263
  127. Bell C 1973 Oestrogen-induced sensitization of the uterine artery of the guinea pig to acetylcholine. *Br J Pharmacol* 49:595–601
  128. Weiner C, Liu KZ, Thompson L, Herrig J, Chestnut D 1991 Effect of pregnancy on endothelium and smooth muscle: their role in reduced adrenergic sensitivity. *Am J Physiol* 261:H1275–H1283
  129. Goetz RM, Morano I, Calovini T, Studer R, Holtz J 1994 Increased expression of endothelial constitutive nitric oxide synthase in rat aorta during pregnancy. *Biochem Biophys Res Commun* 205:905–910
  130. Van Buren GA, Yang DS, Clark KE 1992 Estrogen-induced uterine vasodilatation is antagonized by L- nitroarginine methyl ester, an inhibitor of nitric oxide synthesis. *Am J Obstet Gynecol* 167:828–833
  131. Williams SP, Shackelford DP, Iams SG, Mustafa SJ 1988 Endo-

- thelium-dependent relaxation in estrogen-treated spontaneously hypertensive rats. *Eur J Pharmacol* 145:205–207
132. **Gisclard V, Miller VM, Vanhoutte PM** 1988 Effect of 17 beta-estradiol on endothelium-dependent responses in the rabbit. *J Pharmacol Exp Ther* 244:19–22
  133. **Collins P, Shay J, Jiang C, Moss J** 1994 Nitric oxide accounts for dose-dependent estrogen-mediated coronary relaxation after acute estrogen withdrawal. *Circulation* 90:1964–1968
  134. **Jiang C, Sarrel PM, Poole-Wilson PA, Collins P** 1992 Acute effects of 17 B-estradiol on rabbit coronary artery contractile responses to endothelin-1. *Am J Physiol* 263:H271–H275
  135. **Miller VM, Vanhoutte PM** 1990 17b-Estradiol augments endothelium-dependent contractions to arachidonic acid in rabbit aorta. *Am J Physiol* 258:R1502–R1507
  136. **Weiner CP, Lizasoain I, Baylis SA, Knowles RG, Charles RG, Moncada S** 1994 Induction of calcium-dependent nitric oxide synthases by sex hormones. *Proc Natl Acad Sci USA* 91:5212–5216
  137. **Hishikawa K, Nakaki T, Marumo T, Suzuki H, Kato R, Saruta T** 1995 Up-regulation of nitric oxide synthase by estradiol in human aortic endothelial cells. *FEBS Lett* 360:291–293
  138. **Miyahara K, Kawamoto T, Sase K, Yui Y, Toda K, Yang LX, Hattori R, Aoyama T, Yamamoto Y, Doi Y** 1994 Cloning and structural characterization of the human endothelial nitric-oxide-synthase gene. *Eur J Biochem* 223:719–726
  139. **Yanagisawa M, Kurihara H, Kimura S, Tomobe Y, Kobayashi M, Mitsui Y, Yazaki Y, Goto K, Masaki T** 1988 A novel potent vasoconstrictor peptide produced by vascular endothelial cells. *Nature* 332:411–415
  140. **Emori T, Hirata Y, Ohta K, Shichiri M, Marumo F** 1989 Secretory mechanism of immunoreactive endothelin in cultured bovine endothelial cells. *Biochem Biophys Res Commun* 160:93–100
  141. **Schini VB, Hendrickson H, Heublein DM, Burnett Jr JC, Vanhoutte PM** 1989 Thrombin enhances the release of endothelin from cultured porcine aortic endothelial cells. *Eur J Pharmacol* 165:333–334
  142. **Boulanger C, Luscher TF** 1990 Release of endothelin from the porcine aorta. Inhibition by endothelium-derived nitric oxide. *J Clin Invest* 85:587–590
  143. **Muck AO, Seeger H, Korte K, Lippert TH** 1993 The effect of 17 beta-estradiol and endothelin 1 on prostacyclin and thromboxane production in human endothelial cell cultures. *Clin Exp Obstet Gynecol* 20:203–206
  144. **Davies MG, Hagen PO** 1993 The vascular endothelium. A new horizon. *Ann Surg* 218:593–609
  145. **Wakasugi M, Noguchi T, Kazama YI, Kanemaru Y, Onaya T** 1989 The effects of sex hormones on the synthesis of prostacyclin (PGI<sub>2</sub>) by vascular tissues. *Prostaglandins* 37:401–410
  146. **Chang WC, Nakao J, Orimo H, Murota SI** 1980 Stimulation of prostaglandin cyclooxygenase and prostacyclin synthetase activities by estradiol in rat aortic smooth muscle cells. *Biochim Biophys Acta* 620:472–482
  147. **Crandall ME, Keve TM, McLaughlin MK** 1990 Characterization of norepinephrine sensitivity in the maternal splanchnic circulation during pregnancy. *Am J Obstet Gynecol* 162:1296–1301
  148. **Fallgren B, Bjellin L, Edvinsson L** 1988 Effect of pregnancy and sex steroids on alpha 1-adrenoceptor mechanisms in the guinea-pig uterine vascular bed. *Pharmacol Toxicol* 63:375–381
  149. **Gisclard V, Flavahan NA, Vanhoutte PM** 1987 Alpha adrenergic responses of blood vessels of rabbits after ovariectomy and administration of 17 beta-estradiol. *J Pharmacol Exp Ther* 240:466–470
  150. **Colucci WS, Gimbrone Jr MA, McLaughlin MK, Halpern W, Alexander RW** 1982 Increased vascular catecholamine sensitivity and alpha-adrenergic receptor affinity in female and estrogen-treated male rats. *Circ Res* 50:805–811
  151. **Miller VM, Vanhoutte PM** 1985 Endothelial  $\alpha_2$ -adrenoceptors in canine pulmonary and systemic blood vessels. *Eur J Pharmacol* 118:123–129
  152. **Angus JA, Cocks TM, Satoh K** 1986 The alpha adrenoceptors on endothelial cells. *Fed Proc* 45:2355–2359
  153. **Ford SP, Reynolds LP, Farley DB, Bhatnagar RK, Van Orden DE** 1984 Interaction of ovarian steroids and periarterial alpha 1-adrenergic receptors in altering uterine blood flow during the estrous cycle of gilts. *Am J Obstet Gynecol* 150:480–484
  154. **Hamlet MA, Rorie DK, Tyce GM** 1980 Effects of estradiol on release and disposition of norepinephrine from nerve endings. *Am J Physiol* 239:H450–H456
  155. **Ahmad S, Chretien P, Daniel EE, Shen SH** 1990 Characterization of beta adrenoceptors on cultured endothelial cells by radioligand binding. *Life Sci* 47:2365–2370
  156. **Roberts JM, Riemer RK, Bottari SP, Wu YY, Goldfien A** 1989 Hormonal regulation myometrial adrenergic responses: the receptor and beyond. *J Dev Physiol* 11:125–134
  157. **Klangkalya B, Chan A** 1988 The effects of ovarian hormones on beta-adrenergic and muscarinic receptors in rat heart. *Life Sci* 42:2307–2314
  158. **Adams DJ, Barakeh J, Laskey R, Van Breemen C** 1989 Ion channels and regulation of intracellular calcium in vascular endothelial cells. *FASEB J* 3:2389–2400
  159. **Khalil R, Lodge N, Saida K, Van Breemen C** 1987 Mechanism of calcium activation in vascular smooth muscle. *J Hypertens Suppl* 5:S5–S15
  160. **Bean BP** 1991 Pharmacology of calcium channels in cardiac muscle, vascular muscle, and neurons. *Am J Hypertens* 4:406S–411S
  161. **Bossu JL, Elhamdani A, Feltz A, Tanzi F, Aunis D** 1992 Voltage-gated Ca entry in isolated bovine capillary endothelial cells: evidence of a new type of Bay K8644-sensitive channel. *Pflügers Arch* 420:200–207
  162. **Rendt JM, Toro L, Stefani E, Erulkar SD** 1992 Progesterone increases Ca<sup>2+</sup> currents in myometrial cells from immature and nonpregnant adult rats. *Am J Physiol* 262:C293–C301
  163. **Batra S** 1990 Influence of chronic oestrogen treatment on the density of muscarinic cholinergic receptors and calcium channels in the rabbit uterus. *J Endocrinol* 125:185–189
  164. **Zhang F, Ram JL, Standley PR, Sowers JR** 1994 17 beta-estradiol attenuates voltage-dependent Ca<sup>2+</sup> currents in A7r5 vascular smooth muscle cell line. *Am J Physiol* 266:C975–980
  165. **Morishita S** 1986 Prompt effect of progesterone on the adrenergic response of smooth muscles. *Jpn J Pharmacol* 42:289–296
  166. **Jiang C, Poole-Wilson PA, Sarrel PM, Mochizuki S, Collins P, MacLeod KT** 1992 Effect of 17 beta-oestradiol on contraction, Ca<sup>2+</sup> current and intracellular free Ca<sup>2+</sup> in guinea-pig isolated cardiac myocytes. *Br J Pharmacol* 106:739–745
  167. **Blackmore PF, Lattanzio FA** 1991 Cell surface localization of a novel non-genomic progesterone receptor on the head of human sperm. *Biochem Biophys Res Commun* 181:331–336
  168. **Blackmore PF, Neulen J, Lattanzio F, Beebe SJ** 1991 Cell surface-binding sites for progesterone mediate calcium uptake in human sperm. *J Biol Chem* 266:18655–18659
  169. **Meizel S, Turner KO** 1991 Progesterone acts at the plasma membrane of human sperm. *Mol Cell Endocrinol* 11:R1–R5
  170. **Toro L, Stefani E, Erulkar S** 1990 Hormonal regulation of potassium currents in single myometrial cells. *Proc Natl Acad Sci USA* 87:2892–2895
  171. **Sperelakis N, Inoue Y, Ohya Y** 1992 Fast Na<sup>+</sup> channels and slow Ca<sup>2+</sup> current in smooth muscle from pregnant rat uterus. *Mol Cell Biochem* 114:79–89
  172. **Boyle MB, MacLusky NJ, Naftolin F, Kaczmarek LK** 1987 Hormonal regulation of K<sup>+</sup>-channel messenger RNA in rat myometrium during oestrus cycle and in pregnancy. *Nature* 330:373–375