

# Rod outer segments mediate mitochondrial DNA damage and apoptosis in human retinal pigment epithelium

Gui-Feng Jin, John S. Hurst and Bernard F. Godley

Department of Ophthalmology and Visual Sciences, University of Texas Medical Branch, Galveston, USA

## Abstract

**Purpose.** To investigate the interrelationships between DNA damage, mitochondrial activity, and apoptosis in retinal pigment epithelial cells (RPE) after exposure to rod outer segments (ROS).

**Methods.** After incubation of cultured human RPE with ROS, mitochondrial redox function was evaluated from MTT reduction. Mitochondrial (mt) and nuclear (n) DNA damage were determined by quantitative polymerase chain reactions (QPCR). Apoptotic RPE cells were detected by binding of annexin V to phosphatidyl serine (PS) using fluorescence microscopy. The expression of the pro-apoptotic proteins, p53 and p21<sup>waf-1</sup>, and DNA repair enzymes, apurinic/apyrimidinic endonuclease (APE<sup>ref-1</sup>) and DNA polymerase  $\beta$  ( $\beta$ -pol) were quantitatively determined by Western blotting analysis.

**Results.** Mitochondrial function decreased by  $20 \pm 5\%$  and annexin V immunofluorescent binding was enhanced after exposure of cells to physiological levels of ROS ( $3.8 \times 10^6 \text{ cm}^{-2}$ ) for 4 h. MtDNA was preferentially damaged after exposure to ROS with increased lesion frequencies of  $1.49 \pm 0.37$  and  $2.2 \pm 0.14$  per 10 kb base pairs (bp), respectively after 5 and 7 h contact, compared to untreated controls (zero class damage). APE<sup>ref-1</sup> expression increased more than 340% above controls after exposure to ROS for 7 and 24 h. The expression of  $\beta$ -pol in cultures increased 110% above controls after 24 h contact with the ROS. The expression of p53 and p21 in cells increased 100 and 38% above controls after 24 h exposure to the ROS.

**Conclusions.** Exposure of ROS to ROS induced mtDNA damage and dysfunction and activated nDNA repair pathways, which did not prevent apoptosis.

**Keywords:** DNA damage; mitochondrial dysfunction; oxidative stress; retinal pigment epithelium; rod outer segments

## Introduction

In developed societies age-related macular degeneration (ARMD) predominates, as the cause of irreversible profound vision loss in people aged over 60 years, and in the future will consume a major portion of finite health resources as the aging population expands. It is a multifactorial syndrome with genetic and environmental contributions. It is hypothesized that lifelong chronic cumulative oxidative stress constitutes a significant risk factor.<sup>1–5</sup> The pathobiological changes that culminate in the loss and/or dysfunction of macular RPE cells are believed to be central to the development and progression of this disease.

The RPE regulates nutrient and metabolite exchange to the neural retina including the photoreceptors and is essential to their survival. Consequently degeneration of the post-mitotic RPE cells will result in the death of neighboring photoreceptors. Each RPE cell contacts 8–10 photoreceptor cells, and each photoreceptor cell sheds approximately 20–30 of its outer segment discs resulting in the daily phagocytosis by each RPE cell of more than 2000 photoreceptor outer segment discs.<sup>6</sup> The retinal environment favors the generation of reactive oxygen intermediates (ROI) as photoreceptors and RPE cells are metabolically highly reactive. The neural retina, especially the photoreceptor outer segment membranes, are enriched in polyunsaturated fatty acids, such as docosahexaenoic, which are susceptible to attack and degradation by ROI. In addition, the retina is continuously exposed to sunlight, especially hazardous blue light, has high

Received: January 22, 2001

Accepted: June 15, 2001

**Correspondence:** Bernard F. Godley, MD, PhD, FACS, Department of Ophthalmology and Visual Sciences, University of Texas Medical Branch, Galveston, TX 77555-0787, USA. Tel: (409) 772-28107, Fax: (409) 772-28106, E-mail: bgodley@utmb.edu

levels of molecular photosensitizers, and is subjected to high oxygen tension. Thus phagocytosis of ROS could augment oxidative stress to the RPE cells.

During photoreceptor outer segment phagocytosis, the RPE generate ROI such as superoxide<sup>7</sup> and hydrogen peroxide ( $\text{H}_2\text{O}_2$ ).<sup>8</sup> It has been suggested that increased phagocytic and metabolic burden imposed upon the RPE in the accumulation of age-related lipofuscin, derived in part from oxidatively damaged lipids and proteins, are implicated in RPE and photoreceptor death.<sup>9</sup>

Previously we have shown that hydrogen peroxide preferentially damages mtDNA, disrupts mt metabolic function<sup>10</sup> and promotes apoptosis in RPE cells (submitted). To study the interaction of the RPE with photoreceptor outer segments, cultured human RPE cells were exposed ROS for varying times. The effects upon RPE mt and nDNA damage were reported by sensitive QPCR assays. As aging and ROI are known to damage DNA,<sup>11–13</sup> especially mtDNA<sup>14–16</sup> by forming adducts, and generating mutagenic abasic sites, it would also be important to study the effects of ROS on the expression of the DNA base excision repair enzymes (BER), apurinic/apyrimidinic endonuclease ( $\text{APE}^{\text{ref-1}}$ ) and DNA polymerase  $\beta$  ( $\beta$ -pol), as it has been reported that these enzymes are upregulated as an adaptive response to oxidative stress<sup>17,18</sup> and cooperate in DNA repair.<sup>19</sup> We wanted to study potential interactions between factors promoting DNA repair and apoptosis in RPE cells, to determine if BER upregulation followed exposure of the cells to ROS, and also whether this could influence their survival.  $\text{APE}^{\text{ref-1}}$  and  $\beta$ -pol expression was analyzed by Western blotting. Mitochondrial redox function was quantitated by 3-[4,5-dimethyliazol-2-yl]-2,5-diphenyl tetrazolium bromide (MTT) reduction. Apoptosis was qualitatively determined from phosphatidyl serine (PS) externalization as suggested by increased immunofluorescent binding of annexin V to PS, and the expression of the pro-apoptotic and/or cell cycle growth arrest nuclear transcription factors p53 and p21<sup>waf-1</sup> were analyzed by Western blotting.

## Materials and methods

### Reagents

Electrophoretic-grade reagents: sodium chloride, sodium deoxycholate, sodium dodecyl sulfate (SDS), ethylene glycol-bis ( $\beta$ -aminoethyl ether)-N,N,N',N'-tetraacetic acid (EGTA), NP-40, polyacrylamide, Tris, Tween-20, and non-fat dry milk were obtained from Biorad Laboratories (Hercules, CA). Nitrocellulose membranes were obtained from Micron Separations (Westborough, MA). Antibodies raised to p21<sup>waf-1</sup> (monoclonal), and p53 (polyclonal), were purchased from Santa Cruz Biotechnology (Santa Cruz, CA). Polyclonal antibodies raised to  $\beta$ -pol, and  $\text{APE}^{\text{ref-1}}$  were the gifts of Dr. S. H. Wilson, Sealy Center for Molecular Science, University of Texas Medical Branch, Galveston, TX. The electrochemiluminescence (ECL) Western Blotting kit

including horse-radish peroxidase-conjugated secondary antibodies were obtained from Amersham-Pharmacia Biotechnology (Piscataway, NJ). Goat anti-mouse and anti-rabbit antibodies were used respectively with primary monoclonal and polyclonal antibodies. 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl tetrazolium bromide (MTT) was purchased from Promega (Madison, WI). Diethylenediaminetetraacetic acid (EDTA) dimethyl sulfoxide (DMSO) and sucrose were purchased from Sigma-Aldrich Co. (St. Louis, MO). Bradford reagent was purchased from Biorad (Hercules, CA). ApoAlert® Kits were obtained from Clontech Laboratories, Inc., Palo Alto, CA.

### RPE cell cultures

SV40-transformed fetal male human RPE cells (RPE 28 SV4, Coriell Institute, Camden, NJ) were cultured as previously described.<sup>10</sup> The rationale for using this established and well-documented cell line was as follows. A disadvantage with primary cell cultures derived from different donors is that the cellular response to  $\text{H}_2\text{O}_2$  may lack consistency, which would not be a concern with a well-defined cell line, such as the one used in our studies. These transformed cells exhibit epithelioid morphology and retain physiological functions characteristic of primary human RPE cells, such as polarity and the ability to phagocytize rod outer segments. Additional studies also indicated that both primary and transformed cell lines respond similarly to oxidative stress.<sup>20–23</sup>

### Isolation of rod outer segments (ROS)

ROS were aseptically isolated from fresh bovine eyes obtained locally (Dereck and sons, Inc., Santa Fe, TX) using a discontinuous sucrose density gradient.<sup>24</sup> Ocular globes were sectioned and the neural retinae were removed into 43% sucrose dissolved in 10 mM TRIS-buffered saline (TBS, pH 7.0) containing 60 mM KCl, 30 mM NaCl, 2 mM  $\text{MgCl}_2$ , and 0.1 mM EDTA and thoroughly vortexed and centrifuged at 5000 g for 5 min. The supernatants were diluted with TBS and centrifuged at 27 000 g for 15 min. The supernatants were discarded and the precipitates were suspended in TBS containing 33% sucrose. The sucrose content of the suspensions were determined by refractometry and was adjusted to 32.5% with added 45% sucrose solution. The sucrose was overlaid with TBS to create a gradient and the samples were centrifuged at 16 500 g for 2 h. The interphases between the two layers were collected by syringe aspiration and the ROS were counted using a hemocytometer chamber and were stored at  $-80^\circ\text{C}$  till use.

### MTT assay

The MTT reduction assay was used as the index for mitochondrial redox function. Cells were seeded onto 96-well plates at a density of 12 000 cells per well. The medium was replaced 48 h later with one that was serum-free containing varying numbers of ROS ( $0.95\text{--}15 \times 10^6 \text{ cm}^{-2}$ ). Cells were

exposed to ROS for 4 h, then the media containing the ROS were removed and the cells were washed with phosphate buffered saline (PBS). The cells were then incubated with serum-free medium containing  $4.83 \mu\text{M}$  MTT for 4 h at  $37^\circ\text{C}$ . Then  $150 \mu\text{l}$  DMSO was added to each well and the plates were shaken for 10 min. The supernatants were removed into fresh multi-well plates and the optical densities of the solutions were read at  $590 \text{ nm}$ <sup>25</sup> using a scanning multi-well spectrophotometer (Amersham Pharmacia Biotech, Piscataway, NJ). Absorbance values were converted to MTT reduction as previously described.<sup>10</sup> Values were normalized with respect to the untreated control cultures (100% mitochondrial redox function). Determinations were performed in duplicate and experiments were repeated ( $n = 4$ ).

### DNA damage

QPCR was used to determine the average lesion frequency per DNA strand in specified sequences for both template strands.<sup>10,26,27</sup> Only intact DNA sequences were amplified and lesions including strand breaks, apurinic sites and base modifications would prevent amplification. To control for template copy number differences, and damage unrelated to experimental conditions, DNA sequences of approximately 200 kb base pairs were quantitatively amplified. Total RPE cellular DNA was isolated using a genomic tip 20G kit (Quiagen, Inc., Valencia, CA) according to the instructions. Total DNA concentrations were determined by ethidium bromide fluorescence using an A4-filter fluorimeter with an excitation band pass filter at  $360 \text{ nm}$  and emission cut-off filter at  $600 \text{ nm}$  (Optical Devices, Elmsford, NY) using  $\lambda/\text{Hind III}$  DNA as standard. Sample quality was assessed by QPCR of a 222 base pair mtDNA fragment and an 84 base pair fragment of the  $\beta$ -globin gene as previously described.<sup>10</sup> QPCR was performed in a GeneAmp PCR system 9600 with the GeneAmp XLPCR kit (Perkin-Elmer, San Jose, CA). Reaction mixtures contained  $15 \text{ ng}$  genomic DNA as template. The composition and sequences of the primers and the PCR conditions employed for the 16.2 kb mtDNA, 13.4 kb  $\beta$ -globin, 12.2 kb  $\beta$ -pol and the 10 kb hprt products were as previously described.<sup>10</sup> A quantitative control using  $7.5 \text{ ng}$  genomic template was included in every PCR series. PCR products were electrophoretically resolved on 1% agarose gels (dissolved in TBE buffer, Amresco, Solon, OH) at  $80\text{--}90 \text{ v}$  for 4 h. Dried gels were exposed to phosphor screens for 12 h and quantitated using ImageQuant software (Molecular Dynamics, Sunnyvale, CA). DNA lesion frequencies ( $\lambda$ ) were calculated as the amplification of damaged DNA relative ( $A_d$ ) to non-damaged controls ( $A_0$ ), where  $\lambda = -\ln A_d/A_0$ . Experiments to assess DNA damage in genomic and mtDNA were determined 4 times in duplicate.

### Phosphatidyl serine exposure

RPE cells were placed on a microscope slide and exposed to  $3.8 \times 10^6 \text{ cm}^{-2}$  ROS for 4 h. Cells were washed in PBS and

then the binding buffer supplied with the ApoAlert® kit.  $5 \mu\text{l}$  annexin V conjugated to fluoroisothiocyanate (FITC) and  $10 \mu\text{l}$  propidium iodide were added and the cells were covered with a glass coverslip and incubated with the dyes for 10–15 min in the dark. The cells were viewed under a fluorescence microscope, using a dual filter set for FITC and rhodamine. Apoptotic RPE cells stained with FITC-Annexin V fluoresced green, whereas necrotic cells stained with propidium iodide fluoresced red. The experiment was repeated ( $n = 2$ ).

### Western blot analysis

Cultured RPE cells ( $\sim 90\%$  confluent) exposed to  $3.8 \times 10^6 \text{ cm}^{-2}$  for different times were lysed by RIPA buffer ( $1 \text{ mM}$  Tris-HCl, sodium deoxycholate, NaCl, EGTA, 1% NP-40.<sup>28</sup> Cell lysates corresponding to  $50 \mu\text{g}$  protein, determined according to Bradford,<sup>29</sup> and pre-stained molecular weight markers were loaded onto 1% SDS, 10–15% polyacrylamide gels under reducing conditions. Following electrophoretic separation, the proteins were transferred to nitrocellulose membranes. After blocking with 5% non-fat dry milk and washing in  $10 \text{ mM}$  Tris-buffered saline, pH 7.5, containing, 0.1% Tween-20, (T-TBS) the membranes were incubated with the appropriately diluted antibodies to, APE<sup>ref-1</sup>,  $\beta$ -pol, p21<sup>waf-1</sup> or p53 for 3 h at room temperature. The primary antibodies were decanted and the membranes were washed with TBS. The membranes were then incubated for 2 h at room temperature with horseradish peroxidase conjugated secondary antibodies (goat anti-mouse for the monoclonal p53 and goat anti-rabbit for the polyclonal APE<sup>ref-1</sup>,  $\beta$ -pol, P53, and p21<sup>waf-1</sup> antibodies) in T-TBS diluted to the same concentrations as the primary antibodies. The membranes were washed with TBS and incubated with the ECL substrate for 1 min and then exposed to X-OMAT AR film (Eastman Kodak, Rochester, NY). The processed gel images were analyzed using a phosphorimager with the supplied ImageQuant software (Molecular Dynamics, Sunnyvale, CA). Analyses for each protein were repeated ( $n = 2$ ).

### Statistical analysis

Data for the QPCR and MTT are presented as the means  $\pm$  standard deviation (SD). The Students *t*-test was used for statistical evaluation between experimental groups. When the *p* values were  $<0.05$ , values were considered significant.

## Results

### Mitochondrial dysfunction

Mitochondrial redox function as assayed by the MTT test is generally considered an index for cellular survival. Mitochondrial MTT reduction in cells treated for 4 h with 3.8, 7.6, and  $15 \times 10^6 \text{ cm}^{-2}$  ROS was decreased by 20, 36, and 46%,




Figure 1. Effect of rod outer segments on mitochondrial redox function in RPE cells.

Histogram showing the effects of different concentrations of rod outer segments (ROS) on RPE mitochondrial redox function assayed by MTT reduction. RPE were exposed to the indicated concentrations of bovine ROS for 4 h at 37°C. After exposure, cells were rinsed with PBS and incubated with serum-free media containing 4.83  $\mu\text{M}$  MTT. Media were removed, the cells lysed, and absorbance determined at 570 nm. MTT reduction was determined as a percentage of untreated controls. Data are expressed as means  $\pm$  SD from 4 experiments. Each experiment was performed in duplicate.

respectively (Fig. 1). This indicated a loss of mt redox function or dysfunction as a possible consequence of mtDNA damage after exposure to ROS.

### Mitochondrial DNA damage

RPE cells were treated with  $3.8 \times 10^6 \text{ cm}^{-2}$  ROS to determine whether ROS exposure could compromise either mt or nDNA integrity in the RPE cells. This was assessed from lesion frequency in the relative amplification of mt and nDNA sequences. MtDNA damage significantly increased with increased exposure time to ROS. After 5 and 7 h treatment, mtDNA lesion frequencies increased from  $0 \pm 0.0015$  (control RPE cells not exposed to ROS) to  $1.49 \pm 0.39$  and  $2.2 \pm 0.14$  per 10 kb base pairs, respectively (Fig. 2). Whereas nDNA, represented by the three genomic gene clusters,  $\beta$ -globin,  $\beta$ -pol, and hprt was resistant to ROS-induced damage and did not suffer increased lesion frequency (Fig. 2). Moreover, lesion frequency in  $\beta$ -pol was less after 5 and 7 h exposure to ROS, suggesting an increased rate of repair of pre-existing lesions.

### PS exposure

RPE cells were treated with  $3.8 \times 10^6 \text{ cm}^{-2}$  ROS for 4 h to determine if PS externalization, an apoptotic marker, fol-




Figure 2. Time course of DNA damage in RPE cells after exposure to rod outer segments.

RPE cells were exposed to  $3.8 \times 10^6 \text{ cm}^{-2}$  ROS for indicated times. Lesions per 10 kb base pairs nuclear (n) and mitochondrial (mt) DNA were determined by quantitative polymerase chain reactions (QPCR), using the Poisson equation. Untreated controls were considered zero class damage. Data are expressed as the means  $\pm$  SD from 4 experiments. Each experiment was performed in duplicate.

nDNA is represented by genes coding for  $\beta$ -globin, hprt and DNA polymerase  $\beta$  ( $\beta$ -pol).


lowed. Annexin V binding to exposed PS was detected by fluorescence microscopy<sup>30</sup> as shown in Figures 3A and B. Low levels of Annexin V and propidium iodide fluorescence are shown in control cells not exposed to ROS (Fig. 3A). Whereas Annexin V labeling as shown by green fluorescence was noticeably increased in cells treated with ROS (Fig. 3B), suggesting an increase in apoptotic death after interaction with the ROS.

### Effect of ROS exposure on levels of BER genes

Oxidative stress is known to upregulate levels of BER enzymes, such as APE<sup>ref-1</sup>, and  $\beta$ -pol. Therefore experiments were conducted to determine if levels of these enzymes increased in RPE cells after exposure to ROS. Levels of APE<sup>ref-1</sup> were increased in RPE cells after 5, 7, and 24 h exposure times to  $3.8 \times 10^6 \text{ cm}^{-2}$  ROS (Figs. 4A and B). After 5 h, there was a greater than twofold increase above control values. After 7 h, expression was 344% above control values. A similar increase (360% above control values) was also observed in cells treated with ROS for 24 h. The Western blot is representative of 2 determinations from different cell cultures.

Levels of  $\beta$ -pol was increased twofold above control values in cells exposed to  $3.8 \times 10^6 \text{ cm}^{-2}$  ROS for 24 h (Figs. 5A and B). The Western blot is representative of two determinations from different cell cultures.






**Figure 3.** Phosphatidyl serine exposure in RPE cell plasma membranes after exposure to ROS.

Photomicrographs showing phosphatidyl serine (PS) externalization in cultured RPE cells exposed to  $3.8 \times 10^6 \text{ cm}^{-2}$  ROS for 4 h. Cells were immunofluorescently-labeled with propidium iodide, PI, (orange, necrotic cells) and annexin V (green, PS-labeled apoptotic cells). A Untreated control cells, with isolated necrotic and apoptotic cells are shown in (A). ROS-exposed cells with higher levels of apoptotic cells, as indicated by increased green immunofluorescent binding of Annexin V to exposed PS are shown in (B). Photomicrographs are representative of 2 experiments. Magnification was 200 $\times$ .

### P53 and p21<sup>waf-1</sup> expression

Increases in levels of p53 and p21<sup>waf-1</sup> are associated with apoptosis, therefore the influence of ROS exposure on the levels of these proteins in RPE cells was studied. The expression of p53 (Figs. 6A and B) and p21 (Figs. 7A and B) after exposure to  $3.8 \times 10^6 \text{ cm}^{-2}$  were followed in cultured RPE cells as a function of time. Expression of p53 and p21<sup>waf-1</sup> was only affected after 24 h of exposure, when it increased 100 and 38% respectively, above the control values, suggesting that extracellular contact of RPE cells with ROS increased levels of both proteins and that p53 did not apparently induce p21<sup>waf-1</sup> synthesis. The Western blots for both proteins were performed twice for each time-point and the




**Figure 4.** Time course of APE<sup>ref-1</sup> expression in RPE cells exposed to ROS.

RPE cells were exposed to  $3.8 \times 10^6 \text{ cm}^{-2}$  ROS for the indicated times. Protein expression was analyzed by Western blot. Each gel lane (A) was loaded with 50  $\mu\text{g}$  protein. Control lanes represent protein expression in untreated cells. The densitometric analysis of (A) is shown in (B) and represents the mean of 2 experiments.

data shown represents the averaged values obtained from two different experiments.

### Discussion

Oxidative stress is proposed as a major contributory factor in the degeneration of aging RPE cells.<sup>1,4</sup> It is also believed that an increased rate of apoptotic cell death in the RPE layer could mediate and hasten the onset of ARMD.<sup>31</sup> This hypothesized mechanism as an etiological factor for ARMD has recently been supported by the confirmed presence of apoptotic RPE cells in retinal specimens of ARMD patients.<sup>32</sup> Mitochondria are recognized as pivotal mediators of apoptotic signaling<sup>33</sup> and mtDNA is more susceptible to damage from ROI than nDNA because it lacks non-coding introns and protective histones, but especially because it is in close proximity to high levels of ROI generated in the mt membranes. The relatively low capacity to repair mtDNA has been correlated with apoptosis<sup>34</sup> and oxidant-induced damage to mtDNA is at least three times more extensive and more sus-




**Figure 5.** Time course of DNA polymerase  $\beta$  ( $\beta$ -pol) expression in RPE cells exposed to ROS.

RPE cells were exposed to  $3.8 \times 10^6 \text{ cm}^{-2}$  ROS for the indicated times. Protein expression was analyzed by Western blot. Each gel lane (A) was loaded with  $50 \mu\text{g}$  protein. Control lanes represent protein expression in untreated cells. The densitometric analysis of (A) is shown in (B) and represents the mean of 2 experiments.

tained than that to nDNA.<sup>35</sup> Our previous studies reported mitochondrial redox dysfunction with consequential mtDNA damage<sup>10</sup> and apoptosis in RPE cells in response to the prooxidant,  $\text{H}_2\text{O}_2$  (unpublished data).

The daily phagocytosis of spent photoreceptor outer segment discs, enriched in polyunsaturated fatty acids, by the RPE has been identified as a major source of ROI, such as superoxide<sup>7</sup> and  $\text{H}_2\text{O}_2$ .<sup>8</sup> The present studies indicate that this interaction of cultured RPE with ROS could also trigger mitochondrial metabolic dysfunction that correlates with mtDNA damage and also with apoptosis, as suggested by the QPCR assays and the enhanced immunofluorescent binding of annexin V to externalized PS. The increased expression of the pro-apoptotic and cell cycle arrest-growth proteins, p53 and p22<sup>waf-1</sup>, in the RPE cells further support the evidence for apoptosis after contact with ROS. However, despite some similarities in responses by the RPE cells at the molecular level to both  $\text{H}_2\text{O}_2$  and ROS, there were also significant qualitative and quantitative differences. For example, the



**Figure 6.** Time course of p53 expression in RPE cells exposed to ROS.

RPE cells were exposed  $3.8 \times 10^6 \text{ cm}^{-2}$  ROS for the indicated times. Protein expression was analyzed by Western blot. Each gel lane (A) was loaded with  $50 \mu\text{g}$  protein. Control lanes represent protein expression in untreated cells. The densitometric analysis of (A) is shown in (B) and represents the mean of 2 experiments.

response time courses in the RPE cells differed with  $\text{H}_2\text{O}_2$  or ROS treatments. Generally, a delay was observed in cells exposed to ROS when compared to the effects of  $\text{H}_2\text{O}_2$ . Mitochondrial redox activity in RPE cells exposed to ROS was inhibited by 20% after 4h, which was similar to that in cells exposed to  $50\text{--}100 \mu\text{M}$   $\text{H}_2\text{O}_2$  after only 1h.<sup>10</sup> The lesion frequency in the RPE mtDNA with the ROS treatments after 5 or 7h exposure was greater than that previously reported for  $200 \mu\text{M}$   $\text{H}_2\text{O}_2$  after 1h<sup>10</sup> and the compensatory response in the transcriptionally active gene  $\beta$ -pol was also increased.

Low levels of the ROI, hypochlorite and  $\text{H}_2\text{O}_2$ , have been previously reported to activate and induce APE<sup>ref-1</sup> in hamster ovarian and HeLa cells after 3 to 9h lag periods, suggesting involvement of multiple signaling intermediates.<sup>17,36</sup> Our data also showed that APE<sup>ref-1</sup> was also activated/induced in RPE cells after exposure to ROS for 7 and 24h. APE<sup>ref-1</sup> is a multifunctional protein that not only repairs DNA but also regulates the DNA-binding of redox-regulated transcription




Figure 7. Time course of p21<sup>waf-1</sup> expression in RPE cells exposed to ROS.

RPE cells were exposed to  $3.8 \times 10^6 \text{ cm}^{-2}$  ROS for the indicated times. Protein expression was analyzed by Western blot. Each gel lane (A) was loaded with 50  $\mu\text{g}$  protein. Control lanes represent protein expression in untreated cells. The densitometric analysis of (A) is shown in (B) and represents the mean of 2 experiments.

factors, such as nuclear transcription factor  $\kappa\beta$ , Fos, Jun, p21<sup>waf-1</sup> and Bax, etc.<sup>37</sup> and also modulates the activation of p53.<sup>38</sup> In addition,  $\beta$ -pol was also activated/induced after 24h ROS treatment. Previous studies such as those of Chen *et al.* have reported the upregulation of  $\beta$ -pol in murine monocytes and fibroblasts in response to oxidative-stress-inducing agents including H<sub>2</sub>O<sub>2</sub>.<sup>18</sup> Regarding this observation therefore, it may be significant that the reported specific interaction of APE<sup>ref-1</sup> and  $\beta$ -pol is coordinated to assemble the polymerase onto potentially mutagenic abasic nDNA sites to facilitate base excision repairs.<sup>19</sup> However, despite these late adaptive responses to oxidative stress in the cultured RPE cells after exposure to ROS, the putative early pro-apoptotic reactions such as cytochrome *c* and PS migrations already committed some susceptible cells to an activated apoptotic program from which they could not be rescued. Interestingly, Chen *et al.* recently showed that oxidized low density lipoprotein (LDL) down-regulates  $\beta$ -pol in murine monocytes.<sup>39</sup> Oxidized lipids and proteins including LDL are believed to be present in the ROS daily phagocytized by the RPE and constitute part of the drusen and other extracellu-

lar deposits that accumulate with age and may be implicated in ARMD progression. The upregulation of  $\beta$ -pol seen in the RPE cultures exposed to ROS may represent a short-term adaptation to oxidative stress and may not reflect the long-term chronic *in vivo* response which might be different and may result in  $\beta$ -pol down regulation with consequently accelerated cell death.

PS exposure, detected as enhanced fluorescent labeling, was evident in cultures treated for only 1 h with 50–200  $\mu\text{M}$  H<sub>2</sub>O<sub>2</sub>, whereas it was apparent only after 4 h exposure with ROS. This implies that the oxidative stress that followed ROS contact was delayed compared to direct H<sub>2</sub>O<sub>2</sub> exposure, and/or that H<sub>2</sub>O<sub>2</sub> release during ROS phagocytosis could only in part account for the resultant apoptosis.

Expression of p53, known to be an potent cell death effector after activation by mtDNA damage, was increased in RPE cultures above constitutively-expressed levels only after protracted exposure (24h) to ROS, unlike the rapid increase observed after 1 h H<sub>2</sub>O<sub>2</sub> treatments (unpublished data). It is possible that this represents the previously demonstrated stimulation of p53 activity by redox-dependent and independent activities of APE<sup>ref-1</sup> in HeLa cell.<sup>38</sup> Therefore, APE<sup>ref-1</sup> may function as a critical sensor of genotoxic oxidative stress in RPE and other cells and may play an important role in switching the cell cycle program from survival to apoptosis. In the ROS model of oxidative stress, increased p53 activity did not induce p21<sup>waf-1</sup> expression, as it is unlikely that the modest increases in p21<sup>waf-1</sup> levels (38% above control values) could be attributed to upregulation of its gene activity by p53 protein. This contrasted with the ninefold increase in p21<sup>waf-1</sup> protein expression observed in RPE cells, which correlated with raised p53 protein levels after treatment with 200  $\mu\text{M}$  H<sub>2</sub>O<sub>2</sub> (unpublished data). Evidently therefore, apoptosis induced by ROS in the RPE cells was largely a p21<sup>waf-1</sup>-independent process. Experimental evidence from studies with the P21<sup>waf-1</sup> null-mutant mouse demonstrate that p53-promotes apoptosis in the absence of p21<sup>waf-1</sup>.<sup>40,41</sup> support our hypothesis that the apoptosis of RPE cells induced by p53 is not mediated via the p21 pathway. The mechanisms by which p53 is believed to induce apoptosis independent of the p21 pathway is by stimulation of a burst of ROI on the mitochondrial membrane that releases the triggers of the effector pathway of apoptosis.<sup>42</sup>  $\beta$ -pol is reported to be upregulated by p21<sup>waf-1</sup>.<sup>43</sup> However, it is unlikely that the induction of  $\beta$ -pol was due to the modest increase in p21<sup>waf-1</sup> levels seen in our study.

The marked contrasts in the modulation of apoptotic activity by H<sub>2</sub>O<sub>2</sub> and ROS most probably reflect different ROI species. H<sub>2</sub>O<sub>2</sub> participates in ROS-induced apoptosis of RPE cells, but from our studies may not be the major promoter. It is very possible that ROS lipid hydroperoxides and their aldehydic end products, such as 4-hydroxynonenal (HNE) and 4-hydroxyhexenal, are more actively involved. Arachidonate-derived lipid hydroperoxides, such as the-15 isomer are potent inducers of apoptosis in cell lines derived from T lymphocytes<sup>44</sup> and human retinoblastoma cells<sup>45</sup> and

also prevent inhibition of apoptosis by Bcl-2 overexpression.<sup>44</sup> Mitochondrial dysfunction and apoptosis in PC12 and neuronal cells have been reported as consequences of 4-HNE exposure.<sup>46</sup>

In conclusion, our studies show that upregulation of nDNA repair enzymes in response to oxidative stress is correlated with the increased expression of nuclear transcription factors that promote apoptosis rather than arrested cell growth.

## Acknowledgments

This study was supported by the Paul Beeson Physician Aging Research Program, the Research to Prevent Blindness, Inc., and by Grant EY12850-01 from the National Institutes of Health.

## References

1. Bressler NM, Bressler SB, Fine SL. Age-related macular degeneration. *Surv Ophthalmol*. 1988;32:375–413.
2. Snodderly DM. Evidence for protection against age-related macular degeneration by carotenoids and antioxidant vitamins. *Am J Clin Nutr*. 1995;62:1448S–1461S.
3. van der Hagen AM, Yolton DP, Kaminski MS, Yolton RL. Free radicals and antioxidant supplementation: A review of their roles in age-related macular degeneration. *J Am Optom Assoc*. 1993;64:871–878.
4. Winkler BS, Boulton ME, Gottsch JD, Sternberg P. Oxidative damage and age-related macular degeneration. *Mol Vis*. 1999;5:32.
5. Rose RC, Richer SP, Bode AM. Ocular oxidants and antioxidant protection. *Proc Soc Exp Biol Med*. 1998;217:397–407.
6. Young RW. The renewal of photoreceptor cell outer segments. *J Cell Biol*. 1967;33:61–72.
7. Dorey CK, Khouri GG, Syniuta LA, Curran SA, Weiter JJ. Superoxide production by porcine retinal pigment epithelium *in vitro*. *Invest Ophthalmol Vis Sci*. 1989;30:1047–1054.
8. Tate DJ, Jr., Miceli MV, Newsome DA. Phagocytosis and H<sub>2</sub>O<sub>2</sub> induce catalase and metallothionein gene expression in human retinal pigment epithelial cells. *Invest Ophthalmol Vis Sci*. 1995;36:1271–1279.
9. Dorey CK, Wu G, Ebenstein D, Garsd A, Weiter JJ. Cell loss in the aging retina. Relationship to lipofuscin accumulation and macular degeneration. *Invest Ophthalmol Vis Sci*. 1989;30:1691–1699.
10. Ballinger SW, Van Houten B, Jin GF, Conklin CA, Godley BF. Hydrogen peroxide causes significant mitochondrial DNA damage in human RPE cells. *Exp Eye Res*. 1999;68:765–772.
11. Adelman R, Saul RL, Ames BN. Oxidative damage to DNA: Relation to species metabolic rate and life span. *Proc Natl Acad Sci USA*. 1988;85:2706–2708.
12. Ames BN, Saul RL. In: Iversen OH, ed. *Theories of Carcinogenesis*. Washington, DC, Hemisphere Publishing Corp; 1988:203–220.
13. Richter C, Park J-W, Ames BN. Normal oxidative damage to mitochondrial and nuclear DNA is extensive. *Proc Natl Acad Sci USA*. 1988;85:6465–6467.
14. Porn-Ares MI, Samali A, Orrenius S. Cleavage of the calpain inhibitor, calpastatin, during apoptosis. *Cell Death Differ*. 1998;5:1028–1033.
15. Wei YH. Oxidative stress and mitochondrial DNA mutations in human aging. *Proc Soc Exp Biol Med*. 1998;217:53–63.
16. Wei YH. Mitochondrial DNA mutations and oxidative damage in aging and diseases: An emerging paradigm of gerontology and medicine. *Proc Natl Sci Council, Rep China – Part B, Life Sciences*. 1998;22:55–67.
17. Ramana CV, Boldogh I, Izumi T, Mitra S. Activation of apurinic/apyrimidinic endonuclease in human cells by reactive oxygen species and its correlation with their adaptive response to genotoxicity of free radicals. *Proc Natl Acad Sci USA*. 1998;95:5061–5066.
18. Chen KH, Yakes FM, Srivastava DK, Singhal RK, Sobol RW, Horton JK, Van Houten B, Wilson SH. Up-regulation of base excision repair correlates with enhanced protection against a DNA damaging agent in mouse cell lines. *Nucleic Acids Res*. 1998;26:2001–2007.
19. Bennett RA, Wilson DM, III, Wong D, Demple B. Interaction of human apurinic endonuclease and DNA polymerase beta in the base excision repair pathway. *Proc Natl Acad Sci USA*. 1997;94:7166–7169.
20. Kerendian J, Enomoto H, Wong CG. Induction of stress proteins in SV-40 transformed human RPE-derived cells by organic oxidants. *Curr Eye Res*. 1992;11:385–396.
21. Kvant A. Expression and secretion of transforming growth factor-beta in transformed and nontransformed retinal pigment epithelial cells. *Ophthalmic Res*. 1994;26:361–367.
22. Sippy BD, Hofman FM, He S, Osusky R, Sheu SJ, Walker SM, Ryan SJ, Hinton DR. SV40-immortalized and primary cultured human retinal pigment epithelial cells share similar patterns of cytokine-receptor expression and cytokine responsiveness. *Curr Eye Res*. 1995;14:495–503.
23. Kumaki N, Anderson DM, Cosman D, Kumaki S. Expression of interleukin-15 and its receptor by human fetal retinal pigment epithelial cells. *Curr Eye Res*. 1996;15:876–882.
24. van Kuijk FJ, Buck P, Lewis JW, Kliger DS. Preparative scale isolation and partial purification of human rod outer segments. *Exp Eye Res*. 1993;57:249–252.
25. Hansen MB, Nielsen SE. Reexamination and further development of a precise and rapid dye method for measuring cell/growth kill. *J Immunol Methods*. 1989;119:203–210.
26. Kalinowski DP, Illenye S, Van Houten B. Analysis of DNA damage and repair in murine leukemia L1210 cells using a quantitative polymerase chain reaction assay. *Nucleic Acids Res*. 1992;20:3485–3494.



27. Van Houten B, Chen Y, Nicklas JA, Rainville IR, O'Neill JP. Development of long PCR techniques to analyze deletion mutations of the human hprt gene. *Mutat Res.* 1998;403:171–175.
28. Rao GN. Hydrogen peroxide induces complex formation of SHC-Grb2-SOS with receptor tyrosine kinase and activates Ras and extracellular signal-regulated protein kinases group of mitogen-activated protein kinases. *Oncogene.* 1996;13:713–719.
29. Bradford M. A rapid and sensitive method for the quantitation of protein utilizing the principle of protein dye-binding. *Anal Biochem.* 1976;72:248–254.
30. Zhang G, Gurtu V, Kain SR, Yan G. Early detection of apoptosis using a fluorescent conjugate of annexin V. *Biotechniques.* 1997;23:525–531.
31. Cai J, Yang J, Jones DP. Mitochondrial control of apoptosis: The role of cytochrome *c*. *Biochim Biophys Acta.* 1998;1366:139–149.
32. Hinton DR, He S, Lopez PF. Apoptosis in surgically excised choroidal neovascular membranes in age-related macular degeneration. *Arch Ophthalmol.* 1998;116:203–209.
33. Susin SA, Zamzami N, Kroemer G. Mitochondria as regulators of apoptosis: Doubt no more. *Biochim Biophys Acta.* 1998;1366:151–165.
34. LeDoux SP, Wilson GL, Beecham EJ, Stevnsner T, Wassermann K, Bohr VA. Repair of mitochondrial DNA after various types of DNA damage in Chinese hamster ovary cells. *Carcinogenesis.* 1992;13:1967–1973.
35. Yakes FM, Van Houten B. Mitochondrial DNA damage is more extensive and persists longer than nuclear DNA damage in human cells following oxidative stress. *Proc Natl Acad Sci USA.* 1997;94:514–519.
36. Grosch S, Fritz G, Kaina B. Apurinic endonuclease (Ref-1) is induced in mammalian cells by oxidative stress and involved in clastogenic adaptation. *Cancer Res.* 1998;58:4410–4416.
37. Xanthoudakis S, Miao GG, Curran T. The redox and DNA-repair activities of Ref-1 are encoded by nonoverlapping domains. *Proc Natl Acad Sci USA.* 1994;91:23–27.
38. Jayaraman L, Murthy KG, Zhu C, Curran T, Xanthoudakis S, Prives C. Identification of redox/repair protein Ref-1 as a potent activator of p53. *Genes Dev.* 1997;11:558–570.
39. Chen KH, Srivastava DK, Singhal RK, Jacob S, Ahmed AE, Wilson SH. Modulation of base excision repair by low density lipoprotein, oxidized low density lipoprotein and antioxidants in mouse monocytes. *Carcinogenesis.* 2000;21:1017–1022.
40. Deng C, Zhang P, Harper JW, Elledge SJ, Leder P. Mice lacking p21CIP1/WAF1 undergo normal development, but are defective in G1 checkpoint control. *Cell.* 1995;82:675–684.
41. Attardi LD, Lowe SW, Brugarolas J, Jacks T. Transcriptional activation by p53, but not induction of the p21 gene, is essential for oncogene-mediated apoptosis. *EMBO J.* 1996;15:3693–3701.
42. Polyak K, Xia Y, Zweier JL, Kinzler KW, Vogelstein B. A model for p53-induced apoptosis. *Nature.* 1997;389:300–305.
43. Kedar PS, Lowy DR, Widen SG, Wilson SH. Transfected human beta-polymerase promoter contains a ras-responsive element. *Mol Cell Biol.* 1990;10:3852–3856.
44. Sandstrom PA, Pardi D, Tebbey PW, Dudek RW, Terrian DM, Folks TM, Buttke TM. Lipid hydroperoxide-induced apoptosis: Lack of inhibition by Bcl-2 over-expression. *FEBS Lett.* 1995;365:66–70.
45. Vento R, d'Allesandro N, Giuliano M, Lauricella A, Carabillo M, Tesoriere G. Induction of apoptosis by arachidonic acid in human retinoblastoma Y79 cells: Involvement of oxidative stress. *Exp Eye Res.* 2000;70:503–517.
46. Kruman I, Bruce-Keller AJ, Bredesen D, Waeg G, Mattson MP. Evidence that 4-hydroxynonenal mediates oxidative stress-induced neuronal apoptosis. *J Neurosci.* 1997;17:5089–5100.