

# Protein Sequence and Structure

**Proteomes Interactomes and Biological Networks**

5, 7 and 8 November, 2019

Emidio Capriotti

<http://biofold.org/>




**Biomolecules  
Folding and  
Disease**

Department of Pharmacy and  
Biotechnology (FaBiT)  
University of Bologna



# The Central Dogma




from <http://www.scq.ubc.ca>

<https://www.youtube.com/watch?v=9kOGOY7vthk>

# Amino Acid

The side chain (R) determines the type of the amino acid




# Physico-chemical Properties

The properties of the amino acid depends on the side chain


| Amino acid    | Abbrev. | Side chain  | Hydro-phobic | Polar | Charged  | Small | Tiny | Aromatic or Aliphatic | van der Waals volume | Codon                        | Occurrence in proteins (%) |
|---------------|---------|---|--------------|-------|----------|-------|------|-----------------------|----------------------|------------------------------|----------------------------|
| Alanine       | Ala, A  | -CH <sub>3</sub>  | X            | -     | -        | X     | X    | -                     | 67                   | GCU, GCC, GCA, GCG           | 7.8                        |
| Cysteine      | Cys, C  | -CH <sub>2</sub> SH   | X            | -     | -        | X     | -    | -                     | 86                   | UGU, UGC                     | 1.9                        |
| Aspartate     | Asp, D  | -CH <sub>2</sub> COOH   | -            | X     | negative | X     | -    | -                     | 91                   | GAU, GAC                     | 5.3                        |
| Glutamate     | Glu, E  | -CH <sub>2</sub> CH <sub>2</sub> COOH                         | -            | X     | negative | -     | -    | -                     | 109                  | GAA, GAG                     | 6.3                        |
| Phenylalanine | Phe, F  | -CH <sub>2</sub> C <sub>6</sub> H <sub>5</sub>                | X            | -     | -        | -     | -    | Aromatic              | 136                  | UUU, UUC                     | 3.9                        |
| Glycine       | Gly, G  | -H  | X            | -     | -        | X     | X    | -                     | 48                   | GGU, GGC, GGA, GGG           | 7.2                        |
| Histidine     | His, H  | -CH <sub>2</sub> C <sub>3</sub> H <sub>3</sub> N <sub>2</sub> | -            | X     | positive | -     | -    | Aromatic              | 118                  | CAU, CAC                     | 2.3                        |
| Isoleucine    | Ile, I  | -CH(CH <sub>3</sub> )CH <sub>2</sub> CH <sub>3</sub>          | X            | -     | -        | -     | -    | Aliphatic             | 124                  | AUU, AUC, AUA                | 5.3                        |
| Lysine        | Lys, K  | -(CH <sub>2</sub> ) <sub>4</sub> NH <sub>2</sub>              | -            | X     | positive | -     | -    | -                     | 135                  | AAA, AAG                     | 5.9                        |
| Leucine       | Leu, L  | -CH <sub>2</sub> CH(CH <sub>3</sub> ) <sub>2</sub>            | X            | -     | -        | -     | -    | Aliphatic             | 124                  | UUA, UUG, CUU, CUC, CUA, CUG | 9.1                        |
| Methionine    | Met, M  | -CH <sub>2</sub> CH <sub>2</sub> SCH <sub>3</sub>             | X            | -     | -        | -     | -    | -                     | 124                  | AUG                          | 2.3                        |
| Asparagine    | Asn, N  | -CH <sub>2</sub> CONH <sub>2</sub>                            | -            | X     | -        | X     | -    | -                     | 96                   | AAU, AAC                     | 4.3                        |
| Proline       | Pro, P  | -CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> -            | X            | -     | -        | X     | -    | -                     | 90                   | CCU, CCC, CCA, CCG           | 5.2                        |
| Glutamine     | Gln, Q  | -CH <sub>2</sub> CH <sub>2</sub> CONH <sub>2</sub>            | -            | X     | -        | -     | -    | -                     | 114                  | CAA, CAG                     | 4.2                        |
| Arginine      | Arg, R  | -(CH <sub>2</sub> ) <sub>3</sub> NH-C(NH)NH <sub>2</sub>      | -            | X     | positive | -     | -    | -                     | 148                  | CGU, CGC, CGA, CGG, AGA, AGG | 5.1                        |
| Serine        | Ser, S  | -CH <sub>2</sub> OH   | -            | X     | -        | X     | X    | -                     | 73                   | UCU, UCC, UCA, UCG, AGU, AGC | 6.8                        |
| Threonine     | Thr, T  | -CH(OH)CH <sub>3</sub>  | X            | X     | -        | X     | -    | -                     | 93                   | ACU, ACC, ACA, ACG           | 5.9                        |
| Valine        | Val, V  | -CH(CH <sub>3</sub> ) <sub>2</sub>                            | X            | -     | -        | X     | -    | Aliphatic             | 105                  | GUU, GUC, GUA, GUG           | 6.6                        |
| Tryptophan    | Trp, W  | -CH <sub>2</sub> C <sub>6</sub> H <sub>5</sub> N              | X            | -     | -        | -     | -    | Aromatic              | 163                  | UGG                          | 1.4                        |
| Tyrosine      | Tyr, Y  | -CH <sub>2</sub> C <sub>6</sub> H <sub>4</sub> OH             | X            | X     | -        | -     | -    | Aromatic              | 141                  | UAU, UAC                     | 3.2                        |

# Peptide Bond



# Torsion Angles


Backbone torsion angles determine the structure of the protein



# Protein folding

Protein folding is the process by which a protein assumes its native structure from the unfolded structure


T T C C P S I V A R S N F N V C R L P G T P E A L C A T  
Y T G C I I I P G A T C P G D Y A N



# The Anfinsen's hypothesis


The sequence contains all the information to specify 3-D structure

Anfinsen showed that denatured ribonuclease A could be re-activated removing the denaturant.



# Levinthal's paradox

A protein chain composed by 100 residues with 2 possible conformations has  $2^{100}$  ( $\sim 10^{30}$ ) possible conformations. Considering a time-step of  $10^{-12}$  s for visiting each conformation, the folding process would take  $10^{18}$  s, that is longer than the age of our Universe ( $2\text{-}3 \times 10^{17}$ s)



# The Anfinsen's Dogma

**Uniqueness:** requires that the sequence does **not have any other configuration with a comparable free energy.**

**Stability:** **small changes** in the surrounding environment **not affect the structure of the stable conformation.** This can be pictured as a free energy surface that looks more like a funnel and the free energy surface around the native state must be rather steep and high, in order to provide stability.

**Kinetic accessibility:** means that the path in the **free energy surface** from the unfolded to the folded state **must be reasonably smooth** or, in other words, that the folding of the chain must not involve highly complex changes in the shape.



# Aspects of the same problem

The solution of the protein folding consists in the understanding of three different aspects of the problem:

- Estimate the **stability of the native conformation** and thermodynamic of the process.
- Define the mechanism and the **kinetic of the process**.
- Predict the native **three-dimensional structure** of the protein.





# Folding and stability

The folding free energy difference,  $\Delta G_F$ , is typically small, of the order of -5 to -15 kcal/mol for a globular protein (compared to e.g. -30 to -100 kcal/mol for a covalent bond).




# Folding interactions

Several **electrostatic interactions** are contributing to the **stability** of the native state but they are **not the driving forces** in the folding process

| Type                      | Examples           | Binding energy (kcal/mol)  | Change of free energy water to ethanol (kcal/mol) |
|---------------------------|--------------------|--|---|
| Electrostatic interaction | Salt bridge        | $\text{—COO}^- \cdots \text{N}^+ \text{H}_3^+$                                       | -5  |
|                           | Dipole-dipole      |    | +0.3  |
| Hydrogen bond             | Water              |  | -4  |
|                           | Protein backbone   |  | -3  |
| Dispersion forces         | Aliphatic hydrogen |  | -0.03   |
| Hydrophobic forces        | Side chain of Phe  |  | -2.4  |


# Hydrophobic effect

- Water molecules form a cage-like structure around the nonpolar molecule.
- The positive  $\Delta H$  is due to the fact that the cage has to be broken to transfer the nonpolar molecule.
- The positive  $\Delta S$  is due to the fact that the water molecules are less ordered (an increase in the degree of disorder) when the cage is broken.




# Folding kinetics

The protein folding mechanism depends on the form of the free energy profile.  
Higher activation barrier corresponds to longer folding time




# Hierarchical organization of protein structure

Protein structure is defined by four levels of hierarchical organization.




# Secondary structure (I)

- Helices observed in proteins are mostly right-handed.
- Typical  $\phi$ ,  $\psi$  values for residues in  $\alpha$ -helix are around  $-60^\circ$ ;  $-50^\circ$
- Side chains project backward and outward.
- The core of  $\alpha$ -helix is tightly packed.



# Secondary structure (II)



- Typical  $\phi$ ,  $\psi$  values for residues in  $\beta$ -sheet are around  $140^\circ$ ,  $-130^\circ$
- Side chains of neighboring residues project in opposite directions.
- The polypeptide is in a more extended conformation.
- Parallel  $\beta$ -sheets are less stable than anti-parallel  $\beta$ -sheets.



# More complex structures

The arrangements of secondary structural elements form the Tertiary Structure of the protein.

The complex of **two or more protein domains defines the Quaternary Structure**. In the example Four-helix-bundle, EF-hand and SH2 domains together form an integrated phosphoprotein that functions as a negative regulator of many signaling pathways from receptors at the cell surface.



# Protein at the NCBI

The Protein database is a collection of sequences from several resources accessible through Entrez

<https://www.ncbi.nlm.nih.gov/protein/>

# Protein search

Using the name of the protein and the organism we can retrieve a specific protein

## P53 [Homo sapiens]

GenBank: BAC16799.1

[Identical Proteins](#) [FASTA](#) [Graphics](#)

---

[Go to:](#)

|            |   |        |        |                 |
|------------|---|--------|--------|-----------------|
| LOCUS      | BAC16799  | 393 aa | linear | PRI 01-APR-2003 |
| DEFINITION | P53 [Homo sapiens].   |        |        |                 |
| ACCESSION  | BAC16799  |        |        |                 |
| VERSION    | BAC16799.1  |        |        |                 |
| DBSOURCE   | accession <a href="#">AB082923.1</a>  |        |        |                 |
| KEYWORDS   | .   |        |        |                 |
| SOURCE     | Homo sapiens (human)  |        |        |                 |
| ORGANISM   | <u>Homo sapiens</u><br>Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;<br>Mammalia; Eutheria; Euarchontoglires; Primates; Haplorrhini;<br>Catarrhini; Hominidae; Homo.                                    |        |        |                 |
| REFERENCE  | 1   |        |        |                 |
| AUTHORS    | Azuma,K., Shichijo,S., Maeda,Y., Nakatsura,T., Nonaka,Y., Fujii,T.,<br>Koike,K. and Itoh,K.   |        |        |                 |
| TITLE      | Mutated p53 gene encodes a nonmutated epitope recognized by<br>HLA-B*4601-restricted and tumor cell-reactive CTLs at tumor site   |        |        |                 |
| JOURNAL    | Cancer Res. 63 (4), 854-858 (2003)  |        |        |                 |
| PUBMED     | <a href="#">12591737</a>  |        |        |                 |
| REFERENCE  | 2 (residues 1 to 393)   |        |        |                 |
| AUTHORS    | Shichijo,S. and Itoh,K.   |        |        |                 |
| TITLE      | Direct Submission   |        |        |                 |
| JOURNAL    | Submitted (26-MAR-2002) Shigeki Shichijo, Kurume Univ. School of<br>Med., Dep. Immunol.; 67-Asahi-machi, Kurume, Fukuoka 830-0011,<br>Japan (E-mail:shichijo@med.kurume-u.ac.jp, Tel:81-942-31-7551,<br>Fax:81-942-31-7699) |        |        |                 |
| FEATURES   | Location/Qualifiers   |        |        |                 |

# Protein Sequence DB

The main database of protein sequences is UniProt which is composed by SwissProt and TrEMBL

The mission of UniProt is to provide the scientific community with a comprehensive, high-quality and freely accessible resource of protein sequence and functional information.

**UniProtKB**  
UniProt Knowledgebase  
Swiss-Prot (561,176)  
Manually annotated and reviewed.  
Records with information extracted from literature and curator-evaluated computational analysis.

**UniRef**  
The UniProt Reference Clusters (UniRef) provide clustered sets of sequences from the UniProt Knowledgebase (including isoforms) and selected UniParc records.

**UniParc**  
UniParc is a comprehensive and non-redundant database that contains most of the publicly available protein sequences in the world.

**Proteomes**  
A proteome is the set of proteins thought to be expressed by an organism. UniProt provides proteomes for species with completely sequenced genomes.

**Supporting data**

- Literature citations
- Cross-ref. databases
- Taxonomy
- Diseases
- Subcellular locations
- Keywords

## Getting started

### Text search

Our basic text search allows you to search all the resources available



## UniProt data

### Download latest release

Get the UniProt data

### Statistics

View Swiss-Prot and TrEMBL statistics

### BLAST

Find regions of similarity between your sequences

<https://www.uniprot.org>

# UniProt Composition

Database of annotated proteins

- Swiss-Prot: Manually annotated ~560K

UniProtKB

UniProt Knowledgebase

Swiss-Prot (561,176)


 Manually annotated and reviewed.  
Records with information extracted from literature and curator-evaluated computational analysis.

TrEMBL (180,179,667)

 Automatically annotated and not reviewed.  
Records that await full manual annotation.

# The SwissProt

SwissProt contains all the **proteins that have been manually annotated** using information extracted from literature.



<http://www.expasy.org/>

# The function

Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type.  
11 publications

UniProt

UniProtKB

Advanced Search

BLAST Align Retrieve/ID mapping Help Contact

P04637 - P53\_HUMAN

Basket

Protein: Cellular tumor antigen p53  
Gene: TP53  
Organism: Homo sapiens (Human)  
Status: Reviewed - Annotation score: 00000 - Experimental evidence at protein level<sup>1</sup>

Display: None

Function: Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type. Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression. In cooperation with mitochondrial PP1F is involved in activating oxidative stress-induced necrosis; the function is largely independent of transcription. Induces the transcription of long intergenic non-coding RNA p21 (lncRNA-p21) and lncRNA-Mkln1. LncRNA-p21 participates in TP53-dependent transcriptional repression leading to apoptosis and seem to have effect on cell-cycle regulation. Implicated in Notch signaling cross-over. Prevents CDK7 kinase activity when associated to CAK complex in response to DNA damage, thus stopping cell cycle progression. Isoform 2 enhances the transactivation activity of isoform 1 from some but not all TP53-inducible promoters. Isoform 4 suppresses transactivation activity and impairs growth suppression mediated by isoform 1. Isoform 7 inhibits isoform 1-mediated apoptosis. [11 Publications](#)

Cofactor: Zn<sup>2+</sup>  
Note: Binds 1 zinc ion per subunit.

Sites

| Feature key                | Position(s) | Length | Description            | Graphical view | Feature identifier | Actions |
|----------------------------|-------------|--------|------------------------|----------------|--------------------|---------|
| Site <sup>1</sup>          | 120 – 120   |        | 1 Interaction with DNA |                |                    |         |
| Metal binding <sup>1</sup> | 176 – 176   |        | 1 Zinc                 |                |                    |         |
| Metal binding <sup>1</sup> | 179 – 179   |        | 1 Zinc                 |                |                    |         |
| Metal binding <sup>1</sup> | 238 – 238   |        | 1 Zinc                 |                |                    |         |
| Metal binding <sup>1</sup> | 242 – 242   |        | 1 Zinc                 |                |                    |         |

Regions

# Getting the information

The SwissProt **fasta file contains all the sequences** in the database and the **text file contains** all the information including **annotation**.

The fasta and text files can be downloaded using the following links

[http://www.uniprot.org/uniprot/P53\\_HUMAN.fasta](http://www.uniprot.org/uniprot/P53_HUMAN.fasta)  
[http://www.uniprot.org/uniprot/P53\\_HUMAN.txt](http://www.uniprot.org/uniprot/P53_HUMAN.txt)

More complex queries:

[http://www.uniprot.org/help/programmatic\\_access](http://www.uniprot.org/help/programmatic_access)

|    |  |                   |
|----|--|-------------------|
| ID | P53_HUMAN  | Reviewed; 393 AA. |
| AC | P04637; Q15086; Q15087; Q15088; Q16535; Q16807; Q16808; Q16809;        |                   |
| AC | Q16810; Q16811; Q16848; Q2XN98; Q3LRW1; Q3LRW2; Q3LRW3; Q3LRW4;        |                   |
| AC | Q3LRW5; Q86UG1; Q8J016; Q99659; Q9BTM4; Q9HAQ8; Q9NP68; Q9NPJ2;        |                   |
| AC | Q9NZD0; Q9UBI2; Q9UQ61;  |                   |
| DT | 13-AUG-1987, integrated into UniProtKB/Swiss-Prot.                     |                   |
| DT | 24-NOV-2009, sequence version 4.                                       |                   |
| DT | 04-FEB-2015, entry version 228.  |                   |
| DE | RecName: Full=Cellular tumor antigen p53;                              |                   |
| DE | AltName: Full=Antigen NY-CO-13;  |                   |
| DE | AltName: Full=Phosphoprotein p53;                                      |                   |
| DE | AltName: Full=Tumor suppressor p53;                                    |                   |
| GN | Name=TP53; Synonyms=P53;   |                   |
| OS | Homo sapiens (Human).  |                   |
| OC | Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;      |                   |
| OC | Mammalia; Eutheria; Euarchontoglires; Primates; Haplorrhini;           |                   |
| OC | Catarrhini; Hominidae; Homo.   |                   |
| OX | NCBI_TaxID=9606;   |                   |
| RN | [1]  |                   |
| RP | NUCLEOTIDE SEQUENCE [MRNA] (ISOFORM 1).                                |                   |
| RX | PubMed=4006916;  |                   |
| RA | Zakut-Houri R., Bienz-Tadmor B., Givol D., Oren M.;                    |                   |
| RT | "Human p53 cellular tumor antigen: cDNA sequence and expression in COS |                   |
| RT | cells.";   |                   |
| RL | EMBO J. 4:1251-1255(1985).   |                   |

# Exercise

From the UniProt FTP web site (<ftp://ftp.expasy.org/databases/uniprot/>) download the Human protein UP000005640\_9606 in fasta format.

- What is the total number of human proteins in the SwissProt and TrEMBL dataset?
- Given the fasta file containing the protein sequence of P53 what is the total number of residues?

# The Protein Data Bank

The largest repository of macromolecular structures obtained mainly by X-ray crystallography and NMR


The screenshot shows the RCSB PDB homepage. At the top, a dark blue header bar contains the text "RCSB PDB" and several navigation links: Deposit, Search, Visualize, Analyze, Download, Learn, and More. To the right of these is a yellow "MyPDB" button. Below the header is the RCSB PDB logo and a banner stating "157296 Biological Macromolecular Structures Enabling Breakthroughs in Research and Education". A search bar with the placeholder "Search by PDB ID, author, macromolecule, sequence, or ligands" and a "Go" button is positioned next to the banner. Below the search bar are links for "Advanced Search" and "Browse by Annotations". The main content area features a large map of the world with various molecular structures overlaid. On the left, a sidebar menu lists "Welcome", "Deposit", "Search", "Visualize", "Analyze", "Download", and "Learn". The central content area includes a section titled "A Structural View of Biology" with text about the archive's purpose and its role in biomedicine and agriculture. It also features a "Job Opportunities for Biocurators and Developers" section with an illustration of a computer monitor displaying a molecular model and a red chair. To the right, a section titled "November Molecule of the Month" displays a 3D molecular model of Phospholipase A2, composed of numerous blue spheres representing atoms.

<http://rcsb.org>


<http://ftp.rcsb.org/pub/pdb/>

# The Bovine Ribonuclease A




Ribonuclease A (RNase A) is a pancreatic ribonuclease that cleaves single-stranded RNA.



# Bonds and interactions



Examples of salt bridge, disulfide bond and hydrogen bonds in ribonuclease A



# PDB File

The most important information are the atomic coordinates.

|      |      | AT | RES | CH | POS | X       | Y      | Z       |      |       |
|------|------|----|-----|----|-----|---------|--------|---------|------|-------|
| ATOM | 2145 | N  | GLU | B  | 10  | 150.341 | 72.309 | 103.145 | 1.00 | 99.90 |
| ATOM | 2146 | CA | GLU | B  | 10  | 150.096 | 71.519 | 101.907 | 1.00 | 99.90 |
| ATOM | 2147 | C  | GLU | B  | 10  | 150.425 | 70.046 | 102.190 | 1.00 | 99.90 |
| ATOM | 2148 | O  | GLU | B  | 10  | 151.326 | 69.770 | 102.983 | 1.00 | 99.90 |
| ATOM | 2149 | CB | GLU | B  | 10  | 150.963 | 72.057 | 100.790 | 1.00 | 99.90 |
| ATOM | 2150 | N  | PRO | B  | 11  | 149.661 | 69.092 | 101.595 | 1.00 | 99.90 |
| ATOM | 2151 | CA | PRO | B  | 11  | 149.856 | 67.644 | 101.778 | 1.00 | 99.90 |
| ATOM | 2152 | C  | PRO | B  | 11  | 150.783 | 66.845 | 100.844 | 1.00 | 99.90 |
| ATOM | 2153 | O  | PRO | B  | 11  | 151.938 | 66.593 | 101.185 | 1.00 | 99.90 |
| ATOM | 2154 | CB | PRO | B  | 11  | 148.425 | 67.108 | 101.722 | 1.00 | 99.90 |
| ATOM | 2155 | CG | PRO | B  | 11  | 147.816 | 67.948 | 100.672 | 1.00 | 99.90 |
| ATOM | 2156 | CD | PRO | B  | 11  | 148.333 | 69.350 | 101.000 | 1.00 | 99.90 |
| ATOM | 2157 | N  | SER | B  | 12  | 150.258 | 66.422 | 99.691  | 1.00 | 99.90 |
| ATOM | 2158 | CA | SER | B  | 12  | 150.965 | 65.585 | 98.710  | 1.00 | 99.90 |
| ATOM | 2159 | C  | SER | B  | 12  | 150.922 | 64.167 | 99.292  | 1.00 | 99.90 |
| ATOM | 2160 | O  | SER | B  | 12  | 150.493 | 63.222 | 98.632  | 1.00 | 99.90 |
| ATOM | 2161 | CB | SER | B  | 12  | 152.410 | 66.042 | 98.440  | 1.00 | 99.90 |
| ATOM | 2162 | OG | SER | B  | 12  | 152.907 | 65.499 | 97.219  | 1.00 | 99.90 |

# Exercise

Download the PDB file of the Ribonuclease A (PDB: 7RSA) from the web (<http://ftp.rcsb.org/pub/pdb/data/structures/all/pdb/pdb7rsa.ent.gz>) and perform the following tasks

- Run a shell command to calculate the number of residues of the protein?
- Write a python script to parse the PDB file.
- Modify the program to calculate the distance between two atoms and residues.
- Calculate the average and standard deviation of the distance between two consecutive  $\alpha$  carbons?


# Defining protein structure

Basic information for the characterization of the protein three-dimensional structures are:


- $\phi, \psi$  values for each residue in the protein chain
- secondary structure
- solvent accessible area

# Ramachandran Plot

The backbone of the protein structure can be defined providing the list of  $\phi$ ,  $\psi$  angles for each residue in the chain.



# Ramachandran Analysis



<http://mordred.bioc.cam.ac.uk/~rapper/rampage.php>

# DSSP program

Program that implements the algorithm “Define Secondary Structure of Proteins”.

The method calculates different **features of the protein structure** such as the  $\phi$ ,  $\psi$  angles for each residue, its secondary structure and the solvent accessible area.

| # | RESIDUE | AA | STRUCTURE | BP1 | BP2 | ACC | ... | PHI | PSI | X-CA  | Y-CA   | Z-CA  |       |
|---|---------|----|-----------|-----|-----|-----|-----|-----|-----|-------|--------|-------|-------|
| 1 | 10      | B  | E         |     |     | 0   | 0   | 153 | ... | 360.0 | 144.2  | 150.1 |       |
| 2 | 11      | B  | P         |     | +   | 0   | 0   | 83  | ... | -90.2 | -84.0  | 149.9 |       |
| 3 | 12      | B  | S         | S   | >>  | S+  | 0   | 0   | 60  | ...   | 77.6   | -51.1 | 151.0 |
| 4 | 13      | B  | A         | T   | 34  | S+  | 0   | 0   | 6   | ...   | -82.3  | 73.7  | 151.3 |
| 5 | 14      | B  | D         | T   | 3>  | S+  | 0   | 0   | 39  | ...   | -154.6 | -41.3 | 147.5 |
| 6 | 15      | B  | W         | H   | <>  | S+  | 0   | 0   | 170 | ...   | -60.8  | -41.6 | 148.0 |
| 7 | 16      | B  | L         | H   | X   | S+  | 0   | 0   | 0   | ...   | -62.9  | -38.5 | 150.2 |
| 8 | 17      | B  | A         | H   | >   | S+  | 0   | 0   | 3   | ...   | -62.0  | -58.1 | 147.4 |
| 9 | 18      | B  | T         | H   | X   | S+  | 0   | 0   | 72  | ...   | -56.4  | -34.0 | 144.9 |


**SS**                    **SAA**                    **PHI**                    **PSI**  
**cols=**            **17**                    **36-38**                    **104-109**            **110-115**

DSSP: <ftp://ftp.cmbi.ru.nl/pub/software/dssp>  
more details at <https://swift.cmbi.umcn.nl/gv/dssp/>

Kabsch W, and Sander C, (1983). Biopolymers. 22 2577-2637.

# Relative solvent accessibility

The relative solvent accessible area is obtained dividing the accessible area of the residue by an estimation of its maximum accessible surface.



# Exercise

Download the DSSP file of the Ribonuclease A (PDB: 7RSA) from the web (<ftp://ftp.cmbi.umcn.nl/pub/molbio/data/dssp/7rsa.dssp>) and answer the following questions

- What is the total number of residues in helical and extended conformations?
- What is the average value of the  $\phi$  and  $\psi$  angles for the residues in helical and extended conformations?
- Are the average values falling the the correct region of the Ramachandran plot?
- Considering the solvent accessibility values reported in the DSSP file, calculate the relative solvent accessible area for Lysine, Valine and Glutamine with maximum solvent accessible area of 205, 142 and 198 respectively
- Are this value compatible with the physico-chemical properties of the residues?