

Systems Biology

A Model of the Regulatory Network Controlling Dendritic Cells Differentiation

Mauricio Pérez-Martínez^{1,2} and Luis Mendoza^{2,*}

¹Doctorado en Ciencias Biológicas, Universidad Nacional Autónoma de México. CDMX CP04510. México.

²Instituto de Investigaciones Biomédicas, Universidad Nacional Autónoma de México. CDMX CP04510. México.

*To whom correspondence should be addressed.

Associate Editor: XXXXXXXX

Received on XXXXX; revised on XXXXX; accepted on XXXXX

Abstract

Motivation: Dendritic cells have an important role during the immune response by acting as the main antigen-presenting cells. While there is experimental evidence about some key transcription factors and extracellular signals controlling their differentiation process, it is not yet clear the nature and dynamical properties of the regulatory network conformed by such molecules.

Results: We inferred the regulatory network that controls the differentiation process of murine dendritic cells. We then transformed the regulatory network into a discrete dynamical system, in the form of a Boolean model, to study its global properties. Our results are consistent with reported experimental expression patterns during the differentiation of these cells.

Contact: lmendoza@biomedicas.unam.mx

Supplementary information: Supplementary data are available at *Bioinformatics* online.

1 Introduction

The vertebrate immune system generates both first-line response, as well as delayed protection against pathogens. Dendritic cells (DCs) constitute a central element linking the innate and the adaptive immune system. Originally, the description of DCs were based on their morphology, namely, as “large cells with cytoplasm arranged in a variety of cell shapes ranging from bipolar elongate cells stellate or dendritic ones” (Steinman and Cohn, 1973). Later functional studies resulted in a more modern classification with three main groups: classical DCs (cDCs), epidermal or Langerhans cells (LCs), and plasmacytoid dendritic cells (pDCs) (Schuler *et al.*, 1985; Colonna *et al.*, 2004). Gene expression data has yet to provide a distinctive molecular signature for each of the three DC types. In fact, there might exist a tissue-specific molecular signature of DCs even if they are classified as classical or plasmacytoid subtypes (Alcantara-Hernandez *et al.*, 2017). Broadly speaking, DCs express some hematopoietic markers such as CD45, MHC-II, FLT3, and CD11c, while at the same time lack markers corresponding to T, B, NKs, granulocytes and erythrocytes lineage markers (Merad *et al.*, 2013). Also, they express some transcription factors broadly found in other immune cells, such as IRF8, IRF4, Id2, Zbtb46, E2-2, Batf3, the STATs, RelB, Ikaros, Notch, RBP-J, and PU.1 (Perié and Naik, 2015). Additionally, DCs express a variety of pattern recognition

receptor (PRRs), such as TLRs 1-11, and secrete some cytokines like IL-6, IL-10, and IL-12 (Manicassamy and Pulendran, 2009).

DCs have been the focus of much attention due to their potential therapeutic use, given that these cells are present at hematopoietic sites, environmental contact sites, filtering sites, and immune priming sites (Merad and Manz, 2009), as well as their ample repertoire of PRRs to recognize and uptake antigens, with the subsequent potential to stimulate specific T cell populations (Banchereau and Steinman, 1998; Shortman and Naik, 2007). These characteristics has prompted the efforts to generate *in vitro* particular DCs populations to generate specific adaptive responses (Harmon, 2011). For instance, some experimental approaches have been created against influenza and HIV infections (Boonnak *et al.*, 2013; Jacobson *et al.*, 2016), and recently there have been some promising results in the treatment of metastatic breast cancer (Zacharakis *et al.*, 2018).

DCs develop either from a common myeloid progenitor (CMP) or from a common lymphoid progenitor (CLP) (Manz *et al.*, 2001). FLT3L, which is produced by fibroblasts in the bone marrow, has been identified as the main external signal controlling the maturation process of DCs (Lisovsky *et al.*, 1996; Satpathy *et al.*, 2011; Murphy, 2016). Nonetheless, the precise sequence of molecular events during the differentiation of DCs still remain unknown.

Regulatory network modeling has been used successfully to study the global properties of several regulatory modules related to hematopoietic subprocesses (Mendoza and Méndez, 2015; Martínez-Sánchez *et al.*,




Fig. 1. Differentiation of dendritic cells. Arrows represent transitions from a progenitor to a more differentiated state. Names above arrows represent the main transcription factors reported during the process. Abbreviations are as follows: CDP, common dendritic progenitor, pre-cDC, pre-classical Dendritic Cell, pre-cDC1 and pre-cDC2, pre-classical Dendritic Cell type 1, and type 2, respectively. CD4+ cDCs and CD8+ cDCs, CD4+, and CD8+ classical dendritic cells respectively. pDC plasmacytoid dendritic cell.

2015; Méndez and Mendoza, 2016; Goode *et al.*, 2016; Hamey *et al.*, 2017; Ramirez and Mendoza, 2018; Liqutaya-Montiel and Mendoza, 2018). These studies have helped to understand how cells present specific molecular patterns under different physiological and experimental conditions as a direct result of the dynamical properties of an underlying regulatory network. We hereby present the regulatory network controlling the differentiation of DCs, inferred from a manual curation of experimental literature. The network is then converted into a Boolean model to study its global dynamical properties. The model recovers four attractors, which can be interpreted as the stable patterns of expression of the common dendritic progenitor (CDP), the pre-classical dendritic cell (pre-cDC), the pre-cDC type 1 (pre-cDC1), and the CD8+ classical dendritic cells (CD8+ cDCs). Furthermore, the model is able to describe the molecular patterns observed for single gain and loss-of-function mutations. Finally, we show that the model is able to describe the observed differentiation process from CDP to CD8+ cDCs.

2 Methods

2.1 CD8+ cDCs

The regulatory network controlling cDCs differentiation was inferred manually by reading a vast amount of experimental data available in the literature. Given the diversity of DCs types and sub-populations, and the varying amounts of molecular data available, it was decided to limit the study to the differentiation process of CD8+ cDCs splenic population in mice. The differentiation of CD8+ cDCs from a CMP is schematized in Fig. 1. These cells belong to DC1 DCs branch (Schlitzer *et al.*, 2015), they are tissue-resident, and their development is BATF3 and IRF8-dependent. Functionally, CD8+ cDCs recognize pathogens via TLR3, are capable of antigen cross-presentation, and stimulate CD8+ T cells (Shortman and Heath, 2010). It has been proposed that CD8+ cDCs murine have a human equivalent population defined as CD141+ (Villadangos and Shortman, 2010).

2.2 Modeling framework

The regulatory network was converted into a dynamical system, specifically to a Boolean network, to study its global properties. A Boolean network is a set of nodes representing molecules or molecular complexes as well as a set of regulatory directed interactions among them. In a Boolean network every node is described with a binary variable: 0 (OFF, repressed, or inactive) or 1 (ON, expressed, or active). The specific value of a node x_i is a function of the state of its regulators, so that $x_i(t+1) = F_i[x_1(t), x_2(t), \dots, x_n(t)]$, where F_i is a Boolean function containing any combination of the logic operators AND, OR, and NOT. The vector containing the values of all nodes at a given time, i.e., $X(t) = x_1(t), \dots, x_n(t)$, is the state of the network. Given the Boolean network has finite number of possible states, specifically 2^n , and that the system is deterministic, the trajectory created by successive iterations eventually reaches a previously visited network state. The trajectory consisting of a set of network states that repeat after n time steps is a period- n attractor. Period-1 attractors are also known as fixed points. Attractors represent phenotypes or cellular behaviors that can be directly compared to experimentally reported molecular patterns (Kauffman, 1969; Albert and Thakar, 2014).

We first studied the dynamical behavior of the wild type network, and then we explored the behavior of all possible gain and loss-of-function single mutants. Mutants were simulated by fixing a specific node to a value of 0 to simulate a null-mutation, or to a value of 1 to represent a constitutive activation. The attractors of the wild-type and mutant models were obtained using the R package BoolNet 2.1.3 (Müssel *et al.*, 2010).

A cell fate map of the model was reconstructed by performing dynamical simulation starting the system at a given attractor, and then making a bit-flip on the state of one node and then letting the network converge to an attractor. This temporary change of state was performed systematically on all the nodes of the network, for all the attractors of the model. For clarity, only those perturbations that move the system from one attractor to another were reported.

3 Results

3.1 The molecular basis of the regulatory network

The regulatory network controlling differentiation of splenic CD8+ cDCs contains 21 nodes representing cytokines, receptors, transcription factors, and other molecular markers. There are a total of 64 regulatory interactions among the nodes, where 46 are activations and 18 inhibitions (Figure 2). Sixty-two interactions were inferred from experimental reports, detailed ahead, and it was necessary to incorporate two hypothetical regulatory interactions in order to recover the experimentally observed behavior.

The CD8+ cDCs differentiation process starts with the FLT3 receptor activation by FLT3L, which is secreted by fibroblasts and stromal cells in the bone marrow, as well as by spleen cells (Lisovsky *et al.*, 1996; Miloud *et al.*, 2012). FLT3 is expressed in CDP, pre-cDC preDC1 progenitors, and in the final differentiated state. Constant activation of FLT3L/FLT3 is essential for dendropoiesis (Hochweller *et al.*, 2009). Once activated, FLT3 signals downstream to regulate its target genes including STAT3 (Li and Watowich, 2012). STAT3 can auto-activate (Ichiba *et al.*, 1998), and also activate PU.1 (Laouar *et al.*, 2003), NFIL3 (Vallania *et al.*, 2009), IFN α/β (Moore *et al.*, 2001), and BCL6 (Walker *et al.*, 2013). PU.1 self-activates (Okuno *et al.*, 2005), and is a positive regulator of FLT3 (Carotta *et al.*, 2010), BCL6 (Wei *et al.*, 2009), and IRF8 (Schonheit *et al.*, 2013). ID2 might have a self-activating regulation (Florio *et al.*, 1998; Liu *et al.*, 2000), and also activates BATF3 and IRF8 (Jaiswal *et al.*, 2013). ID2 and E2-2 mutually inhibit each other, thus regulating cell fate; ID2 suppression by E2-2 drives the cell fate to pDCs, and E2-2 suppression by ID2 drives the cell fate to cDCs (Jackson *et al.*, 2011). BATF3 can be activated by NFIL3 (Kashiwada *et al.*, 2011), and in turn it activates




Fig. 2. The regulatory network controlling the differentiation of splenic CD8+ cDCs in mice. Nodes (black squares) represent molecules. Sharp continuous arrows represent activations, while blunt discontinuous arrows represent inhibitions.

IRF8 and IRF4 (Grajales-Reyes *et al.*, 2015). IRF8 self-regulates positively (Kantakamalakul *et al.*, 1999; Bornstein *et al.*, 2014; Grajales-Reyes *et al.*, 2015) and has a mutual inhibition with IRF4 by which regulates the cell fate to CD8+ cDCs or CD4+ cDCs (Tamura *et al.*, 2005). Furthermore, both genes can be activated by IFN α/β (Guang-Nian *et al.*, 2015). In turn, IFN- α/β can be inhibited by IL-10 signalling (Ma *et al.*, 2015).

Experimentally, DCs can be generated using GM-CSF, which activates STAT5 and PU.1 at early differentiation states (Schiavoni *et al.*, 2002; Esashi and Liu, 2008). PU.1 has a mutual inhibition with GATA1, which in turn presents a self-activatory regulation (Zhang *et al.*, 1999; Nishimura *et al.*, 2000; Stopka *et al.*, 2005). GATA1 is able to activate IRF8 (Schonheit *et al.*, 2015), ID2 (Ji *et al.*, 2008), and NFIL3 (Choi *et al.*, 2015). Finally, NFIL3 is likely a positive regulator of ID2 (Xu *et al.*, 2015).

STAT3 and STAT5 have a central role during DCs differentiation (Onai and Manz, 2008). STAT3 is activated by TLR3, and is also part of the IL-10 signal transduction mechanism (Moore *et al.*, 2001; Williams *et al.*, 2004). Moreover, STAT3 is inhibited by GATA1 (Ezoe *et al.*, 2005) and by NF- κ B. Regarding STAT5, it most likely has a self-activating regulation (Metser *et al.*, 2016), activates ID2 (Li *et al.*, 2012) and RUNX1 (Ogawa *et al.*, 2008), inhibits IRF8 (Esashi and Liu, 2008; Waight *et al.*, 2014), and presents a mutual inhibition with BCL6 (Walker *et al.*, 2013). In turn, BCL6 activates IRF4 (Ci *et al.*, 2009), and inhibits both RUNX1 (Hatzi *et al.*, 2015) and NF- κ B (Perez-Rosado *et al.*, 2008).

Dendritic cell maturation involves pathogen recognition through the interaction between a pathogen-associated molecular pattern (PAMP) and a PRR. CD8+ cDCs express the toll-like receptor TLR3 induced by IFN α/β (Miettinen *et al.*, 2001). TLR3 activates the NF- κ B response (Jiang *et al.*, 2004), which in turn positively regulates PU.1 (Bonadies *et al.*, 2009), STAT5 (Katerndah *et al.*, 2017), ID2 (Wang *et al.*, 2012), IRF8 (Simon *et al.*, 2015), RUNX1 (Zhou *et al.*, 2015; Tang *et al.*, 2018), and IL-12 (T.L. *et al.*, 1995; Grohmann *et al.*, 1998; Hoentjen *et al.*, 2005), while inhibiting STAT3 (Ahmad *et al.*, 2015). RUNX1 is inhibited by IRF4 (Cao *et al.*, 2010). In turn, IL-12 is negatively regulated by IRF4 and NFIL3, but positively regulated by IRF8 (Akbari *et al.*, 2014; Kobayashi *et al.*, 2014). Finally, IL-12 can also be negatively regulated by IL-10 in an autocrine or niche-dependent way (Rennick *et al.*, 1997; Corinti *et al.*, 2001), and is self-regulated positively (Knolle *et al.*, 1998).

Despite the amount of published experimental data, we had to make some hypotheses regarding the existence of some regulatory interactions. Specifically, since there was no available information regarding the molecular basis of the regulation of CD8 expression, we included a self-activation, as well as its positive regulation over RUNX1, as originally proposed by a previous regulatory network model (Mendoza and Méndez, 2015).




Fig. 3. Attractors of the Boolean model. Wild-type model attractors. Active nodes are marked in black, and inactive nodes in gray. The corresponding cell types are indicated in the leftmost column.

3.2 Dynamics of the regulatory network

We constructed a Boolean model by assigning logical functions describing the activation of individual nodes as a function of its regulators (Table 1). Notice that two nodes, FLT3L and GM-CSF, act as input to the network. Either of these molecules are necessary for the differentiation of Dendritic Cells. Therefore, during the modeling process these two nodes are always active, unless stated otherwise. An exhaustive analysis of the resulting discrete dynamical system showed that the network has only four attractors, each of these being fixed-points that correspond to the general molecular activation patterns observed experimentally for the cell types CDP, precDC, pre-cDC1, and CD8+ cDCs (Figure 3).

There are small discrepancies between the model and the experimental data. Specifically, it has been reported that BCL6 expression is detected starting at the pre-cDC progenitor (Zhang *et al.*, 2014). In the model, however, BCL6 is detected also at the CDP stage. This is an indication that the model can be improved by looking maybe for a negative regulator. Then, it has been suggested that NFIL3 expression is restricted to CDP or pre-DC states (Kashiwada *et al.*, 2011; Merad *et al.*, 2013; Murphy *et al.*, 2016), while in the attractors this node is always in its off state. The model is thus probably missing a positive regulation over NFIL3.

Despite the relative minor discrepancies mentioned above, the model describes qualitatively the stationary expression patterns of key cell types during the differentiation of CD8+ cDCs. Furthermore, the model is able to qualitatively reproduce the differentiation process itself. CDPs are induced to differentiate by the constant presence of the extracellular signal FLT3L in the bone marrow or spleen, or by constant GM-CSF *in vitro* (Karsunky *et al.*, 2003; Xu *et al.*, 2007; Conti and Gessani, 2008; Helft *et al.*, 2015). An initial developmental decision is the commitment to either the pDC or cDC cell fates. This lineage separation is regulated by the mutual inhibition between ID2 and E2-2 (Belz and Nutt, 2012). The fate map (Figure 4) shows that in the model, under the constant input of either FLT3L or GM-CSF the activation of either BATF3 or ID2 moves the system from the CDP attractor to the pre-cDC attractor. It is worth mentioning that there might be a synergistic effect between ID2 and BATF3 during the differentiation process (Jaiswal *et al.*, 2013). Moreover, notice than in the fate map the presence of BATF3 is able to move the system from CDP trough pre-cDC and pre-CD1 and finally to cDC CD8+. This characteristic supports the idea that these cells might be characterized as BATF3-dependent cDCs (Murphy *et al.*, 2016).

IL-10 repression has been recognized as a maturation indicator during DC differentiation (Morel *et al.*, 1997). This event is also qualitatively recovered in the model, as the fate map shows that the inactivation of IL-10 moves the system from pre-cDC to pre-cDC1. Experimentally reports, however, suggest a gradual loss of sensibility of DCs to IL-10 (MacDonald *et al.*, 1999), thus preventing spontaneous activation of T cells (Corinti *et al.*, 2001).

Finally, results shown on the fate map suggest that the inactivation of BCL6 blocking, or the activation of E2-2 might move a pre-cDC to a CDP-like attractor (Figure 4) which exhibit the potential to differentiate to pDC or cDC branches given the expression of E2-2, ID2 and STAT3 but with the absence of NFIL3. This transition can be interpretes as a

Table 1. The logic rules determining the activity of all nodes in the network.

NODE	REGULATORY INTERACTIONS
BCL6	(STAT3 OR PU.1) AND NOT STAT5
BATF3	ID2 OR NFIL3
CD8	CD8
E2-2	NOT ID2
FLT3	FLT3L AND PU.1
FLT3L	is an input
GATA1	GATA1 AND NOT PU.1
GM-CSF	is an input
ID2	(NFIL3 OR STAT5 OR NF- κ B) AND NOT E2-2
IFN α/β	STAT3 AND NOT IL-10
IL-10	IL-10
IL-12	(NF- κ B AND IRF8) AND NOT IL-10 AND NOT NFIL3
IRF4	BATF3 AND BCL6 OR (IFN α/β AND NOT IRF8)
IRF8	(IRF8 OR IFN α/β OR BATF3 OR NF- κ B OR PU.1 OR ID2 OR RUNX1 OR GATA1) AND NOT IRF4 AND NOT STAT5
NF- κ B	TLR3 AND NOT BCL6
NFIL3	STAT3 AND GATA1
PU.1	(PU.1 OR GM-CSF OR STAT3 OR NF- κ B) AND NOT GATA1
RUNX1	(CD8 OR STAT5 OR NF- κ B) AND NOT IRF4 AND NOT BCL6
STAT3	(STAT3 AND TLR3) OR (FLT3 OR IL-10) AND NOT NF- κ B AND NOT GATA1
STAT5	(STAT5 AND GM-CSF OR NF- κ B) AND NOT BCL6
TLR3	IFN α/β

de-differentiation, from a more differentiated state to a progenitor or less differentiated state. This transition has not been reported experimentally, thus it is prediction of the present model.

3.3 Simulation of mutants and perturbations

We analyzed all possible single loss- and gain-of-function mutants of the model, however only those with biological relevance are reported and discussed in the main text (see Table 2). The attractors of all the simulated mutants are reported in the Supplementary Information. Importantly, models BCL6(+), BATF3(+), E2-2(-), FLT3(+), FLT3L(-), GATA1(-), ID2(+) IRF8(+), NF- κ B(-), NFIL3(-), PU.1(+), RUNX1(+), STAT3(+), and SATAT5(+) show no differences with respect to wild-type model in terms of the attained attractors.

FLT3L and GM-CSF act as partially redundant inputs to the network. Thus, eliminating only one signal does not affect the behavior of the model, but the double negative mutant FLT3L(-)/GM-CSF(-) does (Supplementary Table 43), where all the attractors are lost.

BCL6(-) mutant model shows a complete repression of IRF4 (Supplementary Table 1), but otherwise the resulting attractor are very similar to those obtained in the wild-type model. Experimentally, it is that IRF4 has an essential role during late CD4+ cDCs differentiation. Therefore, the lack of a major effect in our model is consistent with reports on the decreasing but not complete ablation of CD8+ cDCs number in BCL6 knockout models (Ohtsuka *et al.*, 2011). A somewhat similar effect is observed in the BATF3(-) mutant model (Supplementary Table 3), showing a repression of both IRF4 and IL-12. This might correlate with the experimental observation that BATF3 suppression results in an inefficient stimulation of CD8+ cDCs, rather than a reduction in their number (Montagna *et al.*, 2015).

PU.1 is involved the differentiation process of some hematopoietic lineages, particularly at the level of progenitors (Fisher and Scott, 1998; Iwasaki *et al.*, 2005). In the model, the PU.1(-) mutant results in the loss of all attractors (Supplementary Table 33), which is consistent with the report that this mutant shows impaired DC formation *in vitro* (Carotta *et al.*, 2010).

The IRF8(-) mutant model (Supplementary Table 27) has an extra attractor somewhat similar to that expected for CD4+ cDCs, characterized by the FLT3L/FLT3 dependence, the presence of ID2 and IRF4, as well as the absence of IL-12 production, and the expression of TLR3. However, given the some of the key molecular markers of these cells (such as NOTCH




Fig. 4. Fate map. Arrows represent transition between a progenitor cellular state into a more differentiated state, and names above represent possible single node effects leading these transitions. Dashed arrow represent theoretical possible transition (see discussion section).

Table 2. Some relevant mutants

Mutant	Effect on the model	Experimental counterpart
IRF8 loss	cDC CD4-like attractor	Murphy et al., 2016
FLT3L loss	No effect	Karsunky et al., 2003
NFIL3 gain	No IL-12	Kashiwada et al. 2011
PU.1 loss	Unspecific attractors	Carotta et al., 2010
BCL6 loss	IRF4 repression	Ohtsuka et al., 2011
BATF3 loss	IRF4 repression, no IL-12	Montagna et al., 2015
ID2 loss	pDC-like attractor	Ghosh et al., 2010
STAT5 loss	pDC-like attractor	Li et al., 2012

signalling, some IRF4 regulators, IL-23 production, etc.) are not included in the present version of the model, it is not possible to unequivocally describe a CD4+ attractor.

The ID2(-) and STAT5(-) mutant models (Supplementary Tables 17 and 39) reveal a pDC-like attractor characterized by the absence of cDC-related regulators IRF4, BATF3, NFIL3, no IL-12 production and no TLR3 expression, but with the FLT3L/FLT3 dependence, and STAT3, IRF8 and E2-2 presence. It is known that ID2 and STAT5 suppression leads the E2-2 and STAT3 action, both related to pDC differentiation program (Cisse et al., 2008; Li et al., 2012).

4 Discussion

DCs are a key component of the cellular immune response, but the regulatory mechanisms controlling their differentiation remains unclear. We used publicly available information to reconstruct a regulatory network involved in the differentiation of splenic CD8+ DC from the common dendritic progenitor. The model of the network as a Boolean model recovers the expression patterns observed in CDPs, pre-cDCs, pre-cDC1s, and CD8+ cDCs. While there have been previous efforts to uncover the molecular networks associated to dendritic cells (see, for example Pandey et al. (2013); Lin et al. (2015)), to the best of our knowledge this is the

first network of its kind analyzed as a dynamical system to describe the differentiation process of CD8+ cDCs.

The definition of DC lineages has been cause of controversy for many years, given that they may have either a myeloid or a lymphoid origin. Accordingly, there have been some attempts to redefine DCs lineages based on specific molecular characteristics (Perié and Naik, 2015; Murphy et al., 2016). The network presented here contains nodes that are shared with other myeloid and lymphoid lineages (Mendoza and Méndez, 2015; Méndez and Mendoza, 2016; Ramirez and Mendoza, 2018; Liquitaya-Montiel and Mendoza, 2018), and hence it allows for its integration into a more complex network that may help in the understanding of the different possible differentiation routes to generate dendritic cells during hematopoiesis.

Acknowledgements

This paper constitutes a part of the doctoral studies of M.P.-M., who thanks the graduate program Doctorado en Ciencias Biológicas, UNAM.

Funding

This work was supported by CONACYT scholarship 625478 to M.P.-M., and grant UNAM-DGAPA-PAPIIT IN200918 to L.M.

References

Ahmad, S. et al. (2015). Regulation of TNF- α and NF- κ B activation through the JAK/STAT signaling pathway downstream of histamine 4 receptor in a rat model of LPS-induced joint inflammation. *Immunobiology*, **220**(7), 889–898.

Akbari, M. et al. (2014). IRF4 in dendritic cells inhibits IL-12 production and controls Th1 immune responses against *Leishmania major*. *J Immunol*, **192**(5), 2271–2279.

Albert, R. and Thakar, J. (2014). Boolean modeling: A logic-based dynamic approach for understanding signaling and regulatory networks and for making useful predictions. *WIREs Syst Biol Med*, **6**, 353–369.

Alcantara-Hernandez, M. et al. (2017). High-dimensional phenotypic mapping of human dendritic cells reveals interindividual variation and tissue specialization. *Immunity*, **47**, 1037–1050.

Banchereau, J. and Steinman, R. M. (1998). Dendritic cells and the control of immunity. *Nature*, **392**(6673), 245–252.

Belz, G. T. and Nutt, S. L. (2012). Transcriptional programming of the dendritic cell network. *Nat Rev Immunol*, **12**(2), 101–113.

Bonadies, N. et al. (2009). PU.1 is regulated by NF- κ B through a novel binding site in a 17 kb upstream enhancer element. *Oncogene*, **29**, 1062–1072.

Boonnak, K. et al. (2013). Antigen-activated dendritic cells ameliorate influenza A infections. *J Clin Invest*, **123**(7), 2850–2861.

Bornstein, C. et al. (2014). A negative feedback loop of transcription factors specifies alternative dendritic cell chromatin states. *Mol Cell*, **56**(6), 749–762.

Cao, Y. et al. (2010). Interferon regulatory factor 4 regulates thymocyte differentiation by repressing RUNX3 expression. *Eur J Immunol*, **40**(11), 3198–3209.

Carotta, S. et al. (2010). The transcription factor PU.1 controls dendritic cell development and FLT3 cytokine receptor expression in a dose-dependent manner. *Immunity*, **35**(5), 628–641.

Choi, K. et al. (2015). GATA1 overexpression in neurons increases the expression of cell-mediated cytotoxicity-related genes. *Anim Cells Syst*, **20**(1), 31–38.

- Ci, W. *et al.* (2009). The BCL6 transcriptional program features repression of multiple oncogenes in primary B cells and is deregulated in DLBCL. *Blood*, **113**(22), 5536–5548.
- Cisse, B. *et al.* (2008). Transcription factor E2-2 is an essential and specific regulator of plasmacytoid dendritic cell development. *Cell*, **135**(1), 37–48.
- Colonna, M. *et al.* (2004). Plasmacytoid dendritic cells in immunity. *Nat. Immunol.*, **5**, 1219–26.
- Conti, L. and Gessani, S. (2008). GM-CSF in the generation of dendritic cells from human blood monocyte precursors: Recent advances. *Immunobiology*, **213**(9–10), 859–870.
- Corinti, S. *et al.* (2001). Regulatory activity of autocrine IL-10 on dendritic cell functions. *J Immunol*, **166**(7), 4312–4318.
- Esashi, E. and Liu, Y.-J. (2008). E-box protein E2-2 is a crucial regulator of plasmacytoid DC development. *Eur J Immunol*, **38**(9), 2386–2388.
- Ezoe, S. *et al.* (2005). GATA transcription factors inhibit cytokine-dependent growth and survival of a hematopoietic cell line through the inhibition of STAT3 activity. *J Biol Chem*, **280**(13), 13163–13170.
- Fisher, R. C. and Scott, E. W. (1998). Role of PU.1 in hematopoiesis. *Stem cells (Dayton, Ohio)*, **16**(1), 25–37.
- Florio, M. *et al.* (1998). ID2 promotes apoptosis by a novel mechanism independent of dimerization to basic helix-loop-helix factors. *Mol Cell Biol*, **18**(9), 5435–5444.
- Goode, D. *et al.* (2016). Dynamic gene regulatory networks drive hematopoietic specification and differentiation. *Cell*, **36**, 572–587.
- Grajales-Reyes, G. E. *et al.* (2015). BATF3 maintains autoactivation of IRF8 for commitment of a CD8(+) conventional DC clonogenic progenitor. *Nat Immunol*, **16**(7), 708–717.
- Grohmann, U. *et al.* (1998). IL-12 acts directly on DC to promote nuclear localization of NF-kappaB and primes DC for IL-12 production. *Immunity*, **9**(3), 315–323.
- Guang-Nian, Z. *et al.* (2015). Interferon regulatory factors: At the crossroads of immunity, metabolism, and disease. *Biochim Biophys Acta (BBA) - Molecular Basis of Disease*, **1852**(2), 365–378.
- Hamey, F. *et al.* (2017). Reconstructing blood stem cell regulatory network models from single-cell molecular profiles. *PNAS*, **14**(23), 5822–5829.
- Harmon, K. (2011). The patient scientist. *Sci Am*, **306**(1), 54–59.
- Hatzi, K. *et al.* (2015). BCL6 orchestrates Tfh cell differentiation via multiple distinct mechanisms. *J Exp Med*, **212**(4), 539–553.
- Helft, J. *et al.* (2015). GM-CSF mouse bone marrow cultures comprise a heterogeneous population of CD11c(+)MHCII(+) macrophages and dendritic cells. *Immunity*, **42**(6), 1197–1211.
- Hochweller, K. *et al.* (2009). Homeostasis of dendritic cells in lymphoid organs is controlled by regulation of their precursors via a feedback loop. *Blood*, **114**(20), 4411–4421.
- Hoentjen, F. *et al.* (2005). STAT3 regulates NF-κB recruitment to the IL-12p40 promoter in dendritic cells. *Blood*, **105**(2), 689–696.
- Ichiba, M. *et al.* (1998). Autoregulation of the STAT3 gene through cooperation with a cAMP-responsive element-binding protein. *J Biol Chem*, **273**(11), 6132–6138.
- Iwasaki, H. *et al.* (2005). Distinctive and indispensable roles of PU.1 in maintenance of hematopoietic stem cells and their differentiation. *Blood*, **106**(5), 1590–1600.
- Jackson, J. T. *et al.* (2011). ID2 expression delineates differential checkpoints in the genetic program of CD8+ and CD103+ dendritic cell lineages. *The EMBO journal*, **30**(13), 2690–2704.
- Jacobson, J. *et al.* (2016). Dendritic cell immunotherapy for HIV-1 infection using autologous HIV-1 RNA: A randomized, double-blind, placebo-controlled clinical trial. *J Acquir Immune Defic Syndr*, **72**(1), 31–38.
- Jaiswal, H. *et al.* (2013). BATF3 and ID2 have a synergistic effect on IRF8-directed classical CD8+ dendritic cell development. *J Immunol*, **191**(12), 5993–6001.
- Ji, M. *et al.* (2008). ID2 intrinsically regulates lymphoid and erythroid development via interaction with different target proteins. *Blood*, **112**(4), 1068–1077.
- Jiang, Z. *et al.* (2004). Toll-like receptor 3-mediated activation of NF-κB and IRF3 diverges at Toll-IL-1 receptor domain-containing adapter inducing IFN. *PNAS*, **101**(10), 3533–3538.
- Kantakamalakul, W. *et al.* (1999). Regulation of ifn consensus sequence binding protein expression in murine macrophages. *J Immunol*, **162**(12), 7417–7425.
- Karsunky, H. *et al.* (2003). FLT3 ligand regulates dendritic cell development from FLT3+ lymphoid and myeloid-committed progenitors to FLT3+ dendritic cells in vivo. *J Exp Med*, **198**(2), 305–313.
- Kashiwada, M. *et al.* (2011). NFIL3/E4BP4 is a key transcription factor for CD8+ dendritic cell development. *Blood*, **117**(23), 6193–6197.
- Katerndah, C. *et al.* (2017). Antagonism of B cell enhancer networks by STAT5 drives leukemia and poor patient survival. *Nat Immunol*, **18**(6), 694–704.
- Kauffman, S. (1969). Metabolic stability and epigenesis in randomly constructed genetic nets. *J Theor Biol*, **22**(3), 437–467.
- Knolle, P. *et al.* (1998). Interleukin-10 expression is autoregulated at the transcriptional level in human and murine kupffer cells. *Hepatology*, **27**(1), 93–99.
- Kobayashi, T. *et al.* (2014). NFIL3-deficient mice develop microbiota-dependent, IL-12/23-driven spontaneous colitis. *J Immunol*, **192**(4), 1918–1927.
- Laouar, Y. *et al.* (2003). STAT3 is required for FLT3L-dependent dendritic cell differentiation. *Immunity*, **19**(6), 903–912.
- Li, H. and Watowich, S. (2012). A STATus report on DC development. *J Leukoc Biol.*, **92**(3), 445–459.
- Li, H. *et al.* (2012). The signal transducers STAT5 and STAT3 control expression of ID2 and E2-2 during dendritic cell development. *Blood*, **120**(22), 4363–4373.
- Lin, Q. *et al.* (2015). Epigenetic program and transcription factor circuitry of dendritic cell development. *Nucleic Acids Res*, **43**(20), 9680–9693.
- Liquitaya-Montiel, A. J. and Mendoza, L. (2018). Dynamical analysis of the regulatory network controlling Natural Killer cells differentiation. *Front. Physiol*, **9**(1029), 1–10.
- Lisovsky, M. *et al.* (1996). FLT3-ligand production by human bone marrow stromal cells. *Leukemia*, **10**(6), 1012–1018.
- Liu, J. *et al.* (2000). A cysteine residue in the helix-loop-helix domain of ID2 is critical for homodimerization and function. *Biochem Biophys Res Commun*, **273**(3), 1042–1047.
- Ma, X. *et al.* (2015). Regulation of IL-10 and IL-12 production and function in macrophages and dendritic cells. *F1000Research*, **4**(1465), 1–13.
- MacDonald, K. *et al.* (1999). Resistance of rheumatoid synovial dendritic cells to the immunosuppressive effects of IL-10. *J Immunol*, **163**, 5599–5607.
- Manicassamy, S. and Pulendran, B. (2009). Modulation of adaptive immunity with toll-like receptors. *Semin Immunol*, **21**, 185–193.
- Manz, M. *et al.* (2001). Dendritic cell potentials of early lymphoid and myeloid progenitors. *Blood*, **97**, 3333–3341.
- Martinez-Sanchez, M. *et al.* (2015). A minimal regulatory network of extrinsic and intrinsic factors recovers observed patterns of CD4+ T cell differentiation and plasticity. *PLoS Comput Biol*, **11**(6), 1–23.
- Mendoza, L. and Méndez, A. (2015). A dynamical model of the regulatory network controlling lymphopoiesis. *Bio Systems*, **137**, 25–33.
- Merad, M. and Manz, M. (2009). Dendritic cell homeostasis. *Blood*, **113**(15), 3418–3427.
- Merad, M. *et al.* (2013). The dendritic cell lineage: Ontogeny and function of dendritic cells and their subsets in the steady state and the inflamed

- setting. *Annu Rev Immunol*, **31**, 563–604.
- Metser, G. *et al.* (2016). An autoregulatory enhancer controls mammary-specific STAT5 functions. *Nucleic Acids Res*, **44**(3), 1052–1063.
- Miettinen, M. *et al.* (2001). IFNs activate toll-like receptor gene expression in viral infections. *Genes Immun*, **2**, 349–355.
- Miloud, T. *et al.* (2012). Organ-specific cellular requirements for in vivo dendritic cell generation. *J Immunol (Baltimore, Md. : 1950)*, **188**(3), 1125–1135.
- Montagna, G. N. *et al.* (2015). BATF3 deficiency proves the pivotal role of CD8(+) dendritic cells in protection induced by vaccination with attenuated plasmodium sporozoites. *Parasite Immunol*, **37**(10), 533–543.
- Moore, K. *et al.* (2001). Interleukin-10 and the interleukin-10 receptor. *Ann Rev Immunol*, **19**, 683–765.
- Morel, A.-S. *et al.* (1997). Split activity of interleukin-10 on antigen capture and antigen presentation by human dendritic cells: Definition of a maturative step. *Eur. J. Immunol.*, **27**, 26–34.
- Murphy, K. M. (2016). Transcriptional control of dendritic cell development. *Adv Immunol*, **120**, 239–257.
- Murphy, T. L. *et al.* (2016). Transcriptional control of dendritic cell development. *Annu Rev Immunol*, **34**, 93–119.
- Méndez, A. and Mendoza, L. (2016). A network model to describe the terminal differentiation of B cells. *PLoS ONE*, **11**(12), 1–26.
- Müssel, C. *et al.* (2010). Boolnet – an R package for generation, reconstruction and analysis of boolean networks. *Bioinformatics*, **26**(10), 1378–1380.
- Nishimura, S. *et al.* (2000). A GATA box in the GATA-1 gene hematopoietic enhancer is a critical element in the network of GATA factors and sites that regulate this gene. *Mol Cell Biol*, **20**(2), 713–723.
- Ogawa, S. *et al.* (2008). Physical and functional interactions between STAT5 and RUNX transcription factors. *J Biochem*, **143**(5), 695–709.
- Ohtsuka, H. *et al.* (2011). BCL6 is required for the development of mouse CD4+ and CD8+ dendritic cells. *J Immunol (Baltimore, Md. : 1950)*, **186**(1), 255–263.
- Okuno, Y. *et al.* (2005). Potential autoregulation of transcription factor PU.1 by an upstream regulatory element. *Mol Cell Biol*, **27**(5), 2832–2845.
- Onai, N. and Manz, M. G. (2008). The STATs on dendritic cell development. *Immunity*, **28**(4), 490–492.
- Pandey, G. *et al.* (2013). Decoding dendritic cell function through module and network analysis. *J Immunol Methods*, **387**(1–2), 71–80.
- Perez-Rosado, A. *et al.* (2008). BCL6 represses NFkappaB activity in diffuse large B-cell lymphomas. *J Pathol*, **214**(4), 498–507.
- Perié, L. and Naik, S. H. (2015). Toward defining a ‘lineage’ – the case for dendritic cells. *Semin Cell Dev Biol*, **41**, 3–8.
- Ramirez, C. and Mendoza, L. (2018). Phenotypic stability and plasticity in GMP-derived cells as determined by their underlying regulatory network. *Bioinformatics*, **34**(7), 1174–1182.
- Rennick, D. M. *et al.* (1997). Studies with IL-10-/- mice: an overview. *J Leukoc Biol*, **61**(4), 389–396.
- Satpathy, A. T. *et al.* (2011). Transcription factor networks in dendritic cell development. *Semin Immunol*, **23**(5), 388–397.
- Schiavoni, G. *et al.* (2002). ICSBP is essential for the development of mouse type I interferon-producing cells and for the generation and activation of CD8alpha(+) dendritic cells. *J Exp Med*, **196**(11), 1415–1425.
- Schlitzer, A. *et al.* (2015). Identification of cDC1- and cDC2-committed DC progenitors reveals early lineage priming at the common DC progenitor stage in the bone marrow. *Nat Immunol*, **16**(7), 718–728.
- Schonheit, J. *et al.* (2013). PU.1 level-directed chromatin structure remodeling at the IRF8 gene drives dendritic cell commitment. *Cell Rep*, **3**, 1617–1628.
- Schonheit, J. *et al.* (2015). Chromatin dynamics during differentiation of myeloid cells. *J Mol Biol*, **427**(3), 670–687.
- Schuler, G. *et al.* (1985). A comparison of murine epidermal langerhans cells with spleen dendritic cells. *Dermatol*, **85**, 99–106.
- Shortman, K. and Heath, W. R. (2010). The CD8+ dendritic cell subset. *Immunol Rev*, **234**(1), 18–31.
- Shortman, K. and Naik, S. H. (2007). Steady-state and inflammatory dendritic-cell development. *Nat Rev Immunol*, **7**(1), 19–30.
- Simon, P. *et al.* (2015). The NF-κB p65 and p50 homodimer cooperate with IRF8 to activate iNOS transcription. *BMC Cancer*, **15**(770), 1–12.
- Steinman, R. M. and Cohn, Z. A. (1973). Identification of a novel cell type in peripheral lymphoid organs of mice. *J Exp Med*, **137**(5), 1142–1162.
- Stopka, T. *et al.* (2005). PU.1 inhibits the erythroid program by binding to GATA-1 on DNA and creating a repressive chromatin structure. *The EMBO journal*, **24**(21), 3712–3723.
- Tamura, T. *et al.* (2005). IFN regulatory factor-4 and -8 govern dendritic cell subset development and their functional diversity. *J Immunol*, **174**(5), 2573–2581.
- Tang, X. *et al.* (2018). RUNX1: A regulator of NF-κB signaling in pulmonary diseases. *Curr Protein Pept Sci*, **19**(2), 172–178.
- T.L., M. *et al.* (1995). Regulation of interleukin 12 p40 expression through an NF-κB half-site. *Mol Cell Biol*, **15**(10), 5258–5267.
- Vallania, F. *et al.* (2009). Genome-wide discovery of functional transcription factor binding sites by comparative genomics: The case of STAT3. *PNAS*, **106**(13), 5117–5122.
- Villadangos, J. A. and Shortman, K. (2010). Found in translation: The human equivalent of mouse CD8+ dendritic cells. *J Exp Med*, **207**(6), 1131–1134.
- Waight, J. *et al.* (2014). Regulation of the Interferon regulatory factor-8 (IRF-8) tumor suppressor gene by the Signal Transducer and Activator of Transcription 5 (STAT5) transcription factor in chronic myeloid leukemia. *J Biol Chem*, **289**(22), 15642–15652.
- Walker, S. *et al.* (2013). STAT5 outcompetes STAT3 to regulate the expression of the oncogenic transcriptional modulator BCL6. *Mol Cell Biol*, **33**(15), 2879–2890.
- Wang, C. *et al.* (2012). ID2 regulates the proliferation of squamous cell carcinoma in vitro via the NF-κB/Cyclin D1 pathway. *Chin J Cancer*, **31**(9), 430–439.
- Wei, F. *et al.* (2009). PU.1 can recruit BCL6 to DNA to repress gene expression in germinal center B cells. *Mol Cell Biol*, **29**(17), 4612–4622.
- Williams, L. *et al.* (2004). Signal transducer and activator of transcription 3 is the dominant mediator of the anti-inflammatory effects of IL-10 in human macrophages. *J Immunol*, **172**(1), 567–576.
- Xu, W. *et al.* (2015). NFIL3 orchestrates the emergence of common helper innate lymphoid cell precursors. *Cell Rep*, **10**(12), 2043–2054.
- Xu, Y. *et al.* (2007). Differential development of murine dendritic cells by GM-CSF versus FLT3 ligand has implications for inflammation and trafficking. *J Immunol*, **179**(11), 7577–7584.
- Zacharakis, N. *et al.* (2018). Immune recognition of somatic mutations leading to complete durable regression in metastatic breast cancer. *Nat Med*, **24**, 724–730.
- Zhang, P. *et al.* (1999). Negative cross-talk between hematopoietic regulators: GATA proteins repress PU.1. *PNAS*, **96**(15), 8705–8710.
- Zhang, T.-T. *et al.* (2014). Dynamic expression of BCL6 in murine conventional dendritic cells during in vivo development and activation. *PLOS ONE*, **9**(6), 1–12.
- Zhou, J. *et al.* (2015). Aberrant nuclear factor-kappa B activity in acute myeloid leukemia: From molecular pathogenesis to therapeutic target. *Oncotarget*, **6**(8), 5490–5500.