EDUCATIONAL ARTICLE

Unraveling *Leishmania donovani*: Parasite Biology, Pathogenesis & Diagnostic (Part 1)

Anjali Prabha | B.Sc. Zoology Undergraduate | Aspiring Computational Biologist

About this article

This article is an educational overview prepared to consolidate learning about visceral leishmaniasis, the life cycle of *Leishmania donovani*, selected host-parasite interactions, and commonly used diagnostic approaches. It is not an original research paper, a systematic review, clinical guidance, or an official publication of any institute. Diagnostic performance varies with the test, specimen, study population, and clinical setting.

DISEASE CONTEXT

- **>95% Mortality:** Fatal if untreated; high-priority WHO Neglected Tropical Disease.
- **Endemic Hotspot:** Gangetic plains (Bihar, Jharkhand, West Bengal).

PARASITE FORMS

- **Dimorphic Life Cycle:** Extracellular promastigote (vector) vs. intracellular amastigote (host).
- **Immune Interaction:** LPG & GP63 are key molecules studied in host-parasite interactions and immune evasion.

DIAGNOSTIC GOLD STANDARD

- **>98% Sensitivity:** kDNA/18S qPCR offers non-invasive, ultra-sensitive detection.
- **Monitoring:** Helps monitor parasite clearance and assess response to therapy.

1. Disease Context & Research Apex (ICMR-RMNIHR)

The Threat of Kala-Azar (Black Fever)

- **Pathogen & Severity:** Caused by *Leishmania donovani*; untreated VL can be fatal, described by WHO as fatal in >95% of cases if left untreated.
- **Vector:** Transmitted to humans through the bites of infected female phlebotomine sandflies (principally *Phlebotomus argentipes* in South Asia).
- **Endemic Burden:** Historically concentrated across the Gangetic plains, including Bihar, Jharkhand, and West Bengal.
- **Clinical Signs:** Characterized by prolonged irregular fever, weight loss, enlargement of the spleen and liver, and anemia.

ICMR-RMNIHR Patna Research Focus

- **Premier Research Center:** ICMR-RMNIHR in Patna is a premier national research center dedicated to visceral leishmaniasis research.
- **Drug Efficacy & Resistance:** Conducts essential research on drug susceptibility, therapeutic efficacy, and tracking antimonial resistance.
- **Diagnostics Development:** Evaluates field diagnostic tests like rK39 RDTs and refines molecular detection methods like qPCR.
- **Elimination Programs:** Supports regional elimination efforts through vector surveillance and evidence-based epidemiological strategies.

2. Lifecycle Dynamics (Digenetic Transition)

1. Vector Stage	2. Inoculation	3. Intracellular Form	4. Lysis & Spread
• **Midgut Growth:** Amastigotes transform into procyclic promastigotes. • **Differentiation:** Differentiate into infectious metacyclic promastigotes that migrate to the sandfly proboscis.	• **Transmission:** Injected into the host skin during the sandfly blood meal. • **Phagocytosis:** Engulfed by host macrophages and mononuclear phagocytes.	• **Transformation:** Promastigotes lose their flagella and transform into amastigotes inside phagolysosomes. • **Host Survival:** Adapted to survive and replicate within the acidic phagolysosome environment.	• **Binary Fission:** Amastigotes multiply repeatedly until host cell ruptures. • **Dissemination:** Invade adjacent monocytes; spread to spleen, liver, and bone marrow.

Promastigote vs. Amastigote Stage Comparison

Promastigote Stage (Vector Form)	Amastigote Stage (Host / LD Bodies)
• **Host & Niche:** Extracellular; female sandfly midgut & proboscis; 22–28°C, neutral pH (7.0–7.4). • **Morphology:** Elongated, slender, lanceolate (10–20 µm long, 1.5–2.5 µm wide). • **Motility:** Highly motile via single long anterior flagellum (15–20 µm). • **Kinetoplast:** Positioned anterior to the central spherical nucleus. • **Role:** Metacyclic promastigote is the infectious transmission unit.	• **Host & Niche:** Obligate intracellular inside macrophages; 37°C, acidic phagolysosome (pH 4.5–5.5). • **Morphology:** Small, ovoid or spherical (2–4 µm diameter). • **Motility:** Non-motile; flagellum internalized and rudimentary. • **Kinetoplast:** Rod-shaped, adjacent to eccentric nucleus. • **Role:** Replicative, pathogenic driver of clinical visceral leishmaniasis.

3. Host Invasion & Immune Evasion

1. Surface Defense & Entry	2. Phagosome Arrest	3. Signaling Disruption	4. Immune Polarization
• **Surface LPG:** Lipophosphoglycan (LPG) covers the promastigote surface and plays important roles during sandfly midgut attachment and initial complement resistance. • **Entry Mediators:** LPG and surface proteins facilitate binding and silent entry into host macrophages via complement receptors.	• **Maturation Delay:** LPG is recognized for transiently delaying phagosome maturation, helping promastigotes establish infection. • **Differentiation Window:** This temporary delay gives the parasite time to differentiate into the acid-tolerant amastigote form.	• **GP63 Protease:** Major surface metalloprotease (GP63) is a major parasite virulence factor studied for cleaving host cellular substrates. • **Host Signaling Impact:** Modulates host macrophage signaling cascades, dampening microbicidal responses and promoting parasite survival.	• **Immune Balance:** Active visceral leishmaniasis is accompanied by suppression of cell-mediated immunity and impaired Th1 responses. • **Parasite Persistence:** Elevation of anti-inflammatory cytokines like IL-10 downregulates macrophage leishmanicidal mechanisms, promoting persistence.

4. Diagnostic Matrix: What Each Method Can & Cannot Tell Us

1. Serological (rK39 RDT & ELISA)

- **Target Marker:** Circulating host IgG antibodies specific to recombinant kinesin-related antigen rK39 (39-amino-acid repeat).
- **Sensitivity & Specificity:** High sensitivity and specificity in South Asia; rapid, field-deployable strip test format.
- **Invasiveness:** Minimal; requires only a finger-prick whole blood droplet for lateral flow strip testing.
- **Cure Assessment:** Cannot distinguish current active infection from past exposure, as anti-Leishmania antibodies persist after cure.

2. Microscopy (Giemsa Stain)

- **Target Marker:** Direct visual identification of intracellular amastigotes (LD bodies with nucleus and kinetoplast).
- **Sensitivity & Specificity:** High specificity (definitive evidence of infection); sensitivity varies depending on tissue source.
- **Invasiveness:** High; splenic aspiration carries fatal hemorrhage risks; bone marrow causes severe patient discomfort.
- **Cure Assessment:** Can confirm parasitological cure when negative, but repeat aspiration carries procedural risk.

3. Molecular (kDNA / 18S qPCR)

- **Target Marker:** Conserved kDNA minicircles (~10,000 copies/parasite) or nuclear 18S rRNA genes.
- **Sensitivity & Specificity:** Extremely high sensitivity and analytical specificity using conserved target sequences.
- **Invasiveness:** Minimal; requires only routine peripheral venous blood (EDTA) or buffy coat.
- **Cure Assessment:** Enables monitoring of parasite clearance, detection of early relapse, and post-treatment assessment.

5. Key Takeaways & Part 2 Benchtop Preview

Key Takeaways: Diagnostic Paradigm Shift

- **Shift from Invasive Biopsies:** Definitive diagnosis is moving away from hazardous splenic aspirates toward molecular testing.
- **Role of Serology:** rK39 RDTs remain essential for field screening, but persisting antibodies limit their role in monitoring cure.
- **Molecular Monitoring:** Molecular assays provide non-invasive options for tracking parasite clearance, diagnosing PKDL, and identifying relapse.

Preview: Part 2 Benchtop Protocol

- **Wet-Lab Transition:** In Part 2, we transition from theoretical biology to wet-lab protocols: step-by-step procedures for whole-blood collection and genomic DNA extraction.
- **Molecular Workflow:** Spectrophotometric purity QC via NanoDrop (A260/A280 ratios), real-time kDNA qPCR amplification kinetics, and post-amplification gel electrophoresis.

Author's Note

This educational article was prepared for learning and portfolio purposes. It is not an original research publication and does not represent an official statement or endorsement by any institution.