

bioinformatics.ca

bioinformaticsdotca.github.io



CC BY-SA 4.0 DEED

Attribution-ShareAlike 4.0 International

Canonical URL : https://creativecommons.org/licenses/by-sa/4.0/

See the legal code

You are free to:

Share — copy and redistribute the material in any medium or format for any purpose, even commercially.

Adapt — remix, transform, and build upon the material for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:

Attribution — You must give appropriate credit , provide a link to the license, and indicate if changes were made . You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

ShareAlike — If you remix, transform, or build upon the material, you must distribute your contributions under the same license as the original.

No additional restrictions — You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits.

Notices:

You do not have to comply with the license for elements of the material in the public domain or where your use is permitted by an applicable exception or limitation .


No warranties are given. The license may not give you all of the permissions necessary for your intended use. For example, other rights such as publicity, privacy, or moral rights may limit how you use the material.

Module 4: Genome Assembly, Polishing, Evaluation and Annotation

Dr. Ido Hatam

CBW High-Throughput Genomics

November 4-6, 2026




Instructor affiliation logos go here

Learning Objectives

- Explain the difference between de novo and reference-based assembly
- Explain how polishing improves assembly accuracy
- Explain how contiguity, correctness and completeness are evaluated
- Read and interpret Flye, QUAST and BUSCO reports
- Distinguish structural from functional annotation and read their outputs (GFF3, eggNOG table)

Where we are in the workflow




= this module (dashed outline = supplied or reference)

The case: can we rebuild the wolf's genome, one piece per chromosome?

What is genome assembly?


- Sequencing gives millions of pieces: **reads**. None spans a chromosome
- **Assembly**: rebuild the genome from the reads alone
- Overlapping reads are joined; the **consensus** (the base most reads agree on) is kept
- The result: **contigs**, saved in a FASTA file
- Like a jigsaw puzzle with no picture on the box



Schematic: reads shortened for the picture

Repeats: the "blue sky" problem

Genome



Short reads



which copy?

Long read



spans the repeat and both flanks

Read **shorter** than the repeat: its origin is ambiguous, so the assembly breaks

Read **longer** than the repeat: unique flanks anchor it, so the pieces connect

Contigs and scaffolds

Contigs (gap-free)



Scaffold (contigs joined by N gaps)



Contig: a gap-free sequence

Scaffold: contigs joined with N gaps (**N** = unknown base)

Goal for yeast: one contig per chromosome. ScRAP (Module 3) assembled the wolf (BAL_1a) as 16 nuclear contigs, 32 telomeres, 0 gaps. How close will Flye get? Lab 4

De novo vs reference-guided


	De novo	Reference-guided
Idea	Compare reads with each other; no guide	Place each read on a reference genome
Strength	Finds new sequence and structural variants	Fast; little memory
Weakness	Heavy on memory and CPU	Reference bias: misses what the reference lacks
Use when	No reference, or a very different sample	A close reference exists

Reference bias

Reference genome



Your sample



Reads from sequence the reference lacks are lost or misplaced

How different are the wolf (BAL_1a) and the corgi (S288C)? Modules 5 and 6 measure it


Your turn: which strategy?

- A new yeast species with no genome sequence yet
- 200 strains of S. cerevisiae, to find SNVs (single-base differences) in each
- One wild isolate where you suspect large rearrangements
- **Our pipeline:** de novo with Flye first, then the reference for evaluation and variant calling

Long vs short reads for assembly

	Short reads (Illumina)	Long reads (ONT)
Length	about 150-300 bp	thousands to 100,000+ bp (ours, Lab 3: mean about 19.5 kb)
Accuracy	about 99.9% (Q30) for most bases	Chemistry dependent
Repeats	Longer than the read: assembly breaks	Spanned by one read: contigs connect
Assembler	de Bruijn graph	overlap or repeat graph (Flye)
Needs	High coverage	Polishing

Hybrid assembly: long plus short



Illumina reads of this isolate exist (ENA ERR8603864); we do not use them in this session because of time. Polishing carries the accuracy.

How assemblers solve the puzzle


Greedy
merge the best overlap;
stalls on repeats

→

de Bruijn graph
reads cut into k-mers;
standard for short reads

→

Repeat graph (Flye)
long reads; repeats
resolved by spanning reads



Which exit goes with which entrance?

A read spanning A1, the repeat and B1 answers it

Tools you may meet with short reads: assembly

Tool	What it does
SPAdes	De Bruijn graph assembler for short reads (also hybrid)
MaSuRCA	Short-read or hybrid assembly

Not covered in this workshop

Flye: our assembler


- Built for long, noisy reads; builds a **repeat graph**
- **Read-type flag** must match the data
- --nano-raw: older ONT chemistries (up to 20% error)
- --nano-hq: ONT R10 reads, aka Q20 (<3% error)
- Ours: R9.4.1, basecaller unknown; about Q9 (Lab 3) fits --nano-raw

```
flye --nano-raw reads.fastq.gz
  --genome-size 12m
  --iterations 3
  --threads 8
  --out-dir assembly/
```

genome size: our estimate (about 12 Mb)
iterations: Flye's own polishing rounds
runs about 20 minutes
the log reports total length, contigs, N50 and coverage

Why polish?

- The assembler fixes the **order** of the pieces, but single bases can still be wrong
- **Polishing**: correct base errors in the assembly using the reads
- Typical ONT error (Module 3): a lost base in a run of one base (a homopolymer)
- Flye polishes internally first; then **medaka** polishes its consensus
- Polishing fixes bases, not structure



Illustrative example

Polishing: map and vote

Draft assembly (one wrong base)

A	C	G	T	**T**	A	C	G	A	T

Reads mapped to the draft

G	T	**C**	A	C	G	A
G	T	**C**	A	C	G	A
G	T	**C**	A	C	G	A
G	T	**C**	A	C	G	A
G	T	T	A	C	**A**	A

4 reads say C: vote C

Polished assembly

A	C	G	T	**C**	A	C	G	A	T

1 **Map** the reads to the draft

2 **Stack**: look at all bases at each position


3 **Vote**: the majority base replaces the draft base

One read's random error (here A) is outvoted

medaka: a neural-network polisher


- Made by Oxford Nanopore
- A **neural network** reads the stack of reads and decides the consensus
- It learned systematic nanopore error patterns, e.g. homopolymers
- Models are **chemistry-specific**: match flow cell and basecaller
- Takes about 6 minutes here

Our model name, decoded



Hybrid polishing with short reads

- Illumina reads of this isolate (ERR8603864, slide 13): we do not use them in this session because of time
- Very accurate (Q30): map them to the assembly and vote again
- Tools you may meet: Polypolish, pypolca (reimplements POLCA)
- Short reads map poorly inside repeats: less reliable there
- After the workshop: SPAdes vs Flye vs Flye + short-read polishing



dashed = not covered in this workshop


Judging an assembly: three lenses

Contiguity	Correctness	Completeness
How few, how long are the pieces? **contigs, N50, N90, L50** Tool: Flye log, QUAST	Does it agree with the reads and a reference? **misassemblies, mismatches** Tool: QUAST with a reference	Are the expected genes there? **C, S, D, F, M** Tool: BUSCO

Coverage underlies all three: how much evidence the assembly rests on. Never judge on one number


Coverage: how much evidence?

- **Coverage (depth)**: the average number of reads covering each base
- Total sequenced bases ÷ genome size
- Ours (Lab 3): **448 Mb ÷ 12.1 Mb = about 37×**
- Long-read assemblies: aim for 30-50×
- The Flye log reports the mean coverage



Schematic

N50: a worked example



Total = 74; half = 37

Add contigs from the longest until you pass 37

It takes 4 contigs: **L50 = 4**

The last one added has length 8: **N50 = 8**


QUAST: the corgi's card

- **QUAST**: Quality Assessment Tool for Genome Assemblies
- No reference needed for the basics
- Here: the S288C reference, our yardstick: 17 pieces
- GC 38%: the yeast value we compare against; 0 Ns: no gaps
- Report as text and HTML (copied to your laptop in the lab)

S288C reference	QUAST
# contigs	17
Largest contig	1,531,933
Total length	12,157,105
GC (%)	38.15
N50	924,431
N90	439,888
L50	6
# N's per 100 kbp	0.00

Ensembl R64-1-1; QUAST 5.2.0 without -r. N90: the same idea at 90% of the assembly, never larger than N50. Ours: Lab 4


QUAST with a reference



Misassembly: order or orientation of blocks disagrees with the reference

Also reported: mismatches and indels per 100 kbp, and the share of the reference covered


Misassembly or real variant?



XII: the rDNA break is a contig end (contiguity), an assembly limit, not a misassembly

A block missing against the reference: assembly error, or a real difference between the wolf and the corgi? QUAST cannot tell

BUSCO: are the genes there?



BUSCO: Benchmarking Universal Single-Copy Orthologs

Ortholog: the same gene in different species, split by speciation

These genes are expected once in nearly every yeast of the lineage

A biological check: does the assembly contain the genes a yeast needs?

Reading a BUSCO result

C:43.7%[S:43.4%,D:0.3%],F:2.6%,M:53.7%,n:2319


	Meaning	R7.3 chemistry	Genes
C	Complete (C = S + D)	43.7%	1,014
S	Complete, single-copy	43.4%	1,006
D	Complete, duplicated	0.3%	8
F	Fragmented	2.6%	60
M	Missing	53.7%	1,245
n	BUSCO genes searched		2,319

Note: Percentages are of the 2,319 BUSCO genes

What each BUSCO class tells you

- **S**: found once, as expected
- **D**: found more than once: real paralogs, or a region assembled twice
- **F**: only part of the gene: a gap or errors
- **M**: the gene is absent
- **Internal stop codons (E):** a complete gene with a stop inside. Legacy: 70 of 1,014 (6.9%)

A complete gene with an internal stop




stop codon inside the gene

A conserved gene should not have a stop in the middle. Which nanopore errors (Module 3) could cause this?

How many in ours, and why? Lab 4


A cautionary example: the legacy R7.3 assembly

Legacy R7.3 (ERR1539006): BUSCO genes




	Legacy R7.3	S288C (yardstick)	Ours: Lab 4
Contigs	431	17	?
Total length (bp)	10,073,637	12,157,105	?
N50 (bp)	28,584	924,431	?
BUSCO complete (C)	43.7%	not run	?
BUSCO missing (M)	53.7%	not run	?

Annotation: where, then what




Example: ADH1 in the corgi's annotation (S288C, Ensembl)



Finding genes: two approaches

	Ab initio	Evidence-based
Idea	Statistical models find gene patterns: start and stop codons, splice sites	Align known proteins and RNA-seq reads to the genome
Strength	Fast; needs no extra data	Real support; RNA-seq shows what is transcribed
Weakness	False positives if the model is poorly trained	Needs data; proteins find only known genes; RNA-seq misses unexpressed genes

funannotate: the pipeline



Yeast versus larger eukaryotes

	Yeast	Human (a larger eukaryote)
Genome size	about 12 Mb	about 3,100 Mb
Protein-coding genes	about 6,000	about 20,000
Genes with introns	few (about 5%)	most
Alternative splicing	rare	common
Gene prediction	ab initio works well on accurate sequence	needs RNA-seq evidence

A prediction is a hypothesis: compare it with what you expect for the species

How many gene models in ours, and how many are split into pieces? Lab 4

GFF3: the annotation file

seqid	source	type	start	end	score	strand	phase	attributes
XV	sgd	gene	159548	160594	.	-	.	ID=gene:YOL086C;Name=ADH1
XV	sgd	mRNA	159548	160594	.	-	.	ID=transcript:YOL086C_mRNA;Parent=gene:YOL086C
XV	sgd	exon	159548	160594	.	-	.	Parent=transcript:YOL086C_mRNA
XV	sgd	CDS	159548	160594	.	-	0	ID=CDS:YOL086C;Parent=transcript:YOL086C_mRNA


One feature per line; nine tab-separated columns

Parent= links exon and CDS to the mRNA, and the mRNA to the gene

A **gene model** = gene + mRNA + exons + CDS

Functional annotation: orthology

- Many predicted genes are "hypothetical proteins"
- **Homology**: shared ancestry; includes **paralogs**, which may have new functions
- **Orthology**: genes split by speciation usually keep their function
- eggNOG-mapper places each protein in an orthologous group and transfers annotations
- In: proteins (FASTA). Out: one table row per protein



A1 and B: orthologs (split by speciation)
A1 and A2: paralogs (split by duplication)

Reading the eggNOG table

Vocabulary	Describes	Example
GO	function, process, location	GO:0005355: glucose transmembrane transporter activity
KEGG	the gene family (KO) and its pathways	ADH1: KO K13953 in map00010 (Glycolysis / Gluconeogenesis)
EC	the reaction an enzyme catalyses	EC 1.1.1.1: alcohol dehydrogenase
COG	broad functional category (one letter)	K: transcription; P: inorganic ion transport

Example: ADH1 (S288C YOL086C): EC 1.1.1.1, KO K13953

Summary

- De novo assembly needs no reference; reference-guided is fast but biased
- Long reads span repeats; Flye assembles raw ONT reads; medaka polishes with the right model
- Judge an assembly by contiguity (N50), correctness (QUAST) and completeness (BUSCO)
- Lab 4 tells us how the wolf's genome scores on these three checks
- Structural annotation = where (GFF3); functional = what (eggNOG table)

Lab 4 preview: what you will do

1
Read the Flye log

2
Polish with medaka (sbatch)

3
Explore the supplied annotation

4
QUAST, then copy the report with scp

5
BUSCO and the legacy contrast

Bonus
Align the assembly to S288C; view in IGV

Questions are about reading results, not writing code

Coffee Break & Networking Session

Workshop Sponsors







