

# Canadian Bioinformatics Workshops

[www.bioinformatics.ca](http://www.bioinformatics.ca)  
[bioinformaticsdotca.github.io](https://bioinformaticsdotca.github.io)



## CC BY-SA 4.0 DEED

Attribution-ShareAlike 4.0 International

Canonical URL: <https://creativecommons.org/licenses/by-sa/4.0/>

[See the legal code](#)

### You are free to:

**Share** — copy and redistribute the material in any medium or format for any purpose, even commercially.

**Adapt** — remix, transform, and build upon the material for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the license terms.

### Under the following terms:

 **Attribution** — You must give [appropriate credit](#), provide a link to the license, and [indicate if changes were made](#). You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

 **ShareAlike** — If you remix, transform, or build upon the material, you must distribute your contributions under the [same license](#) as the original.

**No additional restrictions** — You may not apply legal terms or [technological measures](#) that legally restrict others from doing anything the license permits.



### Notices:

You do not have to comply with the license for elements of the material in the public domain or where your use is permitted by an applicable [exception or limitation](#).

No warranties are given. The license may not give you all of the permissions necessary for your intended use. For example, other rights such as [publicity, privacy, or moral rights](#) may limit how you use the material.

# RNA-Seq Module 2: Alignment QC


Malachi Griffith, Obi Griffith, Isabel Risch,  
Nicolas Ho, Melisa Acun, Varinder Verma, Mobin Khoramjoo  
RNA-seq Analysis 2025. July 7-9, 2025



# Learning objectives of module 3

- Visualization of RNA-seq alignments in IGV
- Alignment QC Assessment
- BAM read counting and determination of variant allele expression status

# Visualization of RNA-seq alignments in IGV browser




# Alternative viewers to IGV

- Alternative viewers to IGV
  - <http://www.biostars.org/p/12752/>
  - <http://www.biostars.org/p/71300/>
- Artemis, BamView, Chipster, gbrowse2, GenoViewer, MagicViewer, **Savant**, Tablet, tview

# Alignment QC Assessment

- 3' and 5' Bias
- Nucleotide Content
- Base/Read Quality
- PCR Artifact
- Sequencing Depth
- Base Distribution
- Insert Size Distribution

# Alignment QC: 3' & 5' Bias





# Alignment QC: Nucleotide Content

- **Random primers** are used to reverse transcribe RNA fragments into double-stranded complementary DNA (dscDNA)
- Causes certain patterns to be over represented at the beginning (5'end) of reads
- Deviation from expected  $A\%=C\%=G\%=T\%=25\%$

Journal List > Nucleic Acids Res > v.38(12); 2010 Jul > PMC2896536

## Nucleic Acids Research

Nucleic Acids Res. 2010 Jul; 38(12): e131.

Published online 2010 Apr 14. doi: [10.1093/nar/gkq224](https://doi.org/10.1093/nar/gkq224)


### Biases in Illumina transcriptome sequencing caused by random hexamer priming

Kasper D. Hansen,<sup>1,\*</sup> Steven E. Brenner,<sup>2</sup> and Sandrine Dudoit<sup>1,3</sup>

[Author information](#) ▶ [Article notes](#) ▶ [Copyright and License information](#) ▶

This article has been [cited by](#) other articles in PMC.


PI



<http://rseqc.sourceforge.net/>


# Alignment QC: Quality Distribution

- Phred quality score is widely used to characterize the quality of base-calling
- Phred quality score =  $-10 \times \log_{10}(P)$ , here  $P$  is probability that base-calling is wrong
- Phred score of 30 means there is 1/1000 chance that the base-calling is wrong
- The quality of the bases tend to drop at the end of the read, a pattern observed in sequencing by synthesis techniques



# Alignment QC: PCR Duplication


- Duplicate reads are reads that have the same start/end positions and same exact sequence
- In DNA-seq, reads/start point is used as a metric to assess PCR duplication rate
- In DNA-seq, duplicate reads are collapsed using tools such as picard
- How is RNA-seq different from DNA-seq?




<http://rseqc.sourceforge.net/>

# Alignment QC: Sequencing Depth

- **Have we sequenced deep enough?**
- In DNA-seq, we can determine this by looking at the average coverage over the sequenced region. Is it above a certain threshold?
- In RNA-seq, this is a challenge due to the variability in gene abundance
- Use splice junctions detection rate as a way to identify desired sequencing depth
- Check for saturation by resampling 5%, 10%, 15%, ..., 95% of total alignments from aligned file, and then detect splice junctions from each subset and compare to reference gene model.
- This method ensures that you have sufficient coverage to perform alternative splicing analyses





# Alignment QC: Base Distribution




- Your sequenced bases distribution will depend on the library preparation protocol selected

# Alignment QC: Insert Size



<http://thegenomefactory.blogspot.ca/2013/08/paired-end-read-confusion-library.html>


# Alignment QC: Insert Size



Consistent with library size selection?

<http://rseqc.sourceforge.net>

# BAM read counting and variant allele expression status



- A variant C->T is observed in 12 of 25 reads covering this position. Variant allele frequency (VAF)  $12/25 = 48\%$ .
- Both alleles appear to be expressed equally (not always the case) -> heterozygous, no allele specific expression
- How can we determine variant read counts, depth of coverage, and VAF without manually viewing in IGV?



# We are on a Coffee Break & Networking Session

Workshop Sponsors:



Canadian Centre for  
Computational  
Genomics



HPC4Health

