Table 1 Demographics of cohort, medications, and subjective measures of depression and sleep quality

		Groups				
	CTL, n=7	**PD^{dys-}, n=8**	**PD^{dys+}, n=10**	**Stat (df1, df2)**	**p-value**	**Effect-size**
Age, yr	56.14 ± 7.88	59.88 ±10.09	61.30 ± 9.91	ANOVA (2, 22)	0.544	0.054
Gender, % (F/M)	71/29	38/62	70/30	χ^2 (2, -)	0.289	0.288
Disease duration, yr	NA	5.75 ± 3.01	9.98 ± 5.73	t (16, -)	**0.047** [(a)]	**-0.974**
H&Y	NA	2.01 ± 0.11	2.20 ± 0.41	Welch t (9, -)	0.221	-0.002
MDS-UPDRS I	NA	3.57 ± 2.64	4.90 ± 3.63	MW	0.429	-0.039
MDS-UPDRS II	NA	4.75 ± 2.35	8.36 ± 4.45	t (15, -)	0.747	-0.243
MDS-UPDRS III	NA	19.12 ± 10.32	12.18 ± 6.35	t (16, -)	0.282	0.007
MDS-UPDRS IV	NA	2.00 ± 1.41	4.40 ± 3.44	t (10, -)	0.071	0.103
MDS-UPDRS Tot	NA	25.88 ± 13.45	27.60 ±11.72	t (16, -)	0.451	-0.020
AIMS	NA	NA	4.40 ± 3.44		NA	NA
LEDD, mg	NA	481.0 ± 175.82	854.64 ± 335.83	t (16, -)	**0.012** [(b)]	**-1.34**
ESS	5.14 ± 2.61	6.50 ± 4.14	7.70 ± 3.38	KW (2, -)	0.462	-0.021
PSQI	5.43 ± 3.15	5.62 ± 2.50	8.50 ± 3.17	KW (2, -)	0.082	0.214

ADV = advanced patients; AIMS = Abnormal Involuntary Movement Scale; CTL = control; DYS = dyskinetic patients; ESS = Epworth Sleepiness Scale; H&Y = Hoehn and Yahr staging; KW = Kruskal–Wallis; LEDD = L-dopa equivalent daily dose; MDS-UPDRS = Movement Disorder Society–sponsored revision of the Unified Parkinson Disease Rating Scale; MW = Mann–Whitney; NA = not applicable; PSQI = Pittsburgh Sleep Quality Index.

[(a)] $p = 0.036$ between PD^{dys-} and PD^{dys+}

[(b)] $p < 0.001$ between PD^{dys-} and PD^{dys+}