Table 1 Demographics of cohort, medications, and subjective measures of depression and sleep quality

	Groups					
	CTL, n=12	**EPD, n=12**	**ADV, n=13**	**DYS, n=11**	**p-value**	**Effect-size**
Age, yr	58.83 ± 9.63	64.58 ± 6.84	65.92 ± 9.79	60.91 ± 10.47	0.222	0.094
Gender, % (F/M)	42/58	25/75	8/92	55/45	-	-
Disease duration, yr	NA	3.42 ± 2.19 [(a)]	7.54 ± 3.67	8.82 ± 2.96	**3.49e-04**	**0.383**
H&Y	NA	1.25 ± 0.45	1.85 ± 0.38	2.09 ± 0.30	-	-
MDS-UPDRS I	NA	4.32 ± 4.37	6.08 ± 3.82	5.36 ± 3.17	0.700	-0.039
MDS-UPDRS II	NA	5.50 ± 3.78	7.00 ± 3.72	7.00 ± 4.98	0.747	-0.041
MDS-UPDRS III	NA	12.00 ± 8.09	14.46 ± 8.46	11.45 ± 9.06	0.282	0.007
MDS-UPDRS IV	NA	NA	0.69 ± 1.03	2.64 ± 2.80	0.071	0.103
MDS-UPDRS Tot	NA	21.75 ± 14.23	28.23 ± 13.23	26.45 ± 16.05	0.451	-0.020
UDysRS IV	NA	NA	NA	5.80 ± 4.10	NA	NA
UDysRS Tot	NA	NA	NA	26.40 ± 18.28	NA	NA
AIMS	NA	NA	NA	7.64 ± 5.07	NA	NA
LEDD, mg	NA	297.1 ± 169.1 [(b)]	863.0 ± 367.8	1078.7 ± 545.4	**7.64e-05**	**0.323**
HADS-A	3.75 ± 2.09	3.83 ± 3.33	3.85 ± 4.12	3.36 ± 2.38	0.907	-0.045
HADS-D	2.17 ± 2.37	2.50 ± 1.57	3.08 ± 3.15	2.64 ± 2.91	0.837	-0.043
ESS	6.25 ± 3.28	5.67 ± 4.08	7.38 ± 4.44	6.27 ± 1.74	0.509	-0.015
PSQI	5.83 ± 2.89	5.00 ± 3.22	7.38 ± 2.10	6.36 ± 4.18	0.298	0.079

ADV = advanced patients; AIMS = Abnormal Involuntary Movement Scale; CTL= control; DYS = dyskinetic patients; EPD= early-stage patients; ESS = Epworth Sleepiness Scale; Hospital Anxiety and Depression Scale – Anxiety (HADS-A); Hospital Anxiety and Depression Scale – Depression (HADS-D); H&Y = Hoehn and Yahr staging; LEDD = L-dopa equivalent daily dose; MDS-UPDRS = Movement Disorder Society–sponsored revision of the Unified Parkinson Disease Rating Scale; NA = not applicable; PSQI = Pittsburgh Sleep Quality Index; UDysRS = Unified Dyskinesia Rating Scale.

[(a)] $p = 0.0049$ versus ADV, and $p < 0.001$ versus DYS, Tukey post hoc test.

[(b)] $p < 0.001$ versus ADV and DYS, Games-Howell post hoc test.