

# A few Bayesian lectures for the Uninitiated



## Part 5: a mixed-effect model example

Carsten F. Dormann

Biometry & Environmental System Analysis, University of Freiburg

2. April 2022

# Mixed effect model

- ▶ Mixed effect model  $\in$  hierarchical models
- ▶ hierarchy: some parameters describe the response  $y$ , others internal, higher-level *unobservables*
- ▶ PISA school study: all measurements are at the level of the individual pupil, but there are parameters for schools, states and nations in the model
- ▶ mixed effect: subjects are random draws from the population of interest  $\longrightarrow$  *subject-level* effects, and *population-level* effects

# Mixed effect model and its hierarchies

$$y = X\beta + Zu + \varepsilon \text{ (random intercept model)}$$

# Mixed effect model and its hierarchies

$$y = X\beta + Zu + \varepsilon \text{ (random intercept model)}$$

$$y_i = X_i\beta + Zu_i + \varepsilon_i$$

# Mixed effect model and its hierarchies

$$y = X\beta + Zu + \varepsilon \text{ (random intercept model)}$$

$$y_i = X_i\beta + Zu_i + \varepsilon_i$$

$$y_i = X_i\beta + Zu_{j_i} + \varepsilon_i !!$$

# Mixed effect model and its hierarchies

$$y = X\beta + Zu + \varepsilon \text{ (random intercept model)}$$

$$y_i = X_i\beta + Zu_i + \varepsilon_i$$

$$y_i = X_i\beta + Zu_{j_i} + \varepsilon_i !!$$

In words: the  $i$ th data point belongs to a specific  $j$ th random effect.

# Mixed effect model and its hierarchies

$$y = X\beta + Zu + \varepsilon \text{ (random intercept model)}$$

$$y_i = X_i\beta + Zu_i + \varepsilon_i$$

$$y_i = X_i\beta + Zu_{j_i} + \varepsilon_i !!$$

In words: the  $i$ th data point belongs to a specific  $j$ th random effect.

Example: 10 subject were measured 20 times.  $j \in \{1, \dots, 10\}$ , while  $i \in \{1, \dots, 200\}$ .

$Z$  is in this case a dummy matrix with 10 columns, and each column has one estimate for its subject  $u_j$ .

## One final step

The likelihood of  $y_i$  is based on the estimate of population-level  $\beta$ , but each subject gets an additional “offset”  $u$ .



## One final step

The likelihood of  $y_i$  is based on the estimate of population-level  $\beta$ , but each subject gets an additional “offset”  $u$ .

These “offsets”  $u$  are computed relative to their mean. That mean is the actual regression intercept  $\beta_0$ .

## One final step

The likelihood of  $y_i$  is based on the estimate of population-level  $\beta$ , but each subject gets an additional “offset”  $u$ .

These “offsets”  $u$  are computed relative to their mean. That mean is the actual regression intercept  $\beta_0$ .

That means, we use the individual-level intercepts  $u_j$  to estimate the population-level intercept  $\bar{u}$ .

## One final step

The likelihood of  $y_i$  is based on the estimate of population-level  $\beta$ , but each subject gets an additional “offset”  $u$ .

These “offsets”  $u$  are computed relative to their mean. That mean is the actual regression intercept  $\beta_0$ .

That means, we use the individual-level intercepts  $u_j$  to estimate the population-level intercept  $\bar{u}$ .

*That* is the parameter at the other hierarchical level!

## One final step

The likelihood of  $y_i$  is based on the estimate of population-level  $\beta$ , but each subject gets an additional “offset”  $u$ .

These “offsets”  $u$  are computed relative to their mean. That mean is the actual regression intercept  $\beta_0$ .

That means, we use the individual-level intercepts  $u_j$  to estimate the population-level intercept  $\bar{u}$ .

*That* is the parameter at the other hierarchical level!

Let's write this as equations:

# A random intercept mixed effect model

$$y_i \sim N(\mu = \hat{y}_i, \sigma = \hat{s})$$

# A random intercept mixed effect model

$$y_i \sim N(\mu = \hat{y}_i, \sigma = \hat{s})$$

$$\hat{y}_i = X_i\beta + Z_iu_{j_i}$$

# A random intercept mixed effect model

$$y_i \sim N(\mu = \hat{y}_i, \sigma = \hat{s})$$

$$\hat{y}_i = X_i\beta + Z_iu_{j_i}$$

$$u_j \sim N(\mu = \hat{u}, \sigma = \hat{s}_u)$$

# A random intercept mixed effect model

$$y_i \sim N(\mu = \hat{y}_i, \sigma = \hat{s})$$

$$\hat{y}_i = X_i\beta + Z_i u_{j_i}$$

$$u_j \sim N(\mu = \hat{u}, \sigma = \hat{s}_u)$$

$$\hat{u} = c$$



# A random intercept mixed effect model

$$y_i \sim N(\mu = \hat{y}_i, \sigma = \hat{s})$$

$$\hat{y}_i = X_i\beta + Z_i u_{j_i}$$

$$u_j \sim N(\mu = \hat{u}, \sigma = \hat{s}_u)$$

$$\hat{u} = c$$

Final touch: Bayesian here only means that you have to have a prior for the  $\beta$ s, the  $\sigma$ s, the  $z$ s and indeed for  $c$ .

# In JAGS

```
for (i in 1:N){  
  Reaction[i] ~ dnorm(muHat[i], tau)  
  # random intercept model:  
  muHat[i] <- beta0[subjectNr[i]] + beta1 * Days[i]  
}  
# priors  
for (j in 1:k){  
  beta0[j] ~ dnorm(beta00, tau.random) # random intercept  
}  
beta00 ~ dnorm(300, 0.001)           # grand mean  
tau ~ dgamma(0.001, 0.001)  
tau.random ~ dgamma(0.001, 0.001)
```

# Will this never end?

**Next time:**  
**State-space models**