

A few Bayesian lectures for the Uninitiated



Part 5: a mixed-effect model example

Carsten F. Dormann

Biometry & Environmental System Analysis, University of Freiburg

20. April 2022

Mixed effect model

- ▶ Mixed effect model \in hierarchical models
- ▶ hierarchy: some parameters describe the response y , others internal, higher-level *unobservables*
- ▶ PISA school study: all measurements are at the level of the individual pupil, but there are parameters for schools, states and nations in the model
- ▶ mixed effect: subjects are random draws from the population of interest \longrightarrow *subject-level* effects, and *population-level* effects

Mixed effect model and its hierarchies

$$y = X\beta + Zu + \varepsilon \text{ (random intercept model)}$$

Mixed effect model and its hierarchies

$$y = X\beta + Zu + \varepsilon \text{ (random intercept model)}$$

$$y_i = X_i\beta + Zu_i + \varepsilon_i$$

Mixed effect model and its hierarchies

$$y = X\beta + Zu + \varepsilon \text{ (random intercept model)}$$

$$y_i = X_i\beta + Zu_i + \varepsilon_i$$

$$y_i = X_i\beta + Zu_{j_i} + \varepsilon_i !!$$

Mixed effect model and its hierarchies

$$y = X\beta + Zu + \varepsilon \text{ (random intercept model)}$$

$$y_i = X_i\beta + Zu_i + \varepsilon_i$$

$$y_i = X_i\beta + Zu_{j_i} + \varepsilon_i !!$$

In words: the i th data point belongs to a specific j th random effect.

Mixed effect model and its hierarchies

$$y = X\beta + Zu + \varepsilon \text{ (random intercept model)}$$

$$y_i = X_i\beta + Zu_i + \varepsilon_i$$

$$y_i = X_i\beta + Zu_{j_i} + \varepsilon_i !!$$

In words: the i th data point belongs to a specific j th random effect.

Example: 10 subject were measured 20 times. $j \in \{1, \dots, 10\}$, while $i \in \{1, \dots, 200\}$.

Z is in this case a dummy matrix with 10 columns, and each column has one estimate for its subject u_j .

One final step

The likelihood of y_i is based on the estimate of population-level β , but each subject gets an additional “offset” u .

One final step

The likelihood of y_i is based on the estimate of population-level β , but each subject gets an additional “offset” u .

These “offsets” u are computed relative to their mean. That mean is the actual regression intercept β_0 .

One final step

The likelihood of y_i is based on the estimate of population-level β , but each subject gets an additional “offset” u .

These “offsets” u are computed relative to their mean. That mean is the actual regression intercept β_0 .

That means, we use the individual-level intercepts u_j to estimate the population-level intercept \bar{u} .

One final step

The likelihood of y_i is based on the estimate of population-level β , but each subject gets an additional “offset” u .

These “offsets” u are computed relative to their mean. That mean is the actual regression intercept β_0 .

That means, we use the individual-level intercepts u_j to estimate the population-level intercept \bar{u} .

That is the parameter at the other hierarchical level!

One final step

The likelihood of y_i is based on the estimate of population-level β , but each subject gets an additional “offset” u .

These “offsets” u are computed relative to their mean. That mean is the actual regression intercept β_0 .

That means, we use the individual-level intercepts u_j to estimate the population-level intercept \bar{u} .

That is the parameter at the other hierarchical level!

Let's write this as equations:

A random intercept mixed effect model

$$y_i \sim N(\mu = \hat{y}_i, \sigma = \hat{s})$$

A random intercept mixed effect model

$$y_i \sim N(\mu = \hat{y}_i, \sigma = \hat{s})$$
$$\hat{y}_i = X_i\beta + Z_iu_{j_i}$$

A random intercept mixed effect model

$$y_i \sim N(\mu = \hat{y}_i, \sigma = \hat{s})$$

$$\hat{y}_i = X_i\beta + Z_iu_{j_i}$$

$$u_j \sim N(\mu = \hat{u}, \sigma = \hat{s}_u)$$

A random intercept mixed effect model

$$y_i \sim N(\mu = \hat{y}_i, \sigma = \hat{s})$$

$$\hat{y}_i = X_i\beta + Z_iu_{j_i}$$

$$u_j \sim N(\mu = \hat{u}, \sigma = \hat{s}_u)$$

$$\hat{u} = c$$

A random intercept mixed effect model

$$y_i \sim N(\mu = \hat{y}_i, \sigma = \hat{s})$$

$$\hat{y}_i = X_i\beta + Z_i u_{j_i}$$

$$u_j \sim N(\mu = \hat{u}, \sigma = \hat{s}_u)$$

$$\hat{u} = c$$

Final touch: Bayesian here only means that you have to have a prior for the β s, the σ s, the z s and indeed for c .

In JAGS

```
for (i in 1:N){  
  Reaction[i] ~ dnorm(muHat[i], tau)  
  # random intercept model:  
  muHat[i] <- beta0[subjectNr[i]] + beta1 * Days[i]  
}  
# priors  
for (j in 1:k){  
  beta0[j] ~ dnorm(beta00, tau.random) # random intercept  
}  
beta00 ~ dnorm(300, 0.001)           # grand mean  
tau ~ dgamma(0.001, 0.001)  
tau.random ~ dgamma(0.001, 0.001)
```

Easy!

Will this never end?

Next time:
State-space models

Useful links:

[https://www.r-bloggers.com/2022/04/
bayesian-analyses-made-easy-glmms-in-r-package-brms/](https://www.r-bloggers.com/2022/04/bayesian-analyses-made-easy-glmms-in-r-package-brms/)