

Efficacy and Safety of First-line Avelumab Treatment in Patients With Stage IV Metastatic Merkel Cell Carcinoma

A Preplanned Interim Analysis of a Clinical Trial

Sandra P. D'Angelo, MD; Jeffery Russell, MD, PhD; Céleste Lebbé, MD, PhD; Bartosz Chmielowski, MD, PhD; Thilo Gambichler, MD; Jean-Jacques Grob, MD; Felix Kiecker, MD; Guilherme Rabinowits, MD; Patrick Terheyden, MD; Isabella Zwiener, PhD; Marcis Bajars, MD; Meliessa Hennessy, MPH; Howard L. Kaufman, MD

 [Supplemental content](#)

IMPORTANCE Merkel cell carcinoma (MCC) is an aggressive skin cancer that is associated with poor survival outcomes in patients with distant metastatic disease. Results of part A of the JAVELIN Merkel 200 trial (avelumab in patients with Merkel cell carcinoma) showed that avelumab, an anti-programmed cell death ligand 1 (PD-L1) antibody, demonstrated efficacy in second-line or later treatment of patients with metastatic MCC (mMCC).

OBJECTIVE To evaluate the efficacy and safety of avelumab as first-line treatment for patients with distant mMCC.

DESIGN, SETTING, AND PARTICIPANTS JAVELIN Merkel 200 part B is an international, multicenter, single-arm, open-label clinical trial of first-line avelumab monotherapy. Eligible patients were adults with mMCC who had not received prior systemic treatment for metastatic disease. Patients were not selected for PD-L1 expression or Merkel cell polyomavirus status. Data were collected from April 15, 2016, to March 24, 2017, and enrollment is ongoing.

INTERVENTIONS Patients received avelumab, 10 mg/kg, by 1-hour intravenous infusion every 2 weeks until confirmed disease progression, unacceptable toxic effects, or withdrawal occurred.

MAIN OUTCOMES AND MEASURES Tumor status was assessed every 6 weeks and evaluated by independent review committee per Response Evaluation Criteria in Solid Tumors version 1.1. The primary end point was durable response, defined as an objective response with a duration of at least 6 months. Secondary end points include best overall response, duration of response, progression-free survival, safety, and tolerability.

RESULTS As of March 24, 2017, 39 patients were enrolled (30 men and 9 women; median age, 75 years [range, 47-88 years]), with a median follow-up of 5.1 months (range, 0.3-11.3 months). In a preplanned analysis, efficacy was assessed in 29 patients with at least 3 months of follow-up; the confirmed objective response rate was 62.1% (95% CI, 42.3%-79.3%), with 14 of 18 responses (77.8%) ongoing at the time of analysis. In responding patients, the estimated proportion with duration of response of at least 3 months was 93% (95% CI, 61%-99%); duration of response of at least 6 months, 83% (95% CI, 46%-96%). First-line avelumab treatment was generally well tolerated, and no treatment-related deaths or grade 4 adverse events occurred.

CONCLUSIONS AND RELEVANCE High rates of response to first-line avelumab therapy in patients with distant mMCC build on previously reported antitumor activity after second-line or later treatment, and maturing progression-free survival data suggest that responses are durable. These data further support avelumab's approval in the United States and European Union and use as a standard-of-care treatment for mMCC.

TRIAL REGISTRATION clinicaltrials.gov Identifier: [NCT02155647](#)

JAMA Oncol. 2018;4(9):e180077. doi:[10.1001/jamaoncol.2018.0077](#)
Published online March 22, 2018.

Author Affiliations: Author affiliations are listed at the end of this article.

Corresponding Author: Sandra P. D'Angelo, MD, Weill Cornell Medical College and Memorial Sloan Kettering Cancer Center, 300 E 66th St, 12th Floor, New York, NY 10065 (dangelos@mskcc.org).

Metastatic Merkel cell carcinoma (mMCC) is a rare, aggressive skin cancer with poor survival.¹ Merkel cell carcinoma (MCC) is associated with Merkel cell polyomavirus infection, UV light exposure, and advanced age; additional risk factors include fair skin and male sex.^{1,2} Despite MCC being a chemosensitive disease, durable responses are rare.³ In retrospective studies of first-line chemotherapy in patients with mMCC, median progression-free survival (PFS) was 3.1 to 4.6 months,^{4,5} highlighting the need for improved treatment options. Programmed cell death ligand 1 (PD-L1) is expressed on MCC tumor cells and in the MCC tumor microenvironment, indicating a potential role for immune checkpoint inhibitors such as anti-PD-L1 and anti-programmed cell death 1 (PD-1) antibodies.⁶⁻⁸ Trials involving anti-PD-1/PD-L1 antibodies have shown clinical activity and durable responses in patients with advanced MCC.⁹⁻¹¹ Based on findings from the phase 2 trial of avelumab, an IgG1 anti-PD-L1 monoclonal antibody, in patients with MCC (JAVELIN Merkel 200), avelumab became the first treatment for patients with mMCC to receive approval in the United States, the European Union, and Japan.¹¹⁻¹⁵ Herein, we report data from an interim analysis of avelumab as first-line treatment in patients with mMCC.

Methods

In part B of the ongoing JAVELIN Merkel 200 trial (start date, April 15, 2016), eligible patients had stage IV mMCC and had not received prior systemic therapy for metastatic disease. Patients with autoimmune conditions were excluded. The primary end point was durable response, defined as an objective response with a duration of at least 6 months according to Response Evaluation Criteria in Solid Tumors version 1.1, per independent review. Secondary end points included best overall response, duration of response, PFS, and safety. PD-L1 expression and viral status will be investigated in a future analysis. Time-to-event end points were analyzed by Kaplan-Meier methods; medians were calculated with corresponding CIs using the Brookmeyer-Crowley method. Additional eligibility criteria, procedures for statistical analysis, and study design information for JAVELIN Merkel 200 have been previously reported.¹¹ Patients provided written informed consent to enroll in the protocol with institutional review board approval from the participating centers (eTable 1 in the [Supplement](#)) and in accordance with international good clinical practice standards.

Results

From April 15, 2016, to March 24, 2017, 39 of 112 patients initiated avelumab treatment (**Figure 1**), with a median follow-up of 5.1 months (range, 0.3-11.3 months). Thirty patients were men, and 9 were women, and median age was 75 years (range, 47-88 years); additional demographics are described in eTable 2 in the [Supplement](#). Median duration of treatment was 12 weeks (range, 2.0-49.9 weeks). Treatment was ongoing in 24 patients (61.5%), and 15 (38.5%) discontinued


Key Points

Question What is the efficacy of first-line avelumab treatment in patients with distant metastatic Merkel cell carcinoma?

Findings In a preplanned interim analysis of a phase 2 study in 39 patients with metastatic Merkel cell carcinoma, avelumab treatment resulted in a confirmed objective response rate of 62.1% (95% CI, 42.3%-79.3%); 16 of 18 responses (88.9%) occurred by the first planned postbaseline assessment. Grade 3 treatment-related adverse events occurred in 8 of 39 patients (20.5%), with no treatment-related deaths.

Meaning First-line avelumab monotherapy in patients with metastatic Merkel cell carcinoma was associated with early responses, high response rates, and a manageable safety profile.

Figure 1. Flowchart of Study Population



treatment owing to disease progression (7 [17.9%]), adverse events (6 [15.4%]), or death (2 [5.1%]). In this prespecified interim analysis, safety was assessed in 39 patients who received at least 1 dose of avelumab, and efficacy was evaluated in 29 patients with at least 3 months of follow-up. Efficacy was also assessed in a subset of 14 patients with at least 6 months of follow-up.

Of the 29 patients with at least 3 months of follow-up, 4 (13.8%) had a complete response and 14 (48.3%) had a partial response, resulting in a confirmed objective response rate of 62.1% (95% CI, 42.3%-79.3%) (**Table**); 3 additional patients (10.3%) had stable disease. Median time to response was 6.1 weeks (range, 5-17 weeks), with 16 of 18 responses (88.9%) observed at the first scheduled postbaseline assessment approximately 6 weeks from the start of treatment (**Figure 2A**). Among the 14 patients with at least 6 months of follow-up, the confirmed objective response rate was 71.4% (95% CI, 41.9%-91.6%). Four patients (28.6%) had a complete response, and 6 (42.9%) had a partial response; an additional patient (7.1%) had stable disease.

Responses were ongoing at the time of analysis in 14 of 18 responders (77.8%), and the median duration of response was not estimable (range, 1.2-8.3 months). Based on Kaplan-Meier estimates, the proportion of responses with a duration

Table. Outcomes by RECIST Version 1.1, per Independent Review Committee Assessment

Outcome	Patient Follow-up Group	
	≥3 mo	≥6 mo
Response^a		
Confirmed ORR, % (95% CI)	62.1 (42.3-79.3)	71.4 (41.9-91.6)
Confirmed BOR, No. (%)		
Complete response	4 (13.8)	4 (28.6)
Partial response	14 (48.3)	6 (42.9)
Stable disease	3 (10.3)	1 (7.1)
Progressive disease	7 (24.1)	2 (14.3)
Nonevaluable ^b	1 (3.4)	1 (7.1)
Response durability^c		
Median DOR (95% CI), mo	NE (4.0 to NE)	NE (4.0 to NE)
Proportion of responses with duration ≥3 mo, % (95% CI)	93 (61-99)	100 (NE)
Proportion of responses with duration ≥6 mo, % (95% CI)	83 (46-96)	89 (43-98)

Abbreviations: BOR, best overall response; DOR, duration of response; NE, not estimable; ORR, objective response rate; RECIST, Response Evaluation Criteria in Solid Tumors.

^a Includes 29 patients with at least 3 and 14 with at least 6 months of follow-up.

^b Patient died before tumor assessment due to an adverse event unrelated to treatment with avelumab.


^c Includes 18 patients with at least 3 and 10 with at least 6 months of follow-up.

of at least 3 months was 93% (95% CI, 61%-99%); those with a response duration of at least 6 months, 83% (95% CI, 49%-96%). In this still-maturing cohort, the 3-month PFS rate was 67% (95% CI, 48%-80%), and median PFS was 9.1 months (range, 1.9 to not estimable) (Table). Twenty-one of 30 patients (70%) with a follow-up tumor assessment available were reported to have at least 30% reduction in target lesions from baseline (Figure 2B-C).

Among 39 patients evaluable for safety, 28 (71.8%) had a treatment-related adverse event (TRAE), and 8 patients (20.5%) had a grade 3 TRAE (eTable 3 in the [Supplement](#)). No grade 4 TRAEs or treatment-related deaths occurred. Treatment-related infusion-related reactions occurred in 9 patients (23.1%) and were grade 3 in 1 patient (2.6%). Grade 1 immune-related adverse events occurred in 6 patients (15.4%) (eTable 4 in the [Supplement](#)). Avelumab therapy was permanently discontinued because of a TRAE in 1 patient each with cholangitis, elevated aspartate aminotransferase and alanine aminotransferase levels, paraneoplastic syndrome and gait disturbance, and paraneoplastic encephalomyelitis and polyneuropathy and in 2 patients with infusion-related reaction (6 patients [15.4%]). Of these 6 patients, 1 had a complete response ongoing at the time of analysis, 1 had a partial response assessed before the start of subsequent anticancer therapy, and 1 had stable disease before the start of subsequent anticancer therapy.

Discussion

In this prespecified interim analysis of patients with mMCC, first-line avelumab treatment was associated with early and durable responses and a manageable safety profile. The confirmed ob-

Figure 2. Clinical Activity of Avelumab

A, Time to and duration of response and duration of treatment were measured in 18 patients with a confirmed response within the analysis set of patients with at least 3 months of follow-up. B and C, Change in target lesion diameters and percentage of change in target lesion sum diameters from baseline in individual patients were measured in 30 patients with a follow-up tumor assessment available. Parallel broken lines represent Response Evaluation Criteria in Solid Tumors values for progressive disease ($\geq 20\%$ increase in the sum of diameters of target lesions) and partial response ($\geq 30\%$ decrease in the sum of diameters of target lesions).

jective response rate is consistent with results of previous studies of immune checkpoint inhibitors for first-line treatment of

MCC.^{9,10} To our knowledge, this study is the first to investigate immune checkpoint inhibitors exclusively in patients with stage IV MCC. Two anti-PD-1 antibodies have been investigated for the first-line treatment of patients with advanced MCC, including nivolumab in patients with stages II to IV MCC and pembrolizumab in patients with stages IIIB to IV MCC.^{9,10}

Before the approval of avelumab, chemotherapy was commonly used to treat mMCC; it is now recommended for patients who are not candidates for immunotherapy.² In a systematic literature review, first-line chemotherapy produced high initial response rates (range, 53%-61%); however, response durability was short (range, 2.8-8.0 months), and treatment was associated with considerable toxic effects.³ We report promising evidence of response durability among patients with an objective response and at least 3 months of follow-up, given that 83% of patients are estimated to have responses lasting at least 6 months. Our median PFS of 9.1 months for patients receiving first-line treatment exceeds that previously reported with chemotherapy.^{4,5} Our safety analysis showed that avelumab was generally tolerable. These findings suggest overall improvement of outcomes with immune checkpoint inhibitors vs chemotherapy in patients with mMCC.

Our results corroborate prior evidence from part A of the JAVELIN Merkel 200 study,¹¹ which showed that previously treated patients with fewer prior lines of treatment experienced more favorable outcomes. JAVELIN Merkel 200 is the

largest prospective clinical trial performed in stage IV distant mMCC to date and is actively enrolling participants in part B of the study. Findings from this trial led to approval of avelumab for the treatment of mMCC in the United States, the European Union, and Japan.¹²⁻¹⁴

Strengths and Limitations

Analysis of PFS in the present study is based on limited follow-up in some patients and may evolve as the minimum follow-up extends; longer-term follow-up will also permit meaningful analysis of overall survival. Because data with extended follow-up have not been published for nivolumab and pembrolizumab, comparisons of long-term survival outcomes, TRAEs, and economic impact with other immune checkpoint inhibitors for first-line treatment were not possible.^{9,10} Future analysis of first-line avelumab treatment will investigate whether biomarkers, such as PD-L1 expression or viral status, correlate with response to treatment.

Conclusions

First-line avelumab monotherapy had antitumor activity and a manageable safety profile in patients with mMCC. Maturing PFS data suggest durable responses, supporting avelumab as a standard of care in patients with mMCC.

ARTICLE INFORMATION

Accepted for Publication: December 26, 2017.

Published Online: March 22, 2018.

doi:10.1001/jamaoncol.2018.0077

Open Access: This article is published under the [JN-OA license](#) and is free to read on the day of publication.

Author Affiliations: Department of Medicine, Weill Cornell Medical College, New York, New York (D'Angelo); Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, New York (D'Angelo); H. Lee Moffitt Cancer Center, Tampa, Florida (Russell); currently affiliated with Immunocore, Ltd, Conshohocken, Pennsylvania (Russell); Assistance Publique-Hôpitaux de Paris Dermatology and Centre d'Investigation Clinique, University Paris Diderot Institut National de la Santé et de la Recherche Médicale U976, Saint Louis Hospital, Paris, France (Lebbé); Department of Medicine, UCLA (University of California, Los Angeles) Medical Center (Chmielowski); Skin Cancer Center, Department of Dermatology, Ruhr-University Bochum, Bochum, Germany (Gambichler); Department of Dermatology, Venerology, and Cutaneous Oncology, Aix-Marseille University, Assistance Publique-Hôpitaux de Marseille, Timone Hospital, Marseille, France (Grob); Department of Dermatology, Venereology, and Allergology, Charité Universitätsmedizin Berlin, Edmund-Lesser-Haus, Berlin, Germany (Kiecker); Dana-Farber Cancer Institute, Boston, Massachusetts (Rabinowits); Department of Dermatology, Allergology, and Venereology, University of Lübeck, Lübeck, Germany (Terheyden); Merck KGaA, Darmstadt, Germany (Zwiener); EMD Serono, Billerica, Massachusetts (Bajars, Hennessy); Rutgers Cancer Institute of New

Jersey, New Brunswick (Kaufman); currently affiliated with Replimune Inc, Woburn, Massachusetts (Kaufman).

Author Contributions: Dr D'Angelo had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: D'Angelo, Bajars, Kaufman.

Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: D'Angelo, Kaufman.

Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: Terheyden, Zwiener.

Administrative, technical, or material support: Chmielowski, Terheyden, Kaufman.

Study supervision: D'Angelo, Lebbé, Chmielowski, Gambichler, Bajars, Hennessy, Kaufman.

Conflict of Interest Disclosures: Dr D'Angelo reports consulting and advisory roles for EMD Serono, Pfizer, Nektar Therapeutics, and Immune Design. Dr Russell reports consulting and advisory roles for EMD Serono and employment by Immunocore Ltd. Dr Lebbé reports consulting and advisory roles and speaker honoraria for Roche, BMS, Novartis, MSD, and Amgen; research grants from Roche and BMS; and travel accommodations and meeting expenses from Roche, BMS, Novartis, and Amgen. Dr Chmielowski reports consulting and advisory roles for EMD Merck Serono. Dr Gambichler reports consulting and advisory roles for Merck, speaker honoraria for NeraCare, and congress travel support from MSD and Novartis. Dr Grob reports consulting and advisory roles for Roche, MSD, Novartis, Amgen, BMS, Merck & Co, Pfizer, and Pierre Fabre and speaker honoraria for Novartis, Amgen, and BMS. Dr Kiecker reports

consulting and advisory roles for Merck Serono, MSD, BMS, Roche, Novartis, Incyte, and Pfizer; a research grant from Novartis; and speaker honoraria from Merck Serono, MSD, BMS, Roche, Novartis, and Pfizer. Dr Rabinowits reports consulting and advisory roles for EMD Serono and Pfizer and research support (to the institution) from EMD Serono, Exelixis, and Millennium. Dr Terheyden reports consulting and advisory roles for BMS, Merck & Co, Novartis, and Roche and speaker honoraria from BMS, Novartis, and Roche. Dr Zwiener reports employment by Merck KGaA. Dr Bajars and Ms Hennessy report employment by EMD Serono (a business of Merck KGaA). Dr Kaufman reports consulting and advisory roles for Amgen, Celldex, EMD Serono, Merck, Prometheus, and Turnstone Biologics; research grants from Amgen and Merck; and speaker honoraria from Merck. No other disclosures were reported.

Funding/Support: This study was supported by Merck KGaA and is part of an alliance between Merck KGaA and Pfizer Inc.

Role of the Funder/Sponsor: In collaboration with academic authors, representatives of the sponsor participated in design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, and approval of the manuscript; and decision to submit the manuscript for publication.

Additional Contributions: Medical writing support was provided by ClinicalThinking, Inc. The authors thank the patients and their families and the investigators, coinvestigators, and study teams at each of the participating centers and at Merck KGaA and EMD Serono.

REFERENCES

- Schadendorf D, Lebbé C, Zur Hausen A, et al. Merkel cell carcinoma: epidemiology, prognosis, therapy and unmet medical needs. *Eur J Cancer*. 2017;71:53-69.
- National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Merkel Cell Carcinoma, version 1. https://www.nccn.org/store/login/login.aspx?ReturnURL=https://www.nccn.org/professionals/physician_gls/pdf/mcc.pdf. Updated 2017. Accessed October 19, 2017.
- Nghiem P, Kaufman HL, Bharmal M, Mahnke L, Phatak H, Becker JC. Systematic literature review of efficacy, safety and tolerability outcomes of chemotherapy regimens in patients with metastatic Merkel cell carcinoma. *Future Oncol*. 2017;13(14):1263-1279.
- Cowey CL, Mahnke L, Espirito J, Helwig C, Oksen D, Bharmal M. Real-world treatment outcomes in patients with metastatic Merkel cell carcinoma treated with chemotherapy in the USA. *Future Oncol*. 2017;13(19):1699-1710.
- Iyer JG, Blom A, Doumani R, et al. Response rates and durability of chemotherapy among 62 patients with metastatic Merkel cell carcinoma. *Cancer Med*. 2016;5(9):2294-2301.
- Lipson EJ, Vincent JG, Loyo M, et al. PD-L1 expression in the Merkel cell carcinoma microenvironment: association with inflammation, Merkel cell polyomavirus and overall survival. *Cancer Immunol Res*. 2013;1(1):54-63.
- Schadendorf D, Nghiem P, Bhatia S, et al. Immune evasion mechanisms and immune checkpoint inhibition in advanced Merkel cell carcinoma. *Oncoimmunology*. 2017;6(10):e1338237.
- Terheyden P, Becker JC. New developments in the biology and the treatment of metastatic Merkel cell carcinoma. *Curr Opin Oncol*. 2017;29(3):221-226.
- Nghiem PT, Bhatia S, Lipson EJ, et al. PD-1 blockade with pembrolizumab in advanced Merkel-cell carcinoma. *N Engl J Med*. 2016;374(26):2542-2552.
- Topalian SL, Bhatia S, Hollebecque A, et al. Non-comparative, open-label, multiple cohort, phase 1/2 study to evaluate nivolumab (NIVO) in patients with virus-associated tumors (CheckMate 358): efficacy and safety in Merkel cell carcinoma (MCC). *Cancer Res*. 2017;77(13 suppl):Abstract CTO74.
- Kaufman HL, Russell J, Hamid O, et al. Avelumab in patients with chemotherapy-refractory metastatic Merkel cell carcinoma: a multicentre, single-group, open-label, phase 2 trial. *Lancet Oncol*. 2016;17(10):1374-1385.
- Bavencio (avelumab) injection [package insert]. Darmstadt, Germany: Merck KGaA; 2017.
- European Commission approves Bavencio (avelumab) for metastatic Merkel cell carcinoma. Darmstadt, Germany: Merck KGaA and Pfizer Inc; September 21, 2017. https://www.pfizer.com/news/press-release/press-release-detail/european_commission_approves_bavencio_avelumab_for_metastatic_merkel_cell_carcinoma. Accessed October 8, 2017.
- Bavencio (avelumab) approved for Merkel cell carcinoma in Japan. Darmstadt, Germany: Merck KGaA and Pfizer Inc; September 21, 2017. https://www.pfizer.com/news/press-release/press-release-detail/bavencio_avelumab_approved_for_merkel_cell_carcinoma_in_japan. Accessed October 8, 2017.
- Kaufman HL, Russell JS, Hamid O, et al. Updated efficacy of avelumab in patients with previously treated metastatic Merkel cell carcinoma after ≥ 1 year of follow-up: JAVELIN Merkel 200, a phase 2 clinical trial. *J Immunother Cancer*. 2018;6:7.