

Problems

- Functional groups
- Growth independent of biomass.
(ok si on suppose plein stock - à la MTE)
- No competition btw Functional groups
- Sensitivity to perturbations
- Sensitivity of trees to droughts.
(may be handled by "distribution")
- Seed consumption

Expectations

- Logistic growth
- Lower biomass and growth when herbivory
- Whitaker plot
- Escape from herbivory with height
- Seasonality and biomass
- Temp-growth relationship (stops at 0°)

First vibe exercises

- Extract veg. model and write a function with T° , PP as arguments
- Run time series for functional groups
- Re-interpret FG as
 - herbs
 - conif. trees
 - deciduous trees
- Add deciduous - non structural functional group.

Possible solution

Hierarchical model with:

- Regional dynamics with fire disturbance, derive age frequency distr.
- Local dynamics
 - Biomass dynamics according to Harport with $B(t)$ solution

Pr. Problem: $B(t)$ impossible because parameters are time dependent.


other solution: cohorts for vegetation

$$\frac{dB(a,t)}{dt}$$

$$\frac{dC}{dt}, \frac{dT}{dt}, \frac{dH}{dt}$$

H : herbs
C : coniferous
T : deciduous

problème: - coexistence
- taux transition



Unité spatiale: 5m, de sorte que on peut négliger les cas mixtes.