

Model Description: A Hierarchical Vegetation Model with Local and Regional Dynamics

1 Overview

We model vegetation hierarchically, distinguishing local patch dynamics from regional (cell-level) occupancy dynamics. A grid cell is composed of a very large number of small patches, so large that we track only the proportion of patches belonging to each vegetation type and age class. Each patch is small enough to contain at most one tree species, but it always contains herbs.

There are three functional groups: *evergreen trees*, *deciduous trees*, and *herbs*. After a disturbance (fire), all biomass in a patch is reset to zero and herbs are the first to establish. Forest develops through the subsequent growth of tree biomass and progressive shading of herbs by tree canopy.

The fire return interval differs between tree types and drives regional turnover. Differences in community composition across locations emerge from climate-dependent parameters and variation in herbivore pressure.

2 State Variables and Notation

The index i refers to any functional group (herbs, evergreen, or deciduous), while j refers specifically to tree species (evergreen or deciduous only), excluding herbs.

- $B_{i,x,t}$: total leaf biomass (g wet matter) of functional group i at location x and time t , integrated over the cell area.
- $p_{j,a,x,t}$: fraction of patches occupied by tree type j at patch age a (time since last fire)

at location x and time t . The constraint

$$\sum_a \sum_j p_{j,a,x,t} = 1 \quad (1)$$

ensures all patches are accounted for.

- $b_{j,a,x,t}$: leaf biomass density (g wet matter m^{-2}) of tree species j in patches of age a .
- $b_{\text{herb}|j,a,x,t}$: leaf biomass density of herbs in patches of tree type j at age a . Herbs and trees coexist in all patches, but tree species do not coexist within a single patch.
- Total biomass of functional group i is:

$$B_{i,x,t} = A_{\text{cell}} \sum_a p_{i,a,x,t} b_{i,a,x,t} \quad (2)$$

where A_{cell} is the cell area in m^2 . For herbs, the sum runs over both patch types weighted by their respective occupancy.

- $r_{j,x,t}$: relative biomass of tree species j among trees only:

$$r_{j,x,t} = \frac{B_{j,x,t}}{\sum_{j'} B_{j',x,t}} \quad (3)$$

3 Local Biomass Dynamics

The leaf biomass of each functional group in a patch of a given age and type follows a discrete-time difference equation adapted from Harfoot et al. (2014):

$$b_{i,a,x,t+\Delta t} = b_{i,a,x,t} + \Delta_{i,a,x,t}^{\text{Growth}} - \Delta_{i,a,x,t}^{\text{Mort}} \quad (4)$$

3.1 Mortality

$$\Delta_{i,a,x,t}^{\text{Mort}} = \delta \mu_{i,x,t} b_{i,a,x,t} + L_{i,a,x,t} \quad (5)$$

where $\mu_{i,x,t}$ is the mean annual leaf mortality rate (yr^{-1}) of functional group i , δ is a scalar converting the annual rate to the model time step, and $L_{i,a,x,t}$ is the leaf biomass consumed through herbivory during one time step. For constant herbivore populations:

$$L_{i,a,x,t} = \beta_i b_{i,a,x,t} \quad (6)$$

where β_i (yr^{-1}) is the functional-group-specific herbivory rate.

3.2 Growth and Competition

$$\Delta_{i,a,x,t}^{\text{Growth}} = NPP_{x,t} \psi \delta (1 - f_{\text{struct},i}) f_{\text{LeafMort},i} C_{i,a,x,t} \quad (7)$$

where $NPP_{x,t}$ is monthly terrestrial NPP ($\text{kg C m}^{-2} \text{ month}^{-1}$), ψ is the conversion factor from carbon to leaf wet matter, $f_{\text{struct},i}$ is the fractional allocation of primary production to structural tissue, and $f_{\text{LeafMort},i}$ is the fraction of non-structural production allocated to leaves (see Section 5).

Asymmetric competition via light attenuation

NPP is partitioned between herbs and the local tree species through an asymmetric competition term inspired by Beer's law. Trees shade herbs but herbs have no effect on trees. In an evergreen patch:

$$C_{\text{herb},a,x,t} = e^{-\alpha b_{\text{ever},a,x,t}} \quad (8)$$

and in a deciduous patch:

$$C_{\text{herb},a,x,t} = e^{-\alpha b_{\text{decid},a,x,t}} \quad (9)$$

where α ($\text{m}^2 \text{ kg}_C^{-1}$) is the light-extinction coefficient. By the asymmetry assumption:

$$C_{\text{ever},a,x,t} = C_{\text{decid},a,x,t} = 1 - C_{\text{herb},a,x,t} \quad (10)$$

so that $C_{\text{herb}} + C_{\text{tree}} = 1$ within each patch.

4 Regional Occupancy Dynamics

4.1 Aging and Disturbance

Each time step, a fraction $e_{j,x,t}$ of patches of tree type j is disturbed (burned) and returned to age 0. The remaining patches age by one step:

$$p_{j,a,t+\Delta t} = p_{j,a-1,t} (1 - e_{j,x,t}), \quad a \geq 1 \quad (11)$$

The oldest age class, $a_{\max}$, acts as a lumped class accumulating all patches older than $a_{\max}$:

$$p_{j,a_{\max},t+\Delta t} = (p_{j,a_{\max}-1,t} + p_{j,a_{\max},t}) (1 - e_{j,x,t}) \quad (12)$$

4.2 Colonization of Burned Patches

The total flux of patches disturbed at time t is:

$$D_{x,t} = \sum_j \sum_a p_{j,a,x,t} e_{j,x,t} \quad (13)$$

Newly burned patches are colonized by tree species in proportion to their regional relative biomass:

$$p_{j,0,x,t} = D_{x,t} r_{j,x,t} \quad (14)$$

Fire resets all leaf biomass to zero in newly created patches: $b_{j,0} = b_{\text{herb}|j,0} = 0$.

5 Climate-Dependent Parameters from Harfoot et al. (2014)

The following equations follow the terrestrial plant sub-model of Harfoot et al. (2014).

5.1 Net Primary Production

Annual NPP follows the Miami model Lieth1975:

$$NPP_y^T = \min(NPP_T, NPP_P) \quad (15)$$

$$NPP_T = \frac{NPP_{\max}}{1 + e^{c_p - m_p \bar{T}}} \quad (16)$$

$$NPP_P = NPP_{\max} (1 - e^{-\rho P}) \quad (17)$$

where $NPP_{\max}$ is the maximum possible NPP ($\text{kg C m}^{-2} \text{ yr}^{-1}$), c_p and m_p are coefficients relating NPP to mean annual temperature $\bar{T}$ ($^{\circ}\text{C}$), ρ relates NPP to total annual precipitation P (mm yr^{-1}).

Monthly NPP is then obtained by scaling annual NPP by the monthly seasonality factor $\omega_{\text{cell},m}$ (derived from Terra/MODIS remote-sensing data):

$$NPP_m^T = NPP_y^T \cdot \omega_{\text{cell},m} \quad (18)$$

5.2 Fractional Allocation to Structural Tissue

$$f_{\text{struct}} = \min \left(\frac{f_{\text{struct}}^{\min} e^{\theta_{f_{\text{struct}}} NPP_y^T}}{1 + f_{\text{struct}}^{\min} (e^{\theta_{f_{\text{struct}}} NPP_y^T} - 1)}, 0.99 \right) \cdot f_{\text{struct}}^{\max} \quad (19)$$

where $f_{\text{struct}}^{\min} = 0.01$.

5.3 Leaf Mortality Rates

The mean leaf mortality rate is a weighted average of evergreen and deciduous rates:

$$\mu_{\text{Leaf}} = i e^{f_{\text{ever}} \ln \mu_{\text{ever}} + (1 - f_{\text{ever}}) \ln \mu_{\text{decid}}} \quad (20)$$

The evergreen and deciduous leaf mortality rates depend on mean annual temperature $\bar{T}^C$ ($^{\circ}\text{C}$):

$$\mu_{\text{ever}} = e^{(m_e \bar{T}^C - c_e)} \quad (21)$$

$$\mu_{\text{decid}} = e^{-(m_d \bar{T}^C + c_d)} \quad (22)$$

Both rates are bounded: $\mu_{\text{ever}} \in [\mu_e^{\min}, \mu_e^{\max}]$ and $\mu_{\text{decid}} \in [\mu_d^{\min}, \mu_d^{\max}]$.

5.4 Fine Root Mortality Rate

$$\mu_{\text{FineRoot}} = e^{m_r T_{(t)}^C + c_r} \quad (23)$$

where $T_{(t)}^C$ is the monthly mean temperature ($^{\circ}\text{C}$). The rate is bounded: $\mu_{\text{FineRoot}} \in [\mu_r^{\min}, \mu_r^{\max}]$.

5.5 Fractional Leaf Allocation

The fraction of non-structural NPP allocated to leaves (versus fine roots) is:

$$f_{\text{LeafMort}} = \frac{\mu_{\text{Leaf}}}{\mu_{\text{Leaf}} + \mu_{\text{FineRoot}}} \quad (24)$$

5.6 Proportion of Evergreen Productivity

The fraction of NPP produced by evergreen leaves depends on the fraction of the year experiencing frost (F_{frost} , the normalized number of frost days):

$$f_{\text{ever}} = a_{f_{\text{ever}}} F_{\text{frost}}^2 + b_{f_{\text{ever}}} F_{\text{frost}} + c_{f_{\text{ever}}} \quad (25)$$

bounded to $[0, 1]$.

6 Parameter Values

7 Application: Sherbrooke, Canada

Table 2 summarizes the climate-derived parameters computed for Sherbrooke, Québec, Canada (45.4°N , 71.9°W) using the Madingley spatial raster inputs.

References

- [1] Harfoot, M.B.J., Newbold, T., Tittensor, D.P., Emmott, S., Hutton, J., Lyutsarev, V., Smith, M.J., Scharlemann, J.P.W., & Purves, D.W. (2014). Emergent global patterns of ecosystem structure and function from a mechanistic general ecosystem model. *PLOS Biology*, 12(4), e1001841.

- [2] Lieth, H. (1975). Modelling the primary productivity of the world. In: Lieth, H. & Whitaker, R.H. (eds.), *Primary Productivity of the Biosphere*. Springer-Verlag, New York, pp. 237–263.

Table 1: Madingley plant model parameter values (Harfoot et al. 2014, `TerrestrialCarbon.cpp`).

Symbol	Description	Value	Units
<i>Miami NPP model</i>			
$NPP_{\max}$	Maximum possible NPP	0.9616	kg C m ⁻² yr ⁻¹
c_p	Temperature NPP intercept	0.2375	—
m_p	Temperature NPP slope	0.1006	°C ⁻¹
ρ	Precipitation NPP coefficient	0.001184	mm ⁻¹
<i>Structural allocation</i>			
$\theta_{f_{\text{struct}}}$	Scalar for f_{struct}	7.155	—
$f_{\text{struct}}^{\max}$	Maximum structural allocation	0.3627	—
<i>Evergreen leaf mortality</i>			
m_e	Slope vs. temperature	0.04027	°C ⁻¹
c_e	Intercept	1.01307	—
$\mu_e^{\min}$	Minimum rate	0.01	yr ⁻¹
$\mu_e^{\max}$	Maximum rate	24.0	yr ⁻¹
<i>Deciduous leaf mortality</i>			
m_d	Slope vs. temperature	0.02058	°C ⁻¹
c_d	Intercept	−1.1952	—
$\mu_d^{\min}$	Minimum rate	0.01	yr ⁻¹
$\mu_d^{\max}$	Maximum rate	24.0	yr ⁻¹
<i>Fine root mortality</i>			
m_r	Slope vs. temperature	0.04309	°C ⁻¹
c_r	Intercept	−1.4784	—
$\mu_r^{\min}$	Minimum rate	0.01	yr ⁻¹
$\mu_r^{\max}$	Maximum rate	12.0	yr ⁻¹
<i>Evergreen fraction</i>			
$a_{f_{\text{ever}}}$	Quadratic coefficient	1.2708	—
$b_{f_{\text{ever}}}$	Linear coefficient	−1.8286	—
$c_{f_{\text{ever}}}$	Intercept	0.8449	—
<i>Conversion factors</i>			
ψ	C to leaf wet matter (kg C / g dry matter)	$0.476^{-1} \times 0.213^{-1}$	g wet g ⁻¹ C
<i>New model parameters (this study)</i>			
α	Beer-Lambert shading coefficient	5.0	m ² kg _C ⁻¹
β_i	Herbivory rate by functional group	0.01–0.05	yr ⁻¹
e_j	Annual fire disturbance rate by tree type	1/100–1/150	yr ⁻¹
$a_{\max}$	Maximum tracked patch age	600	months

Table 2: Derived parameter values for Sherbrooke, Canada.

Quantity	Value	Units
Mean annual temperature	3.94	°C
Total annual precipitation	1256	mm yr ⁻¹
Total annual AET	488	mm yr ⁻¹
Fraction of year with frost (F_{frost})	0.517	—
Fraction of year with fire (F_{fire})	0.000	—
Annual NPP (Miami model)	0.519	kg C m ⁻² yr ⁻¹
f_{ever}	0.239	—
f_{struct}	0.106	—
μ_{ever}	0.426	yr ⁻¹
μ_{decid}	3.047	yr ⁻¹
μ_{FineRoot}	0.270	yr ⁻¹
$f_{\text{LeafMort,ever}}$	0.612	—
$f_{\text{LeafMort,decid}}$	0.919	—