Biospectra : ISSN: 0973-7057, Vol. 20(2), September, 2025, pp. 117-120 ***Interdisciplinary Science***
An International Biannual Refereed Journal of Life Sciences

ISSN : 0973-7057

Synergistic effects of food-based bioactive compounds and pharmaceuticals: A new frontier in integrative medicine

Surabhi Singh[a], Vijay Kant Pandey[b]* & Santosh Kumar Singh[c]
[a]Department of Botany, S.V.M. Science and Technology P.G. College, Lalganj Pratapgarh, Uttar Pradesh, India
[b]Department of Life Science, Netaji Subhas University, Jamshedpur, Jharkhand, India
[c]Department of Biotechnology, Arka Jain University, Jamshedpur, Jharkhand, India

Received : 15th April, 2025 ; Revised : 16th May, 2025
DOI:-https://doi.org/10.5281/zenodo.18504358

Abstract- Integrative medicine is steering healthcare towards a more holistic approach, blending traditional treatments with evidence-based complementary methods. One exciting area of focus is how food-derived bioactive compounds, or nutraceuticals, can work alongside pharmaceuticals to enhance treatment results and minimize side effects. This review shines a light on the latest research into their synergistic effects, particularly looking at mechanisms like improved pharmacokinetics, multi-target action, and fewer adverse effects. While the potential is promising, there are still hurdles to overcome, such as bioavailability, standardization, and interactions between herbs and drugs. The paper stresses the importance of thorough clinical validation and wraps up by suggesting that a scientifically informed integration of bioactives with medications could lead to more effective and personalized therapies.

Keywords: Integrative medicine, pharmaceutical, pharmacokinetics, bioactives, drugs

INTRODUCTION

The increasing global challenge of chronic diseases like cancer, heart issues, diabetes, and neurodegenerative disorders has highlighted the shortcomings of relying solely on traditional drug therapies.[1] At the same time, both consumers and researchers are becoming more interested in the health benefits of food that go beyond just basic nutrition. This has given rise to the concept of "nutraceuticals," which are bioactive compounds from food sources known to offer physiological benefits or protective effects against various diseases.[2] Integrative medicine aims to blend the best of conventional and alternative treatments, focusing on the individual as a whole and utilizing natural, less invasive method whenever possible. Within this context, the combination of food-based bioactives and pharmaceuticals is not just a coincidence; it's a hot topic of research. The idea is that these compounds can complement medications, potentially leading to better health outcomes through their combined effects. This review delves into this exciting area, looking at the evidence, mechanisms, and implications of merging dietary bioactives with established drugs.

Mechanisms of Synergistic Action

The relationship between nutraceuticals and medications can be divided into two main types of interactions: pharmacokinetic and pharmacodynamic.

Pharmacokinetic Synergy: This refers to how a drug's absorption, distribution, metabolism, or excretion

*Corresponding author :
Phone : 8825290968
E-mail : pandeyvijay00@gmail.com

(often abbreviated as ADME) can be influenced. A classic example is how certain substances interact with the cytochrome P450 (CYP450) enzyme system and drug transporters like P-glycoprotein (P-gp). Take piperine, for instance, which is a key component in black pepper. It's known to inhibit both CYP3A4 and P-gp. Research has indicated that when piperine is taken alongside drugs like curcumin (the active ingredient in turmeric), it can boost their bioavailability by an astonishing 2000%. This effect also extends to other medications, such as phenytoin and propranolol.[3] Likewise, flavonoids found in citrus fruits and green tea can block these metabolic pathways, which might lead to higher drug plasma levels and improved effectiveness, but it's important to manage this carefully to avoid potential toxicity.

Pharmacodynamic Synergy: This type of interaction happens when a bioactive compound and a drug work together at the target site, enhancing the therapeutic effect through complementary actions. This approach, which targets multiple pathways, is especially beneficial for treating complex diseases

Cancer Therapy: Curcumin has been the focus of a lot of research for its potential to make cancer cells more sensitive to chemotherapy. It influences several signaling pathways, such as NF-κB, STAT3, and AKT, which play key roles in cell growth, programmed cell death, and resistance to chemotherapy. When used alongside chemotherapeutic drugs like paclitaxel or 5-fluorouracil, curcumin can boost the effectiveness of cancer cell death and lower the necessary dose of the toxic medication, which helps to minimize side effects.[4]

Cardiovascular Health: Combining statins, which are drugs that lower cholesterol, with omega-3 fatty acids from fish oil or plant sterols creates a strong synergistic effect for managing lipid profiles. Statins mainly work by blocking cholesterol production, while omega-3s help decrease triglyceride levels, and plant sterols prevent the absorption of dietary cholesterol. This multi-faceted approach effectively tackles dyslipidemia.[5]

Amelioration Adverse Drug Reactions

One of the major perks of blending bioactives with traditional medications is the chance to lessen those pesky side effects that come with drugs, ultimately boosting patients' quality of life and their willingness to stick to their treatment plans.

Chemotherapy-Induced Toxicity: Gingerol, which comes from ginger, has shown promise in easing nausea and vomiting caused by chemotherapy. Its ability to combat these symptoms seems to work alongside conventional anti-nausea medications like 5-HT3 receptor antagonists.[6] Likewise, antioxidant-rich compounds such as sulforaphane (found in cruciferous veggies) and resveratrol (from grapes) have been noted for their potential to shield healthy tissues from oxidative harm during radiotherapy and certain chemotherapy treatments, all while keeping their anti-cancer effects intact.

Gastrointestinal Protection: Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) like aspirin and ibuprofen can lead to gastric ulcers. However, flavonoids and other polyphenols, known for their anti-inflammatory and antioxidant benefits, can help safeguard the stomach lining. Taking these compounds alongside NSAIDs might reduce the risk of ulcers.[7]

Table 1: Examples of Synergistic Interactions Between Food-Based Bioactives and Pharmaceuticals

Bioactive Compound (Source)	Pharmaceutical Drug (Class)	Observed Synergistic Effect	Proposed Mechanism
Piperine (Black Pepper)	Curcumin, Phenytoin, Propranolol	Significantly enhanced bioavailability and efficacy	Inhibition of drug-metabolizing enzymes (CYP450) and efflux pump (P-glycoprotein)
Curcumin (Turmeric)	Paclitaxel, 5-Fluorouracil (Chemotherapy)	Enhanced cancer cell death; reduced required chemo dose	Multi-target modulation of signaling pathways (NF-κB, STAT3) inducing apoptosis and overcoming chemo-resistance
Omega-3 Fatty Acids (Fish Oil)	Statins (Lipid-lowering)	Improved overall lipid profile (lower triglycerides & LDL-C)	Complementary mechanisms: Statins inhibit cholesterol synthesis; Omega-3s reduce triglyceride production
Gingerols (Ginger)	5-HT3 Receptor Antagonists (Anti-emetic)	Reduced chemotherapy-induced nausea and vomiting	Complementary anti-emetic mechanisms targeting different pathways in the nausea response
Flavonoids/Polyphenols (e.g., Berries, Tea)	NSAIDs (e.g., Aspirin, Ibuprofen)	Protection against gastric mucosal damage and ulcers	Antioxidant and anti-inflammatory activities that protect the stomach lining from NSAID-induced injury

Challenges and Considerations

While the potential is certainly exciting, the journey toward clinical application is filled with hurdles that need to be tackled.

Standardization and Bioavailability: The levels of bioactive compounds in food can fluctuate significantly depending on factors like growing conditions, processing methods, and storage practices. Additionally, many promising compounds, such as curcumin and resveratrol, struggle with poor solubility in water and low systemic bioavailability. It's essential to create standardized formulations with high bioavailability think nanoparticles, phospholipid complexes, or pairing them with enhancers like piperine to ensure consistent clinical results.[8]

Risk of Antagonistic Interactions: Not every food-drug interaction is a good one; some can actually be harmful. For example, foods rich in vitamin K, like leafy greens, can counteract the blood-thinning effects of warfarin. Similarly, St. John's Wort, a popular herbal supplement, can significantly boost the activity of CYP3A4 and P-gp, which may lower the plasma levels of various medications, including antidepressants, immuno suppressants, and birth control pills, potentially leading to treatment failures.[9] Thorough research is essential to pinpoint and warn against these dangerous interactions.

Challenges and Considerations

Despite the promising potential, the path to clinical application is fraught with challenges that must be addressed.

Standardization and Bioavailability: The concentration of bioactive compounds in food can vary dramatically based on growing conditions, processing, and storage. Furthermore, many promising compounds, like curcumin and resveratrol, suffer from poor aqueous solubility and low systemic bioavailability. Developing standardized, high-bioavailability formulations (e.g., using nanoparticles, phospholipid complexes, or co-administration with enhancers like piperine) is crucial for reproducible clinical outcomes.[8]

Risk of Antagonistic Interactions: Not all food-drug interactions are beneficial. Some can be antagonistic or dangerous. For instance, vitamin K-rich foods (e.g., leafy greens) can antagonize the anticoagulant effect of warfarin. St. John's Wort, a common herbal supplement, is a potent inducer of CYP3A4 and P-gp, leading to reduced plasma levels of many drugs, including antidepressants, immunosuppressants, and oral contraceptives, potentially resulting in therapeutic failure .[9] Rigorous research is needed to identify and contraindicate such harmful interactions.

Evidence Gap and Clinical Translation: While in vitro and animal studies are abundant, well-designed, large-scale human clinical trials are still relatively scarce. More research is needed to establish optimal dosages, treatment durations, and formulations for specific drug-nutraceutical combinations. The regulatory landscape for such combinations also remains complex and varies by region.

CONCLUSION

Bringing together food-based bioactive compounds and pharmaceuticals represents a significant leap in integrative medicine. This strategy is designed to enhance drug effectiveness, cut down on dosages, and lessen side effects, ultimately providing more personalized and effective solutions for chronic, multifaceted diseases. To advance this area, we need clinical trials, innovative delivery systems, clear guidelines, and teamwork across various fields. The fusion of nutraceuticals and pharmaceuticals offers a powerful approach that could exceed the benefits of conventional therapies, paving the way for more customized treatments and transforming how we manage chronic diseases through integrative, evidence-based healthcare practices.

REFERENCES

1. **Seca, A. M., & Pinto, D. C. 2018.** Plant secondary metabolites as anticancer agents: Successes in clinical trials and therapeutic application. *International Journal of Molecular Sciences,* **19(1):** 263.

2. **Kalra, E. K. 2003.** Nutraceutical-definition and introduction. *AAPS PharmSci,* **5(3):**E25.

3. **Dudhatra, G. B., Mody, S. K., Awale, M. M., Patel, H. B., Modi, C. M., Kumar, A., Kamani, D. R., & Chauhan, B. N. 2012.** A comprehensive review on pharmacotherapeutics of herbal bioenhancers. *The Scientific World Journal,* **2012:** 637953.

4. **Giordano, A., & Tommonaro, G. 2019.** Curcumin and cancer. *Nutrients,* **11(10):** 2376.

5. **Backes, J. M., Howard, P. A., & Moriarty, P. M. 2016.** Role of omega-3 fatty acids in the treatment of

hypertriglyceridemia. *Pharmacotherapy: The Journal of Human Pharmacology and Drug Therapy,* **36(5):** 551–562.

6. **Crichton, M., Marshall, S., Marx, W., Isenring, E., & Lohning, A. 2019.** Efficacy of ginger (*Zingiber officinale*) in the prevention of chemotherapy-induced nausea and vomiting: A systematic review. *Journal of Advanced Nursing,* **75(1):** 30–46.

7. **Yuan, L. W., Jiang, X. M., Xu, Y. L., Huang, M. Y., Chen, Y. C., Yu, W. B., & Su, Z. Q. 2021.** Luteolin ameliorates aspirin-induced gastric ulcer in mice by modulating the MAPK/NF-κB signalling pathways. *Immunopharmacology and Immunotoxicology,* **43(4):** 443–452.

8. **Panda, V. S., & Khajuria, R. 2020.** Evaluation of the bioavailability of major withanolides of *Withania somnifera* and the enhanced bioavailability of withaferin A and withanolide A through herbal solid dispersion technique. *Journal of Dietary Supplements,* **17(1):**51–67.

9. **Hu, Z., Yang, X., Ho, P. C., Chan, S. Y., Heng, P. W., Chan, E., Duan, W., Koh, H. L., & Zhou, S. 2005.** Herb-drug interactions: A literature review. *Drugs,* **65(9):** 1239–1282.

***