

From the Editor Desk

Biospectra is an official publication of MSET-ICCB, Ranchi. On behalf of the Editorial Board of the Journal Biospectra and my Co-Editors, I am glad to present the Volume 17, Issue 1 of the Journal. The publication of the Journal started in September, 2006. With the great effort of Former Editor-in-Chief, Late Prof. (Dr.) Mahendra Prasad the Journal is indexed in Clarivate Analytics as Zoological Records. The Journal has a Scientific Journal Impact Factor (SJIF-2022) of 8.379. The Journal is also listed in UGC-CARE Group II through Web of Science (Journals indexed in globally recognized databases). We are putting our effort to list the Journal in UGC-CARE Group I & also preparing for obtaining NAAS rating. As we had NAAS rating of 2.3 in the list of "Rating of Scientific Journals 2012 (Effective from Jan 1, 2013)". From the last issue onwards the Academic Council of MSET-ICCB & Executive Committee of Biospectra has nominated me as the Editor-in-Chief of Biospectra. As I have stated in earlier issue that the papers are published only after checking plagiarism and careful selection through a double-blind peer-review process.

I am happy to state that we have worked hard on our words and the policies of publication. We have connected with several new reviewers from all over India and abroad also.

I extend my great honour and sincerest thanks to all the contributors of the articles, members of trust as well as editorial board and innumerable man power support in bringing out BIOSPECTRA, Volume 17, Issue 1, March, 2022, on ground for the enlightenment of people of Life Sciences.

31.03.2022

(Editor-in-Chief)
Prof. (Dr.) Jyoti Kumar
F.B.S, F.P.S.I,
Former University Professor & Head,
University Department of Botany
Former Dean, Faculty of Sciences,
Ranchi University, Ranchi, Jharkhand
Former Director, IQAC, Ranchi University,
Former Director, UGC HRDC, Ranchi University,

Biospectra : ISSN: 0973-7057, Vol. 17(1), March, 2022, pp. 1-4 **1(AS).**
An International Biannual Refereed Journal of Life Sciences ***Animal Science***

ISSN : 0973-7057

Int. Database Index: 616 www.mjl.clarivate.com

Antagonistic activity of *Lactobacilli* derived from raw milk

Shwet Nisha*, Kumar Ramashankar & Rashmi Kumari
University Department of Zoology, B.N.M. University, Madhepura, Bihar, India

Received : 15th October, 2021 ; Revised : 16th December, 2021

Abstract- Altogether 15 milk samples were collected from local dairy firm of Madhepura district and *Lactobacilli* were isolated. Altogether 3 isolates were obtained, LM1, LM2 and LM3 which were identified as *Lactobacillus casei*, *Lactobacillus brevis* and *Lactobacillus fermentum* respectively. The identification were made on the basis of their morphological, biochemical and sugar test as prescribed in Bergey's manual of systematic Bacteriology (second edition-2001). The antagonistic study of isolated strain was made against four pathogenic bacteria *Pseudomonas aeruginosa*, *Escherichia coli*, *Staphylococcus aureus* and *Proteus sp.*. The inhibition zone against *Pseudomonas aeruginosa, Escherichia coli* and *Proteus sp.*was observed maximum by *Lactobacillus brevis* and against *Staphylococcus aureus*, the maximum inhibition zone was observed by *Lactobacillus fermentum*.

Key words: Niche, Niche Relaxation, Correlation, Amoeba, Copepod

INTRODUCTION

Raw milk is natural growth medium for micro-organism. An integral part of raw milk microflora is species of *Lactobacilli* as *Lactobacillus casei*, *L. plantarum*, *L. rhamnosus*, *L. curvatus*, *L. bravis*, *L. fermentum* and some other Lactic acid bacteria like *Leuconostoc lactis*.

Lactobacilli are rod shaped gram positive, Catalase negative, non-pathogenic bacteria present in milk and milk products. Several *Lactobacillus* strains are used as starter culture for the manufacture of fermented food.[1] These bacteria play a vital role in our daily life for fermentation, preservation, production and of wholesome food.

Lactic acid bacteria produced various compounds as organic acids, diacetyls, hydrogen peroxide, Vitamins and anti-microbial substance, bacteriocin. Antimicrobial activity of *Lactobacillus* against *E. coli*, *S. aureus* and *Solmonella* was reported by several workers.[2-5]

Lactobacillus casei have a wide range of pH and temperature and complement the growth of *L. acidophillus*, a producer of enzyme amylase. *L. sporogenesis* supports the growth of beneficial bacteria and help to maintain a healthy balance of microflora in the intestinal environment. A lot of LAB has been noted to have nutritional benefits, improve Lactose utilization, and have anti-cholesterol, anti-carcinogenic activity. Intestinal Lactic acid bacteria are considered to have health benefits for human being and introduced as probiotic including *L. rhamnosus*, *L. casei* and *L. johnsonii*.[6]

MATERIAL & METHOD

*Corresponding author :
Phone : 9006991000
E-mail : prf.arunkumar@gmail.com

Milk samples were collected from local diary firm of Madhepura in 500ml sterilized bottles and brought to the laboratory in ice boxes. From each sample, 1ml of milk was mixed in 10ml distilled water and serial dilution was made up to 10^{-5}. From each sample, 0.1ml inoculum was inoculated in MRS medium and incubated at 30°C temperature for 48 hours. Colonies were purified by sub-culturing in same medium. From each plate, colonies were examined for morphological study, gram staining, biochemical test and sugar test. On the basis of biochemical test and sugar test, isolates were categorised in following three categories: LM1, LM2 and LM3. The strains were identified on the basis of their morphological character, biochemical and sugar test as prescribed in Bergey's manual of systematic Bacteriology (second edition-2001).

Four pathogens *Pseudomonas aeruginosa*, *Escherichia coli*, *Staphylococcus aureus* and *Proteus sp.*were obtained from local pathological lab. All test pathogenic bacteria were cultured in Nutrient broth.

Antagonistic study:

For the antagonistic study of isolated strains of *Lactobacilli* against pathogenic bacteria was performed by well diffusion method.

Petri dishes containing 20ml of Muller Hinton agar were prepared and inoculated with 0.1ml pathogenic culture. The plates were allowed to solidify and stored in refrigerator for 2 hours. Four wells were made in culture plate and filled using different concentration (0.01ml, 0.02ml, 0.03ml and 0.04ml) of cell free filtrate of each isolate. Plates were incubated at 37°C for 24 hours. The diameter of inhibition zone was measured using Callipers.

Table 1- Biochemical test

Sl. No.	Isolate	Gram staining	Indole	MR	VP	Catalase	Oxidase	Nitrate reducation
1	LM1	+ve	-ve	-ve	-ve	-ve	-ve	+ve
2	LM2	+ve	-ve	-ve	-ve	-ve	-ve	+ve
3	LM3	+ve	-ve	-ve	-ve	-ve	-ve	+ve

Table 3- Antagonistic activity of selected isolates against pathogens

Sl. No.	Pathogens	LM1	LM2	LM3
1	*Pseudomonas aeruginosa*	7mm	13mm	6.5mm
2	*Escherichia coli*	7.5mm	13.5mm	9.5mm
3	*Staphylococcus aureus*	9.5mm	10mm	12.5mm
4	*Proteus sp.*	8mm	14mm	11mm

RESULT

Altogether 15 samples were tested in which 3 isolates were obtained which were assigned as LM1, LM2 and LM3. All isolates were gram positive while Indole, Catalase and Oxidase show negative. In sugar test, LM1 was negative in Glucose and sorbitol and positive in all other tested sugars. LM2 was positive for Xylose and negative for all other sugars. LM3 was negative in Rhamonose and positive in all other test sugars. On the basis of biochemical test and sugar fermentation test, LM1 was identified as *Lactobacillus casei*, LM2 as *Lactobacillus brevis* and LM3 as *Lactobacillus fermentum*.

The result is tabulated in Table no. 01 and 02. All three isolates strains inhibited the growth of all four pathogenic bacteria under test. The inhibition zone varies for different strains for different pathogenic bacteria. The maximum inhibition zone against *Pseudomonas aeruginosa* was of LM2 (13mm) and minimum of LM3 (6.5mm), against *E. coli* the highest inhibition zone was recorded for LM2 (13.5mm) and minimum for LM1 (7.5mm). The maximum inhibition zone against *S. aureus* was recorded for LM3 (12.5mm) and minimum for LM1 (9.5mm) and Inhibition zone against *Proteus* was maximum for LM2 (14mm) and minimum for LM1 (8mm). The result is tabulated in Table no. 03.

CONCLUSION

Altogether 15 milk samples were collected from local area of Madhepura district. *Lactobacilli* from each milk were isolated by culturing in MRS medium, Gram staining, Biochemical test and morphological study of bacteria from

Table 2- Sugar fermentation test

Sugar test	LM1	LM2	LM3
Xylose	+ve	+ve	+ve
Cellubiose	+ve	-ve	+ve
Glucose	-ve	-ve	+ve
Mannitol	+ve	-ve	+ve
Ramonose	+ve	-ve	-ve
Raffinose	+ve	-ve	+ve
Ribose	+ve	-ve	+ve
Sorbitol	-ve	-ve	+ve
Trehalose	+ve	-ve	-ve
Fructose	+ve	-ve	+ve

all samples were performed. Three different isolates were obtained from all samples. On the basis of their biochemical study and sugar test, the isolates were identified as *Lactobacillus casei*, *Lactobacillus brevis* and *Lactobacillus fermentum*. The antagonistic study of these *Lactobacilli* species were performed against four pathogenic bacteria- *Pseudomonas aeruginosa, Escherichia coli, Staphylococcus aureus* and *Proteus* sp.. The antagonistic study was performed by well diffusion method. The inhibition zone against *Pseudomonas aeruginosa, Escherichia coli* and *Proteus sp.* was observed maximum by *Lactobacillus brevis* and against *Staphylococcus aureus*, the maximum inhibition zone was observed by *Lactobacillus fermentum*.

REFERENCE

1. **Badis, A., Guetarni, D., Moussa-Boudjema, B., Henni, D.E., Kihal M. 2004.** Identification and technological properties of lactic acid bacteria isolated from raw goats milk of furnace Algerian races. *Food Microbiol.,* **21(5):** 279-288.
2. **Gopal Pramod, K., Prasad, J., Smart, J., Gill Harsharanjit, S. 2001.** *In vitro* adherence properties of *Lactobacillus rhamnosus* DR20 and *Bifidobacterium lactis* DR10 strains and their antagonistic activity against an enterotoxigenic *Escherichia coli. Int. J. Food Microbiol.,***67(3):**207-216.
3. **Karska-Wysocki, B., Bazo, M., Smoragiewicz, W. 2010**. Antibacterial activity of *Lactobacillus acidophilus* and *Lactobacillus casei* against methicillin-resistant *Staphylococcus aureus* (MRSA). *Microbiol. Res.,* **165(8):** 674-86.
4. **Makras, Lefteris, Vagelis Triantafyllou, Domitille Fayol-Messaoudi, Tom Adriany, Georgia Zoumpopoulou, Effie Tsakalidou, Alain Servin, and Luc De Vuyst. 2006.** Kinetic analysis of the antibacterial activity of probiotic *Lactobacilli* towards *Salmonella enteric serovar Typhimurium* reveals a role for lactic acid and other inhibitory compounds. *Res. Microbiol.,* **157(3):** 241-7.
5. **Saxena, S., Dutta, R. 2011.** Antimicrobial resistance pattern of *Shigella* species over five years at a tertiary-care teaching hospital in North India. *J. Health Popul. Nutr.,* **29(3):** 292-5.
6. **Elazab, N.A., Mendy, J., Gasana, E.R., Vieira, A., Quizon, Forno, E. 2013.** Probiotic administration in early life, atopy, and asthma: a meta-analysis of clinical trials. *Pediatrics,* **132(3):** 666-676.

***