22. **Bruning, J.L. and Kintz, B.L. 1977.** *Computational Handbook of Statistics*, 2nd ed., Glenview, III: Scott, Foresman.

23. **Örn, S., Holbech, H., Madsen, T. H., Norrgren, L., & Petersen, G. I. 2003.** Gonad development and vitellogenin production in zebrafish (*Danio rerio*) exposed to ethinylestradiol and methyltestosterone. *Aquatic Toxicology*. **65(4):** 397-411.

24. **Sonja W. Scholz, Henry Houlden, Claudia Schulte, Manu Sharma, Abi Li, Daniela Berg, Anna Melchers, Reema Paudel, J. Raphael Gibbs, Javier Simon-Sanchez, Coro Paisan-Ruiz, Jose Bras, Jinhui Ding, Honglei Chen, Bryan J. Traynor, Sampath Arepalli, Ryan R. Zonozi, Tamas Revesz, Janice Holton, Nick Wood, Andrew Lees, Wolfgang Oertel Ullrich Wüllner, Stefano Goldwurm, Maria Teresa Pellecchia, Thomas Illig, Olaf Riess, Hubert H. Fernandez, Ramon L. Rodriguez, Michael S. Okun, Werner Poewe, Gregor K. Wenning, John A. Hardy, Andrew B. Singleton, Thomas Gasser MD. 2009.** SNCA variants are associated with increased risk for multiple system atrophy. *Annals of Neurology: Official Journal of the American Neurological Association and the Child Neurology Society*. **65(5):** 610-614.

25. **Dethlefsen, V., Von Westernhagen, H., & Cameron, P. 1996.** Malformations in North Sea pelagic fish embryos during the period 1984–1995. *ICES Journal of Marine Science*. **53(6):** 1024-1035.

26. **Gross-Sorokin, M. Y., Roast, S. D., & Brighty, G. C. 2006.** Assessment of feminization of male fish in English rivers by the Environment Agency of England and Wales. *Environmental Health Perspectives*. **114(Suppl 1):** 147-151.

27. **Lavado, R., Thibaut, R., Raldúa, D., Martýn, R., & Porte, C. 2004.** First evidence of endocrine disruption in feral carp from the Ebro River. *Toxicology and Applied Pharmacology*. **196(2):** 247-257.

28. **Sumpter, J. P., & Jobling, S. 1995.** Vitellogenesis as a biomarker for estrogenic contamination of the aquatic environment. *Environmental Health Perspectives*. **103(suppl 7):** 173-178.

29. **Beronius, A., Rudén, C., Hanberg, A., & Håkansson, H. 2009.** Health risk assessment procedures for endocrine disrupting compounds within different regulatory frameworks in the European Union. *Regulatory Toxicology and Pharmacology*. **55(2):** 111-122.

30. **Purdom, C. E., Hardiman, P. A., Bye, V. V. J., Eno, N. C., Tyler, C. R., & Sumpter, J. P. 1994.** Estrogenic effects of effluents from sewage treatment works. *Chemistry and Ecology*. **8(4):** 275-285.

31. **Länge, R., Hutchinson, T. H., Croudace, C. P., Siegmund, F., Schweinfurth, H., Hampe, P. Panter, G.H. & Sumpter, J. P. 2001.** Effects of the synthetic estrogen 17α ethinylestradiol on the life cycle of the fathead minnow (*Pimephales promelas*). *Environ mental Toxicology and Chemistry: An International Journal*. **20(6):** 1216-1227.

32. **Balch, G. C., Mackenzie, C. A., & Metcalfe, C. D. 2004.** Alterations to gonadal development and reproductive success in japanese medaka (*Oryzias latipes*) exposed to 17α ethinylestradiol. *Environmental Toxicology and Chemistry: An International Journal*. **23(3):**782-791.

33. **Parrott, J.L. and Blunt, B.R., 2005.** Life cycle exposure of fathead minnows (*Pimephales promelas*) to an ethinylestradiol concentration below 1 ng/L reduces egg fertilization success and demasculinizes males. *Environmental Toxicology: An International Journal*. **20(2):**131-141.

34. **Shved, N., Berishvili, G., D'Cotta, H., Baroiller, J. F., Segner, H., Eppler, E., & Reinecke, M. 2007.** Ethinylestradiol differentially interferes with IGF-I in liver and extrahepatic sites during development of male and female bony fish. *Journal of Endocrinology*. **195(3):** 513-524.

35. **Caldwell, D. J., Mastrocco, F., Hutchinson, T. H., Lange, R., Heijerick, D., Janssen, C., Anderson, P.D. & Sumpter, J. P. 2008.** Derivation of an aquatic predicted no-effect concentration for the synthetic hormone, 17α-ethinyl estradiol. *Environmental Science & Technology*. **42(19):** 7046-7054.

***

Biospectra : ISSN: 0973-7057, Vol. 17(1), March, 2022, pp. 147-150
An International Biannual Refereed Journal of Life Sciences

30(AS).
Animal Science

ISSN : 0973-7057

Int. Database Index: 616 www.mjl.clarivate.com

Analytical studies on the effect of cestode parasites on some commercially important freshwater fishes of Madhepura region

Amrita Kumari, Kumar Ramashankar & Arun Kumar
University Department of Zoology, Ranchi University, Ranchi, Jharkhand, India

Received : 15th February, 2022 ; Revised : 16th March, 2022

Abstract- Fish production plays an important role in the economy of a country. They are rich source of protein diet. There are many factors which effects the production of fishes, parasites infection is one of the main reasons for increasing mortality of fish. The present study was performed to find out the effect of parasites on some economically important freshwater fishes with special reference to cestodes (tapeworms), as two more types of parasites were there other than cestodes which can badly affect the fish production such as trematode (flatworms) and nematode (roundworms). A total of 160 commercially important fishes belonging to eight families were taken for this study. 20 each from the following species *Catla catla, Labeo rohita, Channa punctatus, Clarias batrachus, Cirrhinus mrigala, Heteropneustes fossilis, Mystus vittatus, Wallago attu* were taken. Extreme infestations were observed in the fishes. The production of fishes were decreased due to the infestation of cestode parasites. Very few work has been done in this topic till date, especially studies on cestode parasites. It is an attempt to find out those parasites which are particularly affecting these commercially important fishes.

Key words: Cestodes, Nematodes, Trematodes, fishes, infestation.

INTRODUCTION

Fishes are considered as good source of income and government of many countries promote fish farming and provides financial support to their citizens for the production. As fishes are very rich source of protein diet, many doctors also prescribe their patient to consume fish instead of red meat. Fish meat contains ample amount omega-3 fatty acids, vitamin D, vitamin B2, phosphorus, calcium, iron, magnesium, iodine, zinc and potassium.[1] It contains high quality protein with low fat, which keeps our heart and brain healthy. Fishes provide around one-sixth of worlds total protein.[2] Some studies prove that if people consume fish at least once a week, they are less likely to die of heart disease.[3] Fish have had a role in culture for centuries, religious symbols, serving as deities and many more.

Although much smaller than the host, the parasite can cause significant damage to its host. So, parasites are not passive organisms, merely 'riding' with their hosts; To be classified as a parasite, a normal organism must meet criteria that include the fact that its presence has a negative effect on its host.[4] Fish serve as host for parasites as they very diverse. Some parasites usually use fish as their intermediate host. They were found mainly in the intestine of fishes.

*Corresponding author :
Phone : 9006991000
E-mail : prf.arunkumar@gmail.com

MATERIALS & METHODS

Collection of the specimens

For the current study 160 species of commercially important fishes were taken, 20 species each of *Catla catla* (F. Hamilton, 1822), *Labeo rohita* (F. Hamilton, 1822), *Channa punctatus* (Bloch, 1793), *Clarias batrachus* (Linnaeus, 1758), *Cirrhinus mrigala* (Hamilton, 1822), *Heteropneustes fossilis* (Bloch, 1794), *Mystus vittatus* (Bloch, 1794), *Wallago attu* (Bloch & Schneider, 1801). Fishes were purchased from the local waterbodies with the help of fisherman. We have tried to purchase the fish of equal shape and size. Fishes were brought to the laboratory. Total length and weight of the fishes were measured in the laboratory. Their fins and gills were separated and were dissected from the ventral side. The body cavity, spleen, stomach, kidney, liver, intestine etc. all were searched for the presence of parasites. The gastrointestinal tract, the rectum to the esophagus were opened longitudinally and examined carefully.[5]

Table 1. Showing cestode parasites in different host along wwith their habitat

Sl. No.	Host	Cestode Parasites	Habitat
1.	*Catla catla* F. Hamilton, 1822	*Ligula intestinalis L.*	Intestine
2.	*Heteropneustes fossilis* Bloch, 1794	*Crecentovitus biloculus* Murhar 1963	Intestine
3.	*Heteropneustes fossilis* Bloch, 1794	*Djombangia penetrans* Bovien, 1926	Intestine
4.	*Heteropneustes fossilis* Bloch, 1794	*Procamallanus spiculogubernaculus* Agarwal, 1958	Intestine
5.	*Channa punctatus* Bloch, 1793	*Senga raoi* Majid *et al.* 1984	Intestine
6.	*Channa punctatus* Bloch, 1793	*Senga lucknowensis* Johri, 1956	Intestine
7.	*Channa punctatus* Bloch, 1793	*Gangesia paithanensis* Kadam *et al.* 1983	Intestine
8.	*Channa punctatus* Bloch, 1793	*Polyoncobothrium srivastavai* Pande *et al.* 2006	Intestine
9.	*Clarias batrachus* Linnaeus, 1758	*Lytocestus clariae* Tandon, Chakravarty and Das, 2005	Intestine
10.	*Clarias batrachus* Linnaeus, 1758	*Lytocestus longicollis* Ramadevi, 1973	Intestine
11.	*Clarias batrachus* Linnaeus, 1758	*Lytocestus birmanicus* Lynsdale, 1956	Intestine
12.	*Clarias batrachus* Linnaeus, 1758	*Djombangia penetrans* Bovien, 1926	Intestine
13.	*Cirrhinus mrigala* Hamilton, 1822	*Djombangia penetrans* Bovien, 1926	Intestine
14.	*Mystus vittatus* Bloch, 1794	*Lytocestus indicus* (Moghe, 1925) Woodland, 1926	Intestine
15.	*Wallago attu* Bloch & Schneider, 1801	*Gangesia punjabensis*	Intestine
16.	*Wallago attu* Bloch & Schneider, 1801	*Gangesia bengalensis*	Intestine
17.	*Wallago attu* Bloch & Schneider, 1801	*Gangesia vachai*	Intestine
18.	*Labeo rohita* F. Hamilton, 1822	No cestodes	