for further analysis of benthic fauna in the floodplain lakes of this region.

ACKNOWLEDGEMENT

The authors are thankful to the Head, Department of Zoology, B.N. Mandal University, Madhepura, Bihar for providing all the facilities and support. The authors are especially indebted to Dr. A.K. Singh, Professor of Zoology, Patliputra University, Patna, Bihar for his comment and suggestion.

REFERENCES

1. **Odum. E. P. 1971.** Fundamentals of Ecology, 3rd W.B. Saunders Company. Philadelphia, pp. 574.
2. **Reice, S.R. and Wohlenberg, M. 1993.** Monitoring freshwater benthic macro invertebrates and benthic processes: measurement for assessment of ecosystem health. pp. 287-305 in D.M. Rosenberg and V.H.Resh, editors. Freshwater biomonitoring and benthic macro invertebrates. *Chapman & Hall, New York, USA*.
3. **Singh, A.K., Kumari, R. and Kumar, A. 2019.** Diversity and composition of macro-invertebrate in floodplain lakes of North Bihar, India. *Int. J. Fish. Aquat.Stud*., **7(5):** 106-112.
4. **Brickhurst, R.O. 1975.** "Oligochaeta," in Keys to the Water Quality Indicative Organisms of the South Eastern United States, F. K. Parrish, Ed., pp. 69-85, United States Environmental Protection Agency, Office of Research and Development, Environmental Monitoring and Support Laboratory, Cincinnati, Ohio, USA.
5. **Stoffels, R.J., Clarke, K.R. and Closs, G.P. 2005.** Spatial scale and benthic community organization in the littoral zones of large oligotrophic lakes: potential for cross-scale interations. *Freshwater Biology*, **50:** 1131-1145.
6. **Rosenberg, D.M. and Resh, V.H. 1993**. Fresh Water Biomonitoring and Benthic Macro-invertebrates. Chapman and Hall, Inc., New York,pp.820.
7. **Adeogun, A.O. and Fafioye, O.O. 2011.** Impact of effluents on water quality and benthic macro invertebrate fauna of Awba stream and reservoir. *J. Appl. SCI. Environ. Manag*., **15(1):** 105 -113.
8. **Vass, K.K., 2006.** Sustainable fisheries and environmental concerns of floodplain wetlands in India. *Intenal. J. Eco. Environ. Sci*., **32:** 49-62.
9. **Yadava, Y.S., Kolekar, V., Singh R.K. and Chaudhury, M. 1984.** Studies on the macro-benthic fauna of Dinghalibeel (Asam). *Prac, Nat. Acad. Sci. India.* **54(B):** 179-186.
10. **Choudhary, S., Kumar, U. and Kumar, Utpal 2013.** Seasonal changes in the standing crop of macro-invertebrates (Macro-zoobenthos) population in the wetlands (Chaurs) of North Bihar (Begusarai District). *Fishing Chimes*, **33(8):** 72-75.
11. **Malla, S. and Rout, S.K. 2015.** Assessment of the ecological status of Mathura Beel, a flood Plain wetland in Nadia, West Bengal, India by using benthic macro-invertebrates. *NeBIO*, **6(3):** 32-38.
12. **Sharma, G., Dey, A. and Sardana, M. 2011.** Molluscan Diversity. In: Faunal Diversity of Chaurs of North Bihar. *Zool. Surv. India, Wetland Ecosystem Series.* **14:** 21-47.
13. **Sugunan, V.V. and Bhattacharjya, B.K. 2000.** Ecology and fisheries of beels in Assam. Bulletin No.104, *Central Inland Fisheries Research Institute (CIFRI)*, pp. 65.
14. **Edmondson, W.T. 1959.** Freshwater Biology. John Wiley & Sons, New York, USA, pp. 525.
15. **Pennak, R.W. 1978.** Freshwater Invertebrates of the United States. 2nd Ed. John Wiley & Sons, New York, pp. 803.
16. **Tonapi, G.T., 1980.** Fresh Water Animals of India: An Ecological Approach. Oxford & IBH Publishing Co., New Delhi, pp. 341.
17. **Adoni, A.D., Joshi, D.G., Gosh, K., Chourasia, S.K., Vaishya, A.K., Yadav, M. and Verma, H.G., 1985.** Work book on limnology. Pratibha Publishers. C-10 Gour Nagar, Sagar, India**.** pp. 216.
18. **SubbaRao, N. V. 1989.** Handbook Freshwater Molluscs of India. Zoological Survey of India, Calcutta. Edited by the Director, ZSI, pp. 289.

19. **Jhingran, V.G., Natarajan, A.V., Banerjee, S.N. and David, A. 1969.** Methodology on reservoir fisheries investigation in India. *Bull. Cent. Inland Fish. Res. Inst., Barrackpore*, **12:**109 pp.

20. **Khan, R.A. 2002.** The ecology and faunal diversity of two floodplain Ox-bow lakes of South-Eastern West Bengal. *Records of the Zoological Survey of India, Occasional Paper No.* **195:** 1-57.

21. **Sugunan, V.V., Vinci, G.K., B.K. Bhattacharjya, B.K. and Hassan, M.A. 2000.** Ecology and fisheries of beels in West Bengal. Bull. No.103, Central Inland Capture Fisheries Research Institute (ICAR), Barrackpore, West Bengal, pp. 53.

22. **Sharma, R.K. and Agrawal, N. 2011.** Molluscan fauna of Surha Tal, District- Ballia (U. P.), India. *J. Appl. Nat. Sci.*, **3(2):** 295-297.

23. **Patial, P., Hassan, M.A. and A.P. Mishra, A.P. 2015.** Macrobenthic diversity during pre and post drought period of a floodplain wetland in Vaishali district of Bihar. *Int. J. Appl. Biol. Pharm*. **6(2):** 294-298.

24. **Singh, S. and Sharma, R.C. 2020.** Monitoring of benthic macro invertebrates as bio- indicator for assessing the health of the high altitude wetland Dodi Tal, Garhwal Himalaya, India. *Biodiversity Int. J.,* **4(4):** 164-173.

25. **Shyam Sunder and K.K. Vass, 1988.** Seasonal dynamics of benthos in some Kashmir lake. *Proc. Nat. Acad. Sci., India*, **5(11) B**: 193-203.

26. **Soldner, M., Stephen, I., Ramos, L., Angus, R., Wells, N.C., Grosso, A. and Crane, M., 2004.** Relationship between macroinvertebrate fauna and environmental variables in small streams of the Dominican Republic. *Water* Research, **38(4):** 863-874.

27. **Swayne, H., Day, M. and Wetzel, M.J. 2004.** *Limnodrilus hoffmeisteri* (Annelida: Oligochaeta: Tubificidae) in Pop's Cave, Wisconsin, USA. *J. Caves Karst Stud*., **66(1):** 28-31.

28. **Nesemann, H., Sharma, G. and Sinha, R.K. 2004.** Aquatic Annelida (Polychaeta, Oligochaeta, Hirudinea) of the Ganga River and adjacent water bodies in Patna (India, Bihar), with description of a new leech species (Family Salifidae). *Ann.Naturhist. Mus. Wien*., **105 B:** 139-187.

29. **Lenat, D.R. 1993.** A biotic index for the southeastern United States: derivation and list of tolerance values, with criteria for assigning water-quality ratings. *J. North Am. Benthol. Soc*., **12:** 279-290.

30. **Armitage, P., Cranston, P.S. and Pinder, L.C.V. 1995.** The Chironomidae: The biology and ecology of non-biting midges. Chapman and Hall, London, pp. 572.

31. **Clemente, J.M., Mazzeo, N., Gorga, J. and Maerhoff, M. 2005.** Succession and collapse of macrozoobenthos in a subtropical hypertrophic lake under restoration (lake Rodo, Uruguay). *Aquat.Ecol.,* **39:** 455-464.

32. **Mason, C.F. 2002.** Biology of Freshwater Pollution. 4th Edn., Prentice Hall London, pp. 400.

33. **Mophin-Kani, K. and Murugesan, A.G. 2014.** Assessment of river water quality using macro-invertebrate organisms as pollution indicators of Tamirabarani River basin, Tamil Nadu, India. *International Journal of Environmental Protection,* **4(1):** 1-14.

34. **Paine, G.H. and Gaufin, A.R. 1956.** Aquatic Diptera as indicators of pollution in a midwestern stream. *Ohio J. Sci.,***56:** 291-304.

35. **Singh, A.K. 1995.** Effects of sewage on physic-chemical characteristics and macro-zoobenthic community of the river Ganga at Patna, India. *Proc. Nat. Acad. Sci., India*, **65 B (3):** 315-324.

36. **Satyamurti, T.S. 1960.** Land and Freshwater Mollusca. *Bull. Madras Govt. Museum*, pp. 177.

37. **Singh, A.K. 1997.** Abundance of macrozoobenthic organisms in relation to physico-chemical characteristics of river Ganga at Patna (Bihar), India. *J. Environ. Biol.,***18(2):**103-110.

38. **Kumar, A. and Bohra, C. 1999.** Gastropoids as indicators of the pollution status of some wetlands in SantalPargana, Bihar, India. *Indian J. Envir. Ecol*., **2:** 83-87.

***

Biospectra : ISSN: 0973-7057, Vol. 17(1), March, 2022, pp. 175-178
An International Biannual Refereed Journal of Life Sciences

36(AS).
Animal Science

ISSN : 0973-7057

Int. Database Index: 616 www.mjl.clarivate.com

Erratic changes in an air breathing catfish *Heteropneustes fossilis* due to effect of stress

Deepa Kumari[a] & Arun Kumar[b]*

[a]Department of Zoology, T. P College, B. N. M. University, Madhepura, Bihar, India

[b]University Department of Zoology, B. N. M. University, Madhepura, Bihar, India

Received : 15th February, 2022 ; Revised : 16th March, 2022

Abstract- Fish and fish products provide low fat resources and are very healthy substitute of red meat. Fish industry is growing very rapidly, it provides a good source of income. Hence, it is necessary to produce healthy quality of fishes. Fishes get affected by many environmental factors such as temperature, humidity, pH etc. Proper handling of the water body is also necessary for the production. In the current study, the effect of above factors and the erratic changes in air breathing catfish *Heteropneustes fossilis* were studied. The effect of these stressors were studied for a period of two weeks. Subsequent decrease in haemoglobin level and muscle glycogen level was observed. Glycogen content was decreased by 57% when exposed to temperature stress. Haemoglobin level is in relation with erythrocytes count. These erratic changes effects the growth of *Heteropneustes fossilis* which adversely effects the production.

Key words: Erratic changes, Haemoglobin, erythrocytes, *Heteropneustes fossilis*

INTRODUCTION

Fishes are good source of income for human. The fishes are very much dependent on the natural source to feed themselves. They are part of moderate and important food supply in the world.[1] In past these products are thought to be unlimited but now it is realized that these products are finite and in knife edge with the physical chemical and biological factors of the aquatic environment. Fishes are considered as renewable source of income, but some factors such as overfishing pollution, overfishing, alteration in environment has depleted fish production.[2] Fishes are important recourses for human especially as food. Now a day's fish production has become one of the best business worldwide. Fish industry is growing rapidly day by day therefore it is necessary to produce healthy quality of fishes. One of the main obstruction in fish farming is their growth and survival which are effected by many factors. In the current paper we are discussing the erratic changes due to role of stress in the production of fishes. Changes in fish are not completely natural,[3] what precautions should be taken by those who keep fish, is very important. We have taken a number of species of catfish *Heteropneustes fossilis* for this investigation. The erratic changes in *Heteropneustes fossilis* is monitored for a period of two weeks. *Heteropneustes fossilis* are commonly called as singhi and are found in freshwater. It is mainly found in freshwater ditches, marsh lands, swamps, ponds etc. It usually breeds

*Corresponding author :
Phone : 9006991000
E-mail : prf.arunkumar@gmail.com

in monsoon and in constricted water. It is omnivorous in nature. This catfish emanates poison from a gland on its pectoral fin spine to deliver a extremely painful sting to humans. It can tolerate slightly brackish water. It is in great demand due to its medicinal value also.[4-6]

MATERIALS & METHODS

Study area

Madhepura is one among the thirty-eight districts of Bihar situated in 25.920°N latitude and 86.792°E longitudes. The district is 29th largest in terms of area, whereas it is at 34th place in terms of population. The maximum temperature of this district ranges from 35°C to 41°C and the minimum temperature ranges between 7°C to 9°C. The average rain fall in this area is around 1300mm. The district is blessed by a river Kosi, which is an important source of water for agriculture, household use, fish culture etc. The district got its name because it stands at the centre of Kosi ravine, due to which it is termed as Madhyapura (centrally situated) and in due course transformed as Madhepura.

Samples collection

Fifty adult live specimens of *Heteropneustes fossilis* irrespective of sex consideration were collected from the local market of Madhepura. Tanks were maintained at the normal room temperature. The fishes were brought and kept in tanks.The pH of the tanks was maintained near neutral. The pH above and below a certain threshold level is deemed dangerous for the survival of the fishes. The humidity of the lab was also optimum. A good supply of phytoplankton and zooplankton was provided in the tanks from time to time, to ensure proper nutrition to the fishes. Special care was taken while handling the live fishes as their sting can be very painful. It was seen that the tanks donot overcrowd avoiding turbidity. Turbidity usually decreases the photosynthesis rate.

Methods

The fishes were kept in tanks at normal room temperature i.e., 28°C. Simple trial and error methods were used to see the affect of stress such as erratic increase or decrease in temperature, pH,humidity, turbidity, starvation, etc. on fishes. The duration of the experiment was 2 weeks.

Temperature

The tanks were maintained at a normal temperature of 28°C for the first two days of the experiment. This allowed the fishes to get acclimatized or adapted to the new surroundings. After 2 days the temperature of the tank was increasedby 20 with the help of a blower. The fishes were observed at this stressful temperature for 24 hours. Next readings were taken at 48 hours and 96 hours respectively. The movement of fishes both upwards and downwards was seen.

pH

The pH of the tank was kept near neutral for the first two days. The pH of the tank was naturally increased by adding baking soda. The pH increased from around 7 to about 8.3. The fishes were observed for next 24, 48 and 96 hours.

The pH can also be lowered from 7.5 to 6.5 by adding white vinegar to the tank.

Humidity

The humidity on the tank was increased using a vaporizer. The vaporizer locally increases the humidity.

Turbidity

Turbidity affects the amount of sunlight that goes to the bottom of a water body. More turbid water means less sunlight that reaches the water. Less sunlight means less photosynthesis which eventually leads to less food, therefore starvation. Turbidty was increased in the tanks by increasing the number floating plants that cover the surface. The activity of the fish in response to low light was observed.

Starvation

After initial feeding of 2 days, the food provided was minimized by 50%. For the next 24 hours the fishes were observed. The next 48 hours the fishes were provided with only 25% of the food, followed by absolute starvation in the next 96 hours. The metabolic activities were observed carefully.

RESULTS & DISCUSSION

It was seen that introduction of stress in any form be it temperature, pH, humidity, turbidity or starvation leads to restlessness in the initial period. However, if the stress is continued, the fish slowly gets accustomed to the change. If, however, the changes are too erratic the fish's metabolic state is compromised and it slowly settles down at the bottom of the tank. The respiration rate slows down, slowing down all other activities requiring rapid energy as well.