institutional level. Plant should not be collected and natural homes of orchids should not be destroyed without the permission of the local authorities. The market of Orchids can increase throughout the World.



REFERENCES

1. **Kutumbiah, P., 1999.** *Ancient Indian Medicine*. Orient Blackswan.
2. **Bulpitt, C.J., 2005.** The uses and misuses of orchids in medicine. *QJM: An International Journal of Medicine*. **98(9):**625-631.
3. **Abraham, A. & Vatsala, P.9 1981.** Introduction to Orchids ith Illustration and description of 150 South Indian Orchids. TBGRI, Trivandrum, Kerala , pp 533.
4. **Kamemoto, H., Kobayashi, R.S. Amore, T.D. 1989.** Evaluation of 16 seed propagated amphidiploid Dendrobium progenies. Research Extension Series No. 105. Hawaii Institute of Tropical Agriculture and Human Resources, Hawaii, United States of America, p. 127.
5. **Bobisud, C.A. & Kamemoto, H. 1982.** Selection and inbreeding in apmhidiploid Dendrobium (Orchidaceae). *J. Amer. Soc. Hort. Sci*., **107:** 1024-1027.
6. **Singh, F. 1984.** *Cymbidium munronianum*. The Fragrant Cymbidium. *Orchid Reviews*. **92:** 231.
7. **Rogersen, W.P. 1991.** Hybridization of white paphiopedilums, Part-I, *Am. Orchid Soc. Bull*. **60(7):**673-682.
8. **Chen, W.H., Fu, Y.M., Hsieh, R.M., Wu, C.C., Chyou, M.S. and Tsai, W.T. 1995.** Modern breeding in Phalaenopsis orchid. *Taiwan Sugar*. **42(3):** 17-22.
9. **Davidson, B. 1994.** Dendrobium breeding trends. *American Orchids Society Bulletin*. **63:**638-645
10. **Mercy, S.T. and Dale, B. 1997.** Orchids. St. Josephs Press, Thiruvananthapuram, Kerala, India, p. 132.
11. **Fuch, R.F. 1997.** Fabulous Vandaceous intergenerics. *Orchids,* **66:** 350-357.
12. **Kumar, K Madhu and Sheela, V.I. (2007).** Status of breeding in Orchids –A review. *Journal of Ornamental Horticulture*. **10:**199-208.
13. **Bose T.K. and S.K. Battacharjee. 1980.** Orchids in India Naya Parkas Publishers, Calcutta, West Bengal, India.

ADDITIONAL REFERENCES

14. **Singh Foza. 1990.** Indian orchid. *Indian horticulture*. **35(1):**14-15

15. **Taylor, S. 2009.** How to make an Orhid tree.

16. **Friend, GM.R. 2004.** Growing in Orchids in Your Garden Timber press, USA.

17. **Bhattacharjee, S.K. & Das, S.P. 2008.** In: Orchids Botany Breeding, Cultivation, Uses and Post harvest Management. Aavishkar Publishers, Distributers, Jaipur, India, 396 Pb.

18. **De, L.C. 2011.** 'Value Addition in Flower and Orchids' pp.294, Published by New India Publishing Agency, Pitampura, New Delhi-110088.

19. **De. L.C.,Chhetri, Geetamani and Medhi, R.P. 2013.** Orchids –A wonderful Crop for Diversification. *J.Orchid Soc. India*, **279:** 1-20;1-8.

20. **Hajra, P.K & De. Aparajita. 2010.** Phyto-geographic analysis of Orchid flora. J.Orchid soc. India, **24(1/2):**43-46.

21. **Liu H., Chang, L.F., Y.B., Chen B.S., Wang, Z., S. & Gu, H.Y. 2010.** Potential challenges of climate change to orchid conservation in a wild orchid hotspot in south-western China. *Bot. Rev*.,**76:**174-192.

22. **Lopez, R.G. and E.S.Runkle. 2005.** Environmental physiology of growth and flowering orchids. *Hort. Science.* **40(7):**1969-73

23. **Rao A.N. 2004.** Medicinal orchid wealth of Arunachal Pradesh. *Newsletter of Envis Node on Indian Medicinal plant*, **1(2):**1-5.

***

Biospectra : ISSN: 0973-7057, Vol. 17(1), March, 2022, pp. 183-186
An International Biannual Refereed Journal of Life Sciences

38(AS).
Animal Science

ISSN : 0973-7057

Int. Database Index: 616 www.mjl.clarivate.com

Studies on bioaccumulation of heavy metals in gills and muscles of freshwater edible fish species *Clarias batrachus* and *Labeo rohita* of Bara Talab, Bundu, Jharkhand

Shiny E. C. Kachhap* & Anisha Raj Bhengra
Department of Zoology, Marwari College, Ranchi University, Ranchi, Jharkhand, India

Received : 19th February, 2022 ; Revised : 19th March, 2022

Abstract- Agricultural, domestic wastes and industrial discharges have contributed as source of inorganic pollutants in freshwater bodies, that are also a home for many species of edible freshwater fishes. Heavy metals Zinc and Copper are common inorganic components bioaccumulating gradually in different body parts of the fishes, particularly gills and muscles. Consumption of these fishes by people of nearby areas lead to bioaccumulation of toxic heavy metals in the body. In the present study freshwater pond Bara Talab, Bundu, Jharkhand was selected, and bioaccumulation of toxic heavy metals in gills and muscles of two freshwater edible fish *Clarias batrachus* and *Labeo rohita* were studied. Concentration of Zinc in gills of *Clarias batrachus* and *Labeo rohita* was found to be 2.69±0.8 µg/g and 1.8 ±0.6 µg/g respectively, whereas, concentration of Copper was found to be 1.28 ±0.15 µg/g and 0.48 ±0.09 µg/g respectively. Concentration of Zinc in muscles of *C. batrachus* and *L. rohita* was 1.4 ±0.5 µg/g and 0.8 ±0.04 µg/g respectively. Gills showed higher bioaccumulation of both Cu and Zn heavy metal. This result showed that due to deposition of agricultural and domestic waste in Bara Talab, Bundu, freshwater edible fishes are facing the degradation of health as a result of bioaccumulation of toxic heavy metals Zn and Cu.

Key words: Bioaccumulation, *Clarias batrachus, Labeo rohita,* Whatman's filter paper, ANOVA

INTRODUCTION

The term bioaccumulation is defined as uptake, storage, and accumulation of organic and inorganic contaminants by organisms from their environment.[1] Bioaccumulation therefore results from complex interactions between various routes of uptake, excretion, passive release, and metabolization.

For fish, the bioaccumulation process includes two routes of uptake: aqueous uptake of water-borne chemicals and dietary uptake by ingestion of contaminated food particles. Many inorganic pollutants involving toxic metals and different types of salts cause bioaccumulation, that are sourced by agriculture and domestic wastes and industrial discharges. A heavy metal can be defined as a naturally occurring element that has a high atomic weight and high density which is five times greater than that of water.[2] Due to their toxic nature and nature to accumulate in the environment, heavy metals have received a paramount attention among all the pollutants. Usually heavy metals are present in trace amounts in natural waters but many of them are toxic even at very low concentrations.[2] Heavy metals contamination may have devastating effects on the ecological balance of the recipient environment and a diversity of aquatic

*Corresponding author :
Phone : 8051188762
E-mail : shiny.eliza89@gmail.com

organisms. Due to their toxicity and potential bioaccumulation in many aquatic species heavy metals have gained considerable attention in the past decades. High metal concentration in gills reflects high metal concentration in dissolved phase. Zn, although is important for synthesis of important enzymes formation and stabilization of large amount of proteins, it becomes toxic when present in high concentration. Cu also causes Mense disease, Wilson disease, and kidney damage when present in high quantity.

In the present study, bioaccumulation of two heavy metals Zn and Cu in gills and muscles of two important edible freshwater fish species *Clarias batrachus* and *Labeo rohita* found in Bara Talab, Bundu, Jharkhand, was quantitatively studied.

MATERIALS & METHOD

Selection of study area-

In this project Bundu lake, was selected for study because it is widely spread out with abundant flora and fauna. It is a huge lake where freshwater fishes like *Labeo rohita* and *Clarias batrachus* inhabit, other species like *Catla catla* is also found. This was selected for the study mainly because fishes of this lake are brought to the city of Ranchi and are widely sold out. This lake is highly polluted and is vastly covered with water hyacinth. Water hyacinth degrades the water quality and is known to reduce oxygen levels in water bodies.



Fig. 1- partially decomposed plastic pollutants thrown in Bara Talab, Bundu



Fig. 2- Discharge from drains into Bara Talab, Bundu

Fish and tissue collection –

10 specimen of *Clarias batrachus* and *Labeo rohita* each were purchased from the fishermen directly from the site and were transferred to ice box and taken to the laboratory for further assessment. After complete cleaning and removal of scales, fishes were dissected to obtain their gills, accessory respiratory organs and muscles to study the bioaccumulation of heavy metals.

Digestion of tissues-

The tissues were digested following the protocol of Chindah *et al.* (2009)[3]. The tissues were dried in incubator to obtain a constant dry weight 10 ml of concentrated HCl and 1 ml of concentrated HNO_3 was taken in a beaker, 4 g of dry tissue was weighed, ground and added to the beaker gradually. 50 ml of deionized water was added. The mixture was heated in a steam bath to obtain a thick yellow liquid and allowed to cool. The mixture was then filtered using Whatman's filter paper and filtrate obtained was made upto 100 ml using distilled water. Filtrate was stored in plastic bottles and analyzed using atomic absorption spectrophotometer.

Statistical analysis-

All values were expressed as mean of 10 fishes. The difference obtained in the analysis of concentration of metals between two different species were analysed by ANOVA. The difference obtained in the metal concentration in different organs of two different species were statistically analysed by Student's t test to test the significance.

RESULT & DISCUSSION

Table 1- Comparison of concentration of Zinc in gills and muscles of *Clarias batrachus* and *Labeo rohita*

Fish species	Concentration of Zn in Gills (µg/g)	Concentration of Zn in Muscles (µg/g)
Clarias batrachus	2.69±0.8	1.4±0.5*
Labeo rohita	1.8±0.6	0.8±0.04

*=significant at 5% level

Table 2- Comparison of concentration of Copper in gills and muscles of *Clarias batrachus* and *Labeo rohita*

Fish species	Concentration of Cu in Gills (µg/g)	Concentration of Cu in Muscles (µg/g)
Clarias batrachus	1.28±0.15***	0.2±0.08*
Labeo rohita	0.48±0.09	0.08±0.008

*=significant at 5% level