Biospectra : ISSN: 0973-7057, Vol. 17(1), March, 2022, pp. 41-44 **8(AS).**
An International Biannual Refereed Journal of Life Sciences ***Animal Science***

ISSN : 0973-7057

Int. Database Index: 616 www.mjl.clarivate.com

Chemical communication and methods of self-therapy in the honey bee

Asha Ram Meena*
Department of Zoology, University College of Science (Mohanlal Sukhadia University) Udaipur, Rajasthan, India

Received : 20th October, 2021 ; Revised : 27th December, 2021

Abstract- A critical area of physiology of the honey bee is chemical association among individuals and castes in the swarm which balance of its up rightness and limit. The uncommonly friendly interconnection of honeybee is interceded through pheromones. We here discuss the effect of both self-produced organ discharges and foraged hive item on colony health. Explicit honey bee health improving pursuit of honeybee items ought to plainly be recognized from the impact of unblemished nourishment guaranteeing the fundamental resistant ability of honey bee. We should see how the individual working bee with its incredible identification detects consolidates possibly accessible compounds.

Key words: Honey bee, physiology, chemical, colony, health

INTRODUCTION

Honey bee (*Apis mellifera* Linn.) colonies regularly have an enormous number of laborers coming to 15000 to 60,000 individuals of honey bees.[1] Such enormous numbers of honey bees require a lot of food to support their endurance and improvement. Adequate nutrition can support the advancement of honey bee colonies.[2]

Animals incorporating insects search food to get fitting supplements for their food which incorporate development, advancement and proliferation on account of social insects; it isn't simply to fulfill the necessity of people however the scavengers gather the nourishing assets likewise for different individuals living in the hive.[3]

Social resistance sums up all antiparasitic colony level components, not simply decreasing any parasite admission into the colonies yet additionally lessening the speed and transmission among people and colonies.[4] This came while a wondering, given the incredible intra-colonial circumstances for microbes and the evident requirement for control. Nonetheless, colony resistance is administered by individual invulnerable skill as well as by outside safe guard.[5]

Nutrient adjusting is a multiplex process where creatures coordinate taste prompts with post-ingestive data about food quality to acquire ideal nourishment.[6] How rapidly creatures learn signals related with suitable sustenance's. Grasshoppers are bound to figure out how to connect smells and tastes with protein in food assuming they have protein insufficient.[7]

Primary pheromones are proficient means for keeping with social concordance in the colony and their belongings are significant. These pheromones act by influencing the physiology of the beneficiaries with an ensuring change in their way of behaving.[8]

*Corresponding author :
Phone : 6375778159
E-mail : meena05ar@gmail.com

CHEMICAL COMMUNICATION IN THE HONEYBEE

The most bountiful queen mandibular organ pheromone part of keto-2(E) decenoic corrosive (q-ODA) and two fragment parts, 4-hydroxy, 3-hydroxy phenyl ethanol (HVA) and methyl 1p-hydroxy benzoate (HOB) are comparatively communicated.[9] Discharge from the Dufour organ is position controlled.

Martin and Jones (2004)[10] saw that as C28-C38 esters are related with sovereigns and liquor eicosenol is related with non-laying working honeybee. The two esters and eicosenol are synthetically comparable compounds (Both are results of unsaturated fat biosynthesis). 3M2BA may be uniform to certain individuals of Africanized honeybees.[11]

Rhodes *et al.,* (2007)[12] recorded changes in constituent levels from head concentrates of sovereign with expanding age. Non-mated multiday old sovereign had higher normal degree of 9 HDA, 9 ODA and 10 HDA than mated multiday old sovereigns. These outcomes recommend these specific three constituents engage in sexual relations pheromones capacities in the honey bee 9 octa decenoic corrosive decyl decanote from the tarsal organ may likewise take part in the correspondence between genders.

Pheromones functions in the honey bee, 9-octadecenoic acid decyl decanoate from the tergal gland may also participate in communication between sexes. With expanding time of working honey bees, the size of mandibular organ and much 2HPT continuously increments.[13] At the point when the substances are infused into hive, they essentially increment the number of foragers leaving the hive.[14] This recommends that these mixtures might play a pheromonal role in worker prerequisite. An essential quality of eusocial life is a division of work.[15]

Working honeybees for the most part perform various errands in the home (*i.e.,* cell clearing, brood raising brush building, nectar raising for the queen and drones and so on) for the initial three weeks of grown up life and afterward adventure outside to gather food and safeguard the home when they age.[16-18]

In any case, divisions of work in honey bee colonies are unbending.[18, 19] On the connection among wide and working bees a synthetic sign on the outer layer of larvae called brood pheromone is significant.[20,21] A few parts are more every one of the ten individuals compounds show some releaser pheromones impact on adult honey bees.[22] The esters are available in various sums and extents as a component of standing and larvae age.[23]

METHODS OF SELF-THERAPY IN THE HONEY BEE

The additionally influence the retention of synthetic compounds from the climate can serve as pheromones and generally significant for honey bee health, structure a most proficient impediment against the connection to or entrance of the insect cuticle by microorganisms.[24,25] The structure an oily layer on the cuticle which is ceaselessly renewed by glandular emission and keeps organisms and microorganisms from entering the honeybee's body.[26-28] The cuticular hydrocarbon structure fluctuates during individual turn of events and is subject to both the climate and the individual health status.[26] The honey bees cuticular wax layer basically comprises of hydrocarbons, mono and polyester, free unsaturated fat and other polar the substances.[26]

Properties of bee wax have been exhaustively depicted in two monographs by the late.[29,30] Bee wax is discharged by the wax organ and is made out of a perplexing blend of alkanes, alkenes, hydrocarbon 14%, free unsaturated fat 12%, mono esters 35%, diesters 14%, hydroxy monoesters and a few minor constituents (Fatty alchols and hydroxyl diesters).[30,31] The most dynamic compound with upto half of dry weight is melittin. Its high antitoxin and natural action has been read up for a long time in human api-therapy.[31,32] The pollen foragers deposit gathered pollen plates straight for wardly into the capacity cell. The put away pollen is the changed to honey bee bread by in hive honey bees through tactic acid fermentation of pollen blended in with disgorged nectar honey and glandular emission.[2]

Polyfloral pollen further increments endurance rates for stone brood tainted honey bees, as displayed with in vivo tests utilizing honey bee larvae, in contract with monofloral pollen.[33] Protein and amino corrosive quality and amount are the principle measures in regards to the well-being advancing movement of pollen. For instance a new report showed that poly floral pollen isn't really better compared to a monofloral pollen diet of high protein content.[34]

REFERENCES

1. **Southwick, E.E. and Heldmaier, G. 1987.** Temperature control in honey bee colonies. *BioSci.*, **37:**395-399.
2. **Brodschneider R. and Crailsheim K. 2010.** Nutrition and health in honey bees. *Apidologie.* **41:**278-294
3. **Hölldobler, B. and Wilson, E. O. 2009.** The super organism: the beauty, elegance, and strangeness of insect societies. New York: Norton.
4. **Cremer, S., Armitage, S. A. & Schmid-Hempel, P. 2007.** Social immunity. *Current Biology*. **17:**693-702.
5. **Otto, J. C., Mulugu, S., Fridy, P. C., Chiou, S. T., Armbruster, B. N., Ribeiro, A. A. and York, J. D. 2007.** Biochemical analysis of inositol phosphate kinases. *Methods in enzymology*. **434:**171-185.
6. **Simpson, S.J. and Raubenheimer, D. 1997.** Geometric analysis of macronutrient selection in the rat. *Appetite*. **28:** 201-213.
7. **Raubenheimer, D. and Tucker, D. 1997.** Associative learning by locusts: pairing of visual cues with consumption of protein and carbohydrate. *Anim. Behav.* **54:**1449-1459.
8. **Le Conte, Y. and Hefetz, A. 2008.** Primer pheromones in social hymenoptera. *Annual Review of Entomology.* **53:**523–542.
9. **Naumann, K., Winston, M.L., Slessor, K.N., Prestwich, G.D. and Latli, B. 1992.** Intra-nest transmission of aromatic honey bee queen mandibular gland pheromone components: movement as a unit. *Canadian Entomologist.* **124:**917-934.
10. **Martin, S.J. and Jones, G. R. 2004.** Conservation of Bio synthetic pheromone pathways in honeybees *Apis*. *Natur wissen schaften.* **91:**232-236.
11. **Breed, M.D., Guzman-Novoa, E. and Hunt, G.J. 2004.** Defensive behavior of honey bees: Organization, genetics, and comparisons with other bees. *Annual Review of Entomology*. **49:**271-298.
12. **Rhodes, J.W., Lacey, M.J. and Harden, S. 2007.** Changes with age in queen honey bee (*Apis mellifera*) head chemical constituents (Hymenoptera: Apidae). *Sociobiology.* **50:**11-22.
13. **Vallet, A., Cassier, P. and Lensky, Y. 1991.** Ontogeny of the fine structure of the mandibular glands of the honeybee (*Apis mellifera* L.) workers and the pheromonal activity of 2-heptanone. *Journal of Insect Physiology.* **37:**789-804.
14. **Thom, C., Gilley, D.C., Hooper, J. and Esch, H.E. 2007.** The scent of the waggle dance. *Public Library of Science Biology.* **5:**e228.
15. **Pankiw, T. 2004b.** Worker honey bee pheromone regulation of foraging ontogeny. *Natur wissen schaften.* **91:**178-181.
16. **Kolmes, S.A., Winston, M.L. and Fergusson, L.A. 1989.** The division of labor among worker honey bees (Hymenoptera: Apidae): the effects of multiple patrilines. *Journal of the Kansas Entomological Society*. **62:**80-95.
17. **Huang, Z.Y. and Robinson, G.E. 1992.** Honeybee colony integration: Worker-worker interactions mediate hormonally regulated plasticity in division of labor. *Proceedings of the National Academy of Sciences of the USA.* **89:**11726-11729.
18. **Robinson, G.E. and Huang, Z.Y. 1998.** Colony integration in honey bees: genetic, endocrine and social control of division of labor. *Apidologie.* **29:**159-170.
19. **Leoncini, I., Le Conte, Y., Costagliola, G., Plettner, E., Toth, A.L., Wang, M.W., Huang, Z., Becard, J.M., Crauser, D., Slessor, K.N. and Robinson, G.E. 2004.** Regulation of behavioral maturation by a primer pheromone produced by adult worker honey bees. *Proceedings of the National Academy of Sciences* of the USA. **101:**17559-17564.
20. **Le Conte, Y., Sreng, L. and Poitout, S.H. 1995.** Brood pheromone can modulate the feeding behavior of *Apis mellifera* workers (Hymenoptera: Apidae). *Journal of Economic Entomology.* **88:**798-804.
21. **Pankiw, T., Sagili, R.R. and Metz, B.N. 2008.** Brood pheromone effects on colony protein supplement consumption and growth in the honey bee (Hymenoptera: Apidae) in a subtropical winter climate. *Journal of Economic Entomology*. **101:**1749-1755.
22. **Le Conte Y., Mohammedi, A. and Robinson, G. E. 2001.** Primer effects of a brood pheromone on honeybee

behavioural development. *Proceedings of the Royal Society of London B*. **268:**163-168.

23. **Le Conte Y., Sreng, L., Sacher, N., Trouiller, J., Dusticier, G. and Poitout, S. H. 1994/1995.** Chemical recognition of queen cells by honey bee workers *Apis mellifera* (Hymenoptera: Apidae). *Chemoecology.* **5/6:**6-12.

24. **Blomquist, G.J. and Jackson, L.L. 1979.** Chemistry and biochemistry of insect waxes. *Prog. Lipid Res.***17:**319-345

25. **Lockey, K.H. 1988.** Lipids of the insect cuticle: origin, composition and function. *Comp. Biochem. Phys. B.* **89:**595-645.

26. **Blomquist, G.J., Chu, A.J. and Remaley, S. 1980.** Biosynthesis of wax in the honeybee *Apis mellifera* L. *Insect Biochemistry*. **10:**313-321.

27. **Koidsumi, K. 1957.** Antifungal action of cuticular lipids in insects. *Journal of Insect Physiology.***1:**40-51.

28. **Gołębiowski, M., Cerkowniak, M., Boguś, M.I., Włóka, E., Dawgul, M., Kamysz, W. and Stepnowski, P. 2013.** Free fatty acids in the cuticular and internal lipids of *Calliphora vomitoria* and their antimicrobial activity. *Journal of Insect Physiology*. **59:**416-429.

29. **Hepburn, H.R. 1986.** Honeybees and Wax. Springer, Heidelberg.

30. **Hepburn, H.R., Pirk, C.W.W. and Duangphakdee, O. 2014.** Honeybee Nests: Composition, Structure, Function. Springer, Heidelberg.

31. **Mizrahi, A. and Lensky, Y. 1997.** Bee products: Properties, Applications, and Apitherapy. Springer, New York, NY.

32. **Ratcliffe, N.A., Mello, C.B., Garcia, E.S., Butt, T.M. and Azambuja, P. 2011.** Insect natural products and processes: new treatments for human disease. *Insect Biochem.Mol. Biol.*, **41:**747-769.

33. **Foley, K., Fazio, G., Jensen, A.B. and Hughes, W.O.H. 2012.** Nutritional limitation and resistance to opportunistic *Aspergillus* parasites in honey bee larvae. *Journal of Invertebrate Pathology*. **111:** 68-73

34. **Di Pasquale, G., Salignon, M., Le Conte, Y., Belzunces, L.P., Decourtye, A., Kretzschmar, A., Suchail, S., Brunet, J.L. and Alaux, C. 2013.** Influence of pollen nutrition on honey bee health: do pollen quality and diversity matter? *PLoS ONE.* **8:**72016.

***