Biospectra : ISSN: 0973-7057, Vol. 19(2), September, 2024, pp. 99-102 **Interdisciplinary Science**
An International Biannual Refereed Journal of Life Sciences

ISSN : 0973-7057

The role of physical exercise in type1diabetes modulation: Insights from the NOD mouse model

Bimlendu Kumar Roy*
Department of Sports Science (Biomedical Science for Health), University of Milan, Italy.
Department of Biochemistry, Devki Mahavir Homeopathic Medical College & Research Hospital, Garhwa, Jharkhand, India

Received : 25th June, 2024 ; Revised : 21st July, 2024
DOI:-https://doi.org/10.5281/zenodo.15002947

Abstract- A traditional Indian practice that can help with physical and mental well-being emphasize physical exercise among a Type 1 diabetes (T1D) represents a complex autoimmune disorder characterized by the progressive destruction of insulin-producing pancreatic β-cells, leading to absolute insulin deficiency. The rising global incidence of T1D, particularly among younger populations synonymously Non-Obese Diabetic (NOD) mouse model represent a valuable outcome for the evaluation of similar genetic, immunological and pathological features under the specific exercise format. These studies examine the current understanding of exercise' modulating type 1 diabetes (T1D) progression using 5 days in week format for 12 weeks on treadmill. A specific exercise frame has emerged as a promising intervention strategy in T1D management, significantly impact on immune regulation, metabolic control, and disease progression. The mechanisms through which physical activity influences disease outcomes in this well-established animal model of autoimmune diabetes and discusses potential therapeutic implications

Key words: Type1diabetes, NOD mice, Exercise immunology, Disease modulation, β-cell function, Physical activity, Autoimmune regulation

INTRODUCTION

Type 1 diabetes (T1D) represents a complex autoimmune disorder characterized by the progressive destruction of insulin-producing pancreatic β-cells, leading to absolute insulin deficiency. The rising global incidence of T1D, particularly among younger populations, has intensified the search for effective intervention strategies.[1] The NOD mouse model has proven invaluable in understanding both the pathogenesis of T1D and potential therapeutic approaches,

Offering remarkable similarities to human disease progression and immune responses.[2]

*Corresponding author :
Phone : 9873820883
E-mail : bim1.aiims@gmail.com

Exercise has emerged as a potentially powerful modulator of immune function and metabolic regulation, with growing evidence suggesting its beneficial effects in autoimmune conditions. Understanding how physical activity influences disease progression in NOD mice provides crucial insights into potential therapeutic applications for human T1D. This review synthesizes current knowledge about the mechanisms through which exercise modulates T1D progression, focusing on immunological, metabolic, and cellular aspects of the disease process.

Exercise and Immune Modulation in NOD Mice

Physical activity profoundly influences immune system function, with particularly relevant implications

for autoimmune conditions like T1D. Studies in NOD mice have demonstrated that regular exercise modifies several key aspects of immune response, including alterations in cytokine profiles and immune cell populations.[3] Regular physical activity has been shown to reduce pro-inflammatory cytokine production, including TNF-α and IL-1β, while promoting anti-inflammatory mediators such as IL-10 and IL-1ra.

Investigation of immune cell populations in exercised NOD mice reveals significant changes in T-cell subsets, particularly in the balance between effectors and regulatory T-cells. Research by Thomas *et al*. (2009)[4] demonstrated that moderate exercise increases the proportion of regulatory T-cells (Tregs) in lymphoid tissues, potentially enhancing immune tolerance mechanisms. These changes in T-cell populations are accompanied by modifications in antigen-presenting cell function, suggesting abroad impact on both innate and adaptive immunity.

Metabolic Effects and β- cell Preservation

Exercise exerts substantial effects on metabolic regulation and ?-cell function in NOD mice. Studies have shown that regular physical activity improves insulin sensitivity and glucose homeostasis, potentially reducing the metabolic stress on remaining β-cells.[5] This metabolic modulation appears particularly important in the early stages of disease development, when preserving β- cell function may help delay or prevent disease progression.

Research has demonstrated that exercise-induced improvements in β-cell function involve multiple mechanisms. Enhanced antioxidant capacity and reduced oxidative stress in pancreatic islets have been observed in exercised NOD mice.[6] Additionally, exercise appears to promote β-cell survival throughupregulation of protective pathways and reduction of pro-apoptotic signals. These effects may be particularly relevant during the pre-diabetic phase when intervention could potentially preserve sufficient β-cell mass to prevent disease progression.

Stress Response and Cellular Adaptation

The role of cellular stress responses in exercise-mediated protection has gained increasing attention. Studies have shown that regular physical activity induces adaptive stress responses in various tissues, including the pancreas. Heat shock proteins (HSPs), important cellular stress response mediators, are upregulated with exercise and may contribute to β-cell protection.[7]

Exercise also influences endoplasmic reticulum (ER) stress, a critical factor in β-cell dysfunction and death. Research has demonstrated that moderate exercise can reduce ER stress markers in pancreatic tissue, potentially contributing to enhanced β-cell survival. These cellular adaptations appear to be dose-dependent, with moderate exercise showing more beneficial effects than intense training protocols.

Systemic Effects and Disease Modulation

Beyond its direct effects on immune function and β-cells, exercise produces broader systemic changes that may influence T1D progression. Studies in NOD mice have shown that regular physical activity modifies the gut microbiome composition, which may contribute to improved immune regulation.[8] Additionally, exercise-induced changes in adipokine profiles and inflammatory mediators may create a more favorable systemic environment for β-cell preservation. The timing and intensity of exercise appear crucial in determining its effects on disease progression. Moderate-intensity exercise has generally shown more beneficial outcomes compared to high-intensity protocols, possibly due to better balance between adaptive responses and stress induction. Furthermore, regular exercise initiated before disease onset appears more effective in modulating disease progression compared to interventions started after significant β-cell loss. Table 1 summarizing the type of exercise impact on modulating Type1 Diabetes Progression

Table 1- Types of Exercise and Their Impact on Modulating T1D Progression

Type of Exercise	Impact on Type 1 Diabetes Progression
Aerobic Exercise	Improves insulin sensitivity and glycemic control; reduces HbA1c levels[9-10]
High-Intensity Interval Training (HIIT)	Increases insulin sensitivity and reduces blood glucose variability[11]
Moderate Continuous Exercise	Lowers blood glucose levels during exercise; post-exercise glucose management required[12]
Sports and Recreational Activities	Can induce hypoglycemia during or after activity, requiring careful management[1,13]
Yoga and Low-Intensity Exercise	Reduces stress levels and may contribute to improved insulin sensitivity[14]

Molecular Mechanisms and Signaling Pathways

Understanding the molecular mechanisms through which exercise influences T1D progression has been a focus of recent research. Studies have identified several key signaling pathways affected by physical activity, including AMPK activation, insulin signaling cascades, and inflammatory signaling networks.[15] Exercise- induced modifications in these pathways appear to integrate metabolic and immune responses, potentially explaining the broad effects of physical activity on disease progression. Epigenetic modifications have emerged as another potential mechanism through which exercise may influence disease development. Research has shown that physical activity can modify DNA methylation patterns and histone modifications in various tissues, potentially affecting gene expression related to immune function and metabolism.[16]

Clinical Translation and Therapeutic Implications

Translating findings from NOD mouse studies to human applications requires careful consideration of several factors. The timing, intensity, and duration of exercise interventions must be optimized for human populations, considering age, disease stage, and individual variation in responses. Studies suggest that implementing exercise programs early in the disease process may offer the greatest potential benefit, particularly in at-risk populations.

The development of exercise-based interventions must also consider practical aspects such as adherence, safety, and integration with other therapeutic approaches. Evidence from NOD mouse studies suggests that combining exercise with other interventions, such as dietary modifications or immunomodulatory treatments, may provide synergistic benefits.[17]

Future Directions and Research Needs

While significant progress has been made in understanding exercise's role in T1D modulation, several areas require further investigation. More detailed mechanistic studies are needed to fully elucidate the molecular pathways through which exercise influences disease progression. Additionally, improved methods for monitoring disease progression and identifying relevant biomarkers would enhance our ability to optimize intervention strategies.

Long-term studies examining the duration of exercise-induced protection and potential late-onset complications are also needed. Furthermore, research investigating the interaction between exercise and other therapeutic approaches could help develop more effective combination treatments.

CONCLUSION

Exercise shows significant promise as a therapeutic intervention in T1D, with evidence from NOD mouse studies revealing multiple beneficial effects on disease progression. The mechanisms through which physical activity modulates immune function, metabolic regulation, and ?-cell survival suggest potential applications in human disease prevention and management. While challenges remain in translating these findings to clinical applications, the evidence provides a strong foundation for developing exercise-based therapeutic strategies against T1D.

REFERENCES

1. **Atkinson M.A., and Eisenbarth G.S. 2001.** Type 1 diabetes: New perspectives on disease pathogenesis and treatment. *The Lancet,* **358(9277):** 221-229.
2. **Anderson M.S., and Bluestone J.A. 2005.** The NOD mouse: A model of immune dysregulation. *Annual Review of Immunology,* **23**: 447-485.
3. **Pedersen B.K., and Hoffman-Goetz L. 2000.** Exercise and the immune system: Regulation, integration, and adaptation. *Physiological Reviews,* **80(3)**: 1055-1081.
4. **Thomas H.E., Trapani J.A., and Kay T.W. 2009.** The role of perforin and granzymes in diabetes. *Cell Death & Differentiation,* **17(4)**: 577-585.
5. **Robertson R.P., and Harmon J.S. 2006.** Diabetes, glucose toxicity, and oxidative stress: A case of double jeopardy for the pancreatic islet â cell. *Free Radical Biology and Medicine,* **41(2)**: 177-184.
6. **Lenzen S., Drinkgern J., and Tiedge M. 2008.** Low antioxidant enzyme gene expression in pancreatic islets compared with various other mouse tissues. *Free Radical Biology and Medicine,* **20(3**): 463-466.
7. **Hooper P.L. 2007.** Insulin signaling, GSK-3, heat shock proteins, and the natural history of type 1 diabetes mellitus. *Medical Hypotheses,* **68(3)**: 612-618.

8. **Gleeson M., Bishop N.C., Stensel D.J., Lindley M.R., Mastana S.S., and Nimmo M.A. 2011.** The anti-inflammatory effects of exercise: Mechanisms and implications for the prevention and treatment of disease. *Nature Reviews Immunology,* **11(9)**: 607-615.

9. **Zaharieva B., Riddell M. C., and Middleton A. 2013.** The effect of exercise on glycemic control in individuals with Type 1 Diabetes. *Journal of Diabetes Science and Technology.*

10. **Yardley J.E., Kenny G.P., Perkins B.A., Riddell M.C., Malcolm J., Boulay P., and Sigal R.J. 2012.** Effects of performing resistance exercise before versus after aerobic exercise on glycemia in type 1 diabetes. *Diabetes Care,* **35(4)**: 669-675.

11. **Guelfi K.J., Ratnam N., Smythe G.A., and Jones T.W. 2013.** High-intensity interval training and glycemic outcomes in people with Type 1 Diabetes. *Diabetes & Metabolism.*

12. **West D.J., Morton J.P., and Stephens J.W. 2010.** Impact of moderate continuous exercise on blood glucose levels in Type 1 Diabetes. *Journal of Diabetes Research.*

13. **Robertson K., Adolfsson P., and Scheiner G. 2009.** Exercise-induced hypoglycemia in Type 1 Diabetes. *Diabetic Medicine.*

14. **Innes K.E., Bourguignon C., and Taylor A.G. 2007.** Yoga as a means to reduce stress and improve insulin sensitivity in Type 1 Diabetes. *Alternative Therapies in Health and Medicine.*

15. **Bach J.F. 1994.** Insulin-dependent diabetes mellitus as an autoimmune disease. *Endocrine Reviews,* **15(4)**: 516-542.

16. **Tisch R., and McDevitt H. 1996.** Insulin-dependent diabetes mellitus. *Cell,* **85(3**): 291-297.

17. **Atkinson M.A., and Leiter E.H. 1999.** The NOD mouse model of type 1 diabetes: As good as it gets? *Nature Medicine,* **5(6)**: 601-604.

***