

# Supplementary Material

## Contents

<b>1 Supplementary Note 1: Guide on how to understand the reports</b>	<b>2</b>
1.1 Understanding the reports . . . . .	2
1.1.1 Orientation . . . . .	2
1.1.1.1 Toolbar . . . . .	2
1.1.1.2 Main Body . . . . .	4
1.1.1.3 Test Results . . . . .	4
1.1.2 Interpretation . . . . .	4
1.1.2.1 Color . . . . .	4
1.1.2.2 Score . . . . .	5
1.1.3 Paradigms . . . . .	5
1.1.3.1 “Reconstructions” and “Models” . . . . .	5
1.1.3.2 “Lumped” and “Split” Biomass Reaction . . . . .	6
1.1.3.3 “Average” and “Unique” Metabolites . . . . .	6
<b>2 Supplementary Note 2: Validation against experimental data</b>	<b>6</b>
<b>3 Supplementary Methods</b>	<b>9</b>
3.1 Tested models . . . . .	9
3.2 Clustering . . . . .	9
3.3 Test Suite . . . . .	13
3.3.1 Summary of Observations . . . . .	13
3.3.2 Scores . . . . .	13
3.3.3 Independent Section . . . . .	14
3.3.3.1 Consistency . . . . .	14
3.3.3.2 Annotation - Metabolites . . . . .	18
3.3.3.3 Annotation - Reactions . . . . .	33
3.3.3.4 Annotation - Genes . . . . .	46
3.3.3.5 Annotation - SBO Terms . . . . .	60
3.3.4 Specific Section . . . . .	67
3.3.4.1 SBML . . . . .	67
3.3.4.2 Basic Information . . . . .	69
3.3.4.3 Metabolite Information . . . . .	73
3.3.4.4 Reaction Information . . . . .	76
3.3.4.5 Gene-Protein-Reaction (GPR) Association . . . . .	81
3.3.4.6 Biomass . . . . .	83
3.3.4.7 Energy Metabolism . . . . .	88
3.3.4.8 Network Topology . . . . .	90
3.3.4.9 Matrix Conditioning . . . . .	94
<b>Supplementary References</b>	<b>97</b>




Figure S1: Snapshot Report

## 1 Supplementary Note 1: Guide on how to understand the reports

The following text is a verbatim copy modified to work in print taken from memote's documentation at the time of publication. For an updated version please check the latest [memote documentation](#).

### 1.1 Understanding the reports

Memote will return one of four possible outputs. If your preferred workflow is to benchmark one or several genome-scale metabolic models (GSM) memote generates either a snapshot (Figure S1) or a diff report (Figure S2), respectively. For the reconstruction workflow the primary output is a history report (Figure S3). This will only work if the provided input models are formatted correctly in the [systems biology markup language \(SBML\)](#). However, if a provided model is not a valid SBML file, memote composes a report enumerating errors and warnings from the SBML validator the order of appearance. To better understand the output of the error report we refer the reader to this section of the [SBML documentation](#). In this section, we will focus on how to understand the snapshot, diff and history reports.

#### 1.1.1 Orientation

##### 1.1.1.1 Toolbar

In all three reports, the blue toolbar at the top shows (from left to right) the memote logo, a button which expands and collapses all test results, a button which displays the readme and the github icon which links to memote's github page. On the snapshot report, the toolbar will also display the identifier of the tested GEM and a timestamp showing when the test run was initiated.

Independent Section		Specific Section	
Contains tests that are independent of the class of modeled organism, a model's complexity or types of identifiers that are used to describe its components. Parameterization or initialization of the network is not required. See readme for more details.		Covers general statistics and specific aspects of a metabolic network that are not universally applicable. See readme for more details.	
<b>Consistency</b>		<b>Basic Information</b>	
Compared Models		Compared Models	
Stoichiometric Consistency		e_coli_core.xml	e_coli_core_special.xml
100.0%	100.0%	0.69	0.69
Mass Balance		2	2
100.0%	100.0%	72	54
Charge Balance		0.0%	24.3%
100.0%	100.0%	64	54
Metabolite Connectivity		0	0
100.0%	100.0%	72	54
Unbounded Flux In Default Medium		0	0
100.0%	100.0%	7	7
Metabolite Production With Closed Bounds		95	95
98.8%	100.0%	6	6
Metabolite Consumption With Closed Bounds		49	72
100.0%	100.0%	1	1
Sub Total		0	0
89%	92%	25	2
<b>Annotation</b>		Metabolites without Charge	
Compared Models		Metabolites without Formula	
Presence of Metabolite Annotation		e_coli_core.xml	e_coli_core_special.xml
100.0%	100.0%	Medium Components	Medium Components
Metabolite Annotations Per Database		Total Reactions	Total Reactions
pubchem.compound	Info	Reactions without GPR	Reactions without GPR
0.0%	0.0%	Purely Metabolic Reactions	Purely Metabolic Reactions
kegg.compound	100.0%	Purely Metabolic Reactions with Constraints	Purely Metabolic Reactions with Constraints
98.8%	98.3%	Duplicate Reactions	Duplicate Reactions
seed.compound	98.6%	Transport Reactions	Transport Reactions
inchikey	98.1%	Transport Reactions with Constraints	Transport Reactions with Constraints
chebi	0.0%	Fraction of Transport Reactions without GPR	Fraction of Transport Reactions without GPR
hmdb	97.2%	Number of Reversible Oxygen-Containing Reactions	Number of Reversible Oxygen-Containing Reactions
	91.7%	Non-Growth Associated Maintenance Reaction	Non-Growth Associated Maintenance Reaction
	90.7%	Total Genes	Total Genes
		Enzyme Complexes	Enzyme Complexes

Figure S2: Diff Report




Figure S3: History Report

### 1.1.1.2 Main Body

The main body of the reports is divided into an independent section to the left and a specific section to the right.

The tests in the independent section are agnostic of the type of modeled organism, preferred modeling paradigms, the complexity of a genome-scale metabolic model (GEM) or the types of identifiers that are used to describe its components. The tests in this section focus on testing adherence to fundamental principles of constraint-based modeling: mass, charge and stoichiometric balance as well as the presence of annotations. The results in this section can be normalized, and thus enable a comparison of GEMs. The **Score** at the bottom of the page summarises the results to further simplify comparison. While calculating an overall score for this section allows for the quick comparison of any two given models at a glance, we recommend a thorough analysis of all results with respect to the desired use case.

The specific section on the right provides model specific statistics and covers aspects of a metabolic network that can not be normalized without introducing bias. For instance, dedicated quality control of the biomass equation only applies to GEMs which are used to investigate cell growth, i.e., those for which a biomass equation has been generated. Some tests in this section are also influenced by whether the tested GEM represents a prokaryote or a eukaryote. Therefore the results cannot be generalized and direct comparisons ought to take bias into account.

### 1.1.1.3 Test Results

Test results are arranged in rows with the title visible to the left and the result on the right. The result is displayed as white text in a coloured rectangle detailed below in the subsection **Color**.

By default only the minimum information is visible as indicated by an arrow pointing down right of the result. Clicking anywhere in the row will expand the result revealing a description of the concept behind the test, its implementation and a brief summary of the result. In addition, there is a text field which contains plain text representations of Python objects which can be copied and pasted into Python code for follow up procedures.

Some tests carry out one operation on several parameters and therefore deviate slightly from the descriptions above. Expanding the title row reveals only the description, while rows of the individual parameters reveal the text fields.

In the history report, instead of text fields scatterplots show how the respective metrics developed over the commit history for each branch of a repository. By clicking an entry in the legend, it is possible to toggle its visibility in the plot.

## 1.1.2 Interpretation

The variety of constraints-based modeling approaches and differences between various organisms compound the assessment of GSMS. While memote facilitates model assessment it can only do so within limitations. Please bear in mind the diversity of Paradigms that challenge some of memote's results.

### 1.1.2.1 Color

#### Snapshot Report



Figure S4: Snapshot Report Score Gradient



Figure S5: Diff Report Ratio Gradient

Results without highlights are kept in the main blue color of the memote color scheme. Scored results (Figure G1) will be marked with a gradient ranging from red to green denoting a low or a high score respectively:

**Diff Report** The colour in the Diff Report (Figure G2) depends on the ratio of the sample minimum to the sample maximum. Result sets where the sample minimum and the sample maximum are identical will be coloured in the main blue color of the memote color scheme. Result sets where the sample minimum is very small relative to the sample maximum will appear red. This ratio is calculated with as

$$1 - (Min/Max) * 100$$

This is then mapped to the following gradient:

### 1.1.2.2 Score

Each test in the independent section provides a relative measure of completeness with regard to the tested property. The final score is the weighted sum of all individual test results normalized by the maximally achievable score, i.e., all individual results at 100%. Individual tests can be weighted, but it is also possible to apply weighting to entire subsections. Hence the final score is calculated:

$$TotalScore = \frac{\sum_{Subsections} weight_{subsection} \times (\sum_{Tests} weight_{test} \times TestScore)}{MaxScore}$$

Weights for sections and individual tests are indicated by a white number inside a magenta badge. No badge means that the weight defaults to 1.

The subsections “Consistency” and “Annotation - SBO” have weights of 3 and 2, respectively. The test “Stoichiometric Consistency” itself is weighted 3 times stronger than the remaining tests in the “Consistency” subsection. The remaining subsections and tests which cover annotations of metabolites, reactions and genes have weights of 1 (Supplementary Figure G1).

### 1.1.3 Paradigms

#### 1.1.3.1 “Reconstructions” and “Models”

Some authors may publish metabolic networks which are parameterized, ready to run flux balance analysis (FBA), these are referred to simply as ‘models’. Alternatively, others may publish unconstrained metabolic knowledge bases (referred to as ‘reconstructions’), from which several models can be derived by applying different constraints. Both can be encoded in SBML. With having an independent test section, we attempt to make both ‘models’ and ‘reconstructions’ comparable, although a user should be aware that this difference exists and is subject to some discussion. Please note that some tests in the specific section may error for a reconstruction as they require initialization.

### 1.1.3.2 “Lumped” and “Split” Biomass Reaction

There are two basic ways of specifying the biomass composition. The most common is a single lumped reaction containing all biomass precursors. Alternatively, the biomass equation can be split into several reactions each focusing on a different macromolecular component for instance a (1 gDW ash) + b (1 gDW phospholipids) + c (free fatty acids) + d (1 gDW carbohydrates) + e (1 gDW protein) + f (1 gDW RNA) + g (1 gDW DNA) + h (vitamins/cofactors) + x ATP + x H<sub>2</sub>O -> 1 gDCW biomass + x ADP + x H + x Pi. The benefit of either approach depends very much on the use cases which are discussed by the community. Memote employs heuristics to identify the type of biomass which may fail to distinguish edge cases.

### 1.1.3.3 “Average” and “Unique” Metabolites

A metabolite consisting of a fixed core with variable branches such as a membrane lipid is sometimes implemented by averaging over the distribution of individual lipid species. The resulting pseudo-metabolite is assigned an average chemical formula, which requires scaling of stoichiometries of associated reactions to avoid floating point numbers in the chemical formulae. An alternative approach is to implement each species as a distinct metabolite in the model, which increases the total count of reactions. Memote cannot yet distinguish between these paradigms, which means that results in the specific sections that rely on the total number of reactions or scaling of stoichiometric parameters may be biased.

## 2 Supplementary Note 2: Validation against experimental data

To compare model predictions to experimental measurements, a researcher would typically write a short script. The reproducibility of this script may be limited by the original author’s style of writing code, whether the code has been rigorously checked for errors, and whether it is dependent on obsolete libraries. The latter, so called software rot, arises from a lack of active maintenance (Beaulieu-Jones and Greene 2017).

In contrast, with memote researchers may optionally define a configuration file (in YAML format) in which they can set the medium and FBA objective. This file can be used by researchers without prior programming experience. It configures memote to execute clearly defined, formulaic operations, which are unit tested. Lastly, it confers the burden of maintenance to the memote community represented through this consortium. This does not only distribute the necessity for funding onto many shoulders, but also increases the likelihood of the codebase keeping up with advances in its core dependencies, i.e., keeping software rot at bay. The development of the COBRAToolbox (Heirendt et al. 2017) and cobrapy (Ebrahim et al. 2013) are pertinent examples of community projects that operate on a similar strategy. Moreover, frequent versioning ensures that users can return to previous versions to re-run analyses.

Setting up a version-controlled model repository not only allows researchers to publish a ‘default’ unspecific GEM of the investigated organism, but also reproducible instructions on how to obtain a model that is specific to the organism in a defined experimental context including, and validated against the data supporting this context. This formulaic approach of deriving a GEM into a condition-specific form supports Heavner and Price’s (Heavner and Price 2015) call for more transparency and reproducibility in metabolic network reconstruction ([S6](#)).




Figure S6: Experimental tests can be tailored to a specific condition through the use of one or several configuration files (configs). (a) To validate GEMs against experimental data measured in specific conditions, researchers usually write their scripts which constrain the model. This is problematic as scripts can vary a lot and they are, unless actively maintained, susceptible to software rot. (b) With memote, user-defined configuration files replace scripts, which allows the experimental validation of GEMs to be unified and formalized. Bundling the model, configuration files, and experimental data within a version-controlled repository (indicated by the blue asterisk\*) facilitates reproducibility.

### 3 Supplementary Methods

To simplify interpretation, the following figures are grouped by the sections of their corresponding test cases as they appear in a snapshot report. The code that was used to generate the data and figures has been deposited on GitHub <https://github.com/biosustain/memote-meta-study>.

#### 3.1 Tested models

In order to respect the limited resources on the DTU high performance computing infrastructure, we set a maximum time limit for running the memote test suite. This introduced a bias against large models. Additionally, certain models failed the testing procedure. In the following we tabulate the total size of the collections as well as the final number of tested models. The results are shown in Table S1.

#### 3.2 Clustering

In order to perform the clustering analyses, we used all normalized test metrics excluding some particular cases. Excluded are the Sections 3.3.4.2 & 3.3.4.6 because the basic information only contains unnormalized model dimensions and because a biomass formulation is not present in all models. We further removed individual biomass related test cases, as well as the metabolic coverage since that is not properly normalized. Additionally, test cases that contained errors were penalized with the worst metric of one.

To determine the most relevant tests to discriminate between model collections, we built a classifier using a random forest (Breiman 2001) over the collections and normalized test results (0.99 accuracy and 0.01% out-of-bag (OOB) error). Then, the importance of each variable, i.e., test case, was ranked with the Mean Decrease in Accuracy (MDA) (Louppe et al. 2013). This metric measures the total decrease in accuracy, averaged over all trees of the forest, when the value of a given variable is permuted in the OOB samples. Figure S10 represents the 15 most discriminant features on average (see last column) and their independent relevance by collection. The higher the decrease in accuracy, the higher the relative contribution of such a test to differentiate among collections. Thus, the five most discriminant tests are purely metabolic reactions, transport reactions, dead-end metabolites, orphan metabolites, and the presence of a non-growth associated maintenance reaction. Although there is a variable range of importance for each collection, e.g., for CarveMe transport reactions and orphans are more relevant; for Kbase transport reactions; for Ebrahim *et al.* purely metabolic reactions. For a detailed study of the clustering properties, please refer to the *Supplementary Clustering Analysis* notebook.




Figure S7: Depicted are the first two components of a principal components analysis of the normalized test features (metrics).




Figure S8: Depicted are the distances between models in higher order space given by the normalized test features reduced to two dimensions using t-SNE.

Table S1: Number of tested models.

Collection	Number of Models	Tested Models	%
AGORA	818	801	97.9
CarveMe	5587	5511	98.6
Path2Models	2641	2641	100.0
KBase	1637	1632	99.7
BiGG*	36	36	100.0
Ebrahim <i>et al.</i> †	83	80	96.4
OptFlux Models†	100	79	79.0

\* Please note that we removed the large number of *Escherichia coli* strain models from the BiGG collection and only included results from the models iJR904, iAF1260, iJO1366, and iML1515.

† 39 models from these two collections are likely identical based on a filename comparison.




Figure S9: Depicted are the distances between models in higher order space given by the normalized test features reduced to two dimensions using UMAP.




Figure S10: 15 most relevant tests to discriminate among GEM collections, for each collection and the mean. Ranked in decreasing importance according to the *mean decrease in accuracy* metric averaged over all collections (last column), computed over a random forest classification model.

### 3.3 Test Suite

#### 3.3.1 Summary of Observations

- SBO terms are only used by models from KBase and BiGG (Figure [S86](#)).
- Models from Path2Models and Opflux Models are formatted in legacy SBML (< Level 3, Version 1) without FBC package (Figures [S98](#) & [S99](#)).
- Models from the collections of Ebrahim *et al.*, and OptFlux Models are highly variable for many specific tests. Models from automatic reconstruction pipelines (AGORA, CarveMe, Path2Models, and KBase) or the controlled BiGG collection are much more similar within each collection yet still different from each other. This could be due to each collection focusing on a distinct set of taxonomies but could also be related to the algorithms and databases behind each collection (Section [3.3.4.8](#); Figures [S111](#), [S113](#), and [S119](#)).
- On biomass:
  - Only for a minority of models in BiGG, Ebrahim *et al.*, and OptFlux Models memote could not identify a biomass reaction (Figure [S123](#)).
  - A portion of models in the BiGG collections have inconsistent biomass equations followed by OptFlux Models and models in the collection by Ebrahim *et al.*; all models in the CarveMe and Path2Model collections have inconsistent biomass reactions (Figure [S124](#)).
  - Models that cannot be simulated using the default or complete medium exist in Path2Models, BiGG, Ebrahim *et al.*, and OptFlux Models (Figure [S125](#) & [S126](#)).
  - Possible artifacts from automatic reconstruction are present in models from AGORA and KBase that grow despite some biomass precursors being blocked when each precursor is optimized individually in default and complete medium (compare Figures [S127](#) & [S128](#) with [S125](#) & [S126](#)).
- The average fraction of reactions that participate in stoichiometrically-balanced cycles is larger for models from automatic reconstruction pipelines (AGORA, CarveMe, Path2Models, KBase) than for BiGG, Ebrahim *et al.*, and OptfluxModels (Figure [S137](#)). This could be an artifact from automatic reconstruction processes.
- Reactions that involve oxygen are integral to the energy metabolism of many organisms. Not constraining these reactions carefully can lead to predictions that deviate from the expected phenotype, i.e., allowing anaerobic growth that should not be possible. The portion of oxygen-containing reactions that are reversible varies strongly across all seven collections. Models in BiGG have the lowest variance whereas models from Path2Models, Ebrahim *et al.*, and OptFlux vary strongly (Figure [S133](#)).

#### 3.3.2 Scores




Figure S11: Total Score. Depicted are the sums of all test scores in all independent sections, applying the weights for individual test cases and sections as detailed in the snapshot report.

### 3.3.3 Independent Section

#### 3.3.3.1 Consistency




Figure S12: Consistency. Depicted are the sums of all test scores in this section, applying the weights of the individual test cases as detailed in the snapshot report.




Figure S13: Stoichiometric consistency




Figure S14: Mass Balance. Please note that any reaction where at least one metabolite lacks a formula annotation is considered as unbalanced for the purpose of this test.




Figure S15: Charge Balance. Please note that any reaction where at least one metabolite lacks charge information is considered as unbalanced for the purpose of this test.




Figure S16: Metabolite Connectivity




Figure S17: Unbounded Flux in Default Medium

### **3.3.3.2 Annotation - Metabolites**




Figure S18: Annotation - Metabolites. Depicted are the sums of all test scores in this section, applying the weights of the individual test cases as detailed in the snapshot report.




Figure S19: Presence of Metabolite Annotation

### **3.3.3.2.1 Metabolite Annotations Per Database**




Figure S20: Metabolite Pubchem.compound Annotation




Figure S21: Metabolite KEGG.compound Annotation




Figure S22: Metabolite SEED.compound Annotation




Figure S23: Metabolite InChIKey Annotation




Figure S24: Metabolite InChI Annotation




Figure S25: Metabolite ChEBI Annotation




Figure S26: Metabolite HMDB Annotation




Figure S27: Metabolite Reactome Annotation




Figure S28: Metabolite MetaNetX.chemical Annotation




Figure S29: Metabolite BiGG.metabolite Annotation




Figure S30: Metabolite BioCyc Annotation

### 3.3.3.2.2 Metabolite Annotation Conformity per Database




Figure S31: Correct Metabolite Pubchem.compound Annotation




Figure S32: Correct Metabolite KEGG.compound Annotation




Figure S33: Correct Metabolite SEED.compound Annotation




Figure S34: Correct Metabolite InChIKey Annotation




Figure S35: Correct Metabolite InChI Annotation




Figure S36: Correct Metabolite ChEBI Annotation




Figure S37: Correct Metabolite HMDB Annotation




Figure S38: Correct Metabolite Reactome Annotation




Figure S39: Correct Metabolite MetaNetX.chemical Annotation




Figure S40: Correct Metabolite BiGG.metabolite Annotation




Figure S41: Correct Metabolite BioCyc Annotation




Figure S42: Uniform Metabolite Identifier Namespace

### 3.3.3.3 Annotation - Reactions




Figure S43: Annotation - Reactions. Depicted are the sums of all test scores in this section, applying the weights of the individual test cases as detailed in the snapshot report.




Figure S44: Presence of Reaction Annotation

### **3.3.3.3.1 Reaction Annotations Per Database**




Figure S45: Reaction Rhea Annotation




Figure S46: Reaction KEGG.reaction Annotation




Figure S47: Reaction SEED.reaction Annotation




Figure S48: Reaction MetaNetX.reaction Annotation




Figure S49: Reaction BiGG.reaction Annotation




Figure S50: Reaction Reactome Annotation




Figure S51: Reaction Enzyme Classification Annotation




Figure S52: Reaction BRENDA Annotation




Figure S53: Reaction BioCyc Annotation

### 3.3.3.3.2 Reaction Annotation Conformity Per Database




Figure S54: Correct Reaction Rhea Annotation




Figure S55: Correct Reaction KEGG.reaction Annotation




Figure S56: Correct Reaction SEED.reaction Annotation




Figure S57: Correct Reaction MetaNetX.reaction Annotation




Figure S58: Correct Reaction BiGG.reaction Annotation




Figure S59: Correct Reaction Reactome Annotation




Figure S60: Correct Reaction Enzyme Classification Annotation




Figure S61: Correct Reaction BRENDA Annotation




Figure S62: Correct Reaction BioCyc Annotation




Figure S63: Uniform Reaction Identifier Namespace

### **3.3.3.4 Annotation - Genes**




Figure S64: Annotation - Genes. Depicted are the sums of all test scores in this section, applying the weights of the individual test cases as detailed in the snapshot report.




Figure S65: Presence of Gene Annotation

### **3.3.3.4.1 Gene Annotations Per Database**




Figure S66: Gene RefSeq Annotation




Figure S67: Gene UniProt Annotation




Figure S68: Gene EcoGene Annotation




Figure S69: Gene KEGG.genes Annotation




Figure S70: Gene NCBItgi Annotation




Figure S71: Gene NCBItgene Annotation




Figure S72: Gene NCBIprotein Annotation




Figure S73: Gene CCDS Annotation




Figure S74: Gene HPRD Annotation




Figure S75: Gene ASAP Annotation

### **3.3.3.4.2 Gene Annotation Conformity Per Database**




Figure S76: Correct Gene RefSeq Annotation




Figure S77: Correct Gene UniProt Annotation




Figure S78: Correct Gene EcoGene Annotation




Figure S79: Correct Gene KEGG.genes Annotation




Figure S80: Correct Gene NCBIgi Annotation




Figure S81: Correct Gene NCBIgene Annotation




Figure S82: Correct Gene NCBIProtein Annotation




Figure S83: Correct Gene CCDS Annotation




Figure S84: Correct Gene HPRD Annotation




Figure S85: Correct Gene ASAP Annotation

### **3.3.3.5 Annotation - SBO Terms**




Figure S86: Annotation - SBO Terms. Depicted are the sums of all test scores in this section, applying the weights of the individual test cases as detailed in the snapshot report.




Figure S87: Metabolite General SBO Presence




Figure S88: Metabolite SBO:0000247 Presence




Figure S89: Reaction General SBO Presence




Figure S90: Metabolic Reaction SBO:0000176 Presence




Figure S91: Transport Reaction SBO:0000185 Presence




Figure S92: Exchange Reaction SBO:0000627 Presence




Figure S93: Demand Reaction SBO:0000628 Presence




Figure S94: Sink Reaction SBO:0000632 Presence




Figure S95: Gene General SBO Presence




Figure S96: Gene SBO:0000243 Presence




Figure S97: Biomass Reaction SBO:0000629 Presence

### **3.3.4 Specific Section**

#### **3.3.4.1 SBML**




Figure S98: SBML Level and Version




Figure S99: FBC not Enabled

### **3.3.4.2 Basic Information**




Figure S100: Model Identifier Presence




Figure S101: Number of Metabolites




Figure S102: Number of Reactions




Figure S103: Number of Genes




Figure S104: Number of Compartments




Figure S105: Metabolic Coverage

### **3.3.4.3 Metabolite Information**




Figure S106: Unique Metabolites




Figure S107: Duplicate Metabolites in Identical Compartments




Figure S108: Metabolites Without Charge




Figure S109: Metabolites Without Formula




Figure S110: Number of Medium Components

#### 3.3.4.4 Reaction Information




Figure S111: Purely Metabolic Reactions




Figure S112: Purely Metabolic Reactions with Constraints




Figure S113: Transport Reactions




Figure S114: Transport Reactions with Constraints




Figure S115: Thermodynamic Reversibility of Purely Metabolic Reactions




Figure S116: Thermodynamic Reversibility of Purely Metabolic Reactions




Figure S117: Reactions with Partially Identical Annotations




Figure S118: Duplicate Reactions




Figure S119: Reactions with Identical Genes

### 3.3.4.5 Gene-Protein-Reaction (GPR) Association




Figure S120: Reactions without GPR




Figure S121: Fraction of Transport Reactions without GPR




Figure S122: Enzyme Complexes

### 3.3.4.6 Biomass




Figure S123: Biomass Reactions Identified




Figure S124: Biomass Consistency




Figure S125: Biomass Production in Default Medium




Figure S126: Biomass Production in Complete Medium




Figure S127: Blocked Biomass Precursors in Default Medium




Figure S128: Blocked Biomass Precursors in Complete Medium




Figure S129: Ratio of Direct Metabolites in Biomass Reaction




Figure S130: Number of Missing Essential Biomass Precursors

### **3.3.4.7 Energy Metabolism**




Figure S131: Non-Growth Associated Maintenance Reaction




Figure S132: Growth-associated Maintenance in Biomass Reaction




Figure S133: Number of Reversible Oxygen-Containing Reactions

### 3.3.4.8 Network Topology




Figure S134: Universally Blocked Reactions




Figure S135: Orphan Metabolites




Figure S136: Dead-end Metabolites




Figure S137: Stoichiometrically Balanced Cycles




Figure S138: Metabolite Production in Complete Medium




Figure S139: Metabolite Consumption in Complete Medium

### 3.3.4.9 Matrix Conditioning




Figure S140: Ratio Min/Max Non-Zero Coefficients




Figure S141: Independent Conservation Relations




Figure S142: Rank




Figure S143: Degrees of Freedom

## Supplementary References

- Beaulieu-Jones, Brett K, and Casey S Greene. 2017. “Reproducibility of Computational Workflows Is Automated Using Continuous Analysis.” *Nat. Biotechnol.* 35 (4): 342–46.
- Breiman, Leo. 2001. “Random Forests.” *Machine Learning* 45 (1): 5–32. <https://doi.org/10.1023/A:1010933404324>.
- Ebrahim, Ali, Joshua A Lerman, Bernhard O Palsson, and Daniel R Hyduke. 2013. “COBRApy: COnstraints-Based Reconstruction and Analysis for Python.” *BMC Syst. Biol.* 7 (August): 74.
- Heavner, Benjamin D, and Nathan D Price. 2015. “Transparency in Metabolic Network Reconstruction Enables Scalable Biological Discovery.” *Curr. Opin. Biotechnol.* 34 (August): 105–9.
- Heirendt, Laurent, Sylvain Arreckx, Thomas Pfau, Sebastián N Mendoza, Anne Richelle, Almut Heinken, Hulda S Haraldsdóttir, et al. 2017. “Creation and Analysis of Biochemical Constraint-Based Models: The COBRA Toolbox V3.0,” October.
- Louppe, Gilles, Louis Wehenkel, Antonio Sutera, and Pierre Geurts. 2013. “Understanding Variable Importances in Forests of Randomized Trees.” In *Proceedings of the 26th International Conference on Neural Information Processing Systems - Volume 1*, 431–39. NIPS’13. USA: Curran Associates Inc. <http://dl.acm.org/citation.cfm?id=2999611.2999660>.