

# Key access and utilization descriptors for rice genetic resources

This list consists of an initial set of characterization and evaluation descriptors for rice utilization. This strategic set of descriptors, together with passport data, will become the basis for the global accession level information portal being developed by Bioversity International with the support of the Global Crop Diversity Trust (GCDDT). It will facilitate access to and utilization of rice accessions held in genebanks and does not preclude the addition of further descriptors, should data subsequently become available.

Based on the comprehensive list of 'Descriptors for wild and cultivated Rice (*Oryza* spp.)' published by IRRI, WARDA (now Africa Rice Center) and Bioversity International in 2007, the evaluation category was subsequently compared with a number of sources such as results from the Global Public Goods Activity 4.2.1.1, UPOV technical guidelines for rice (2004), Descriptors for Rice (USDA, ARS, GRIN), the IRRI publication 'Standard Evaluation System for Rice' (SES, 2002), and with the traits that were awarded funds for further research by the Global Crop Diversity Trust through the Evaluation Award Scheme (EAS, 2008).

This minimal set defines a first priority set of characteristics to describe, to access and to utilize *Oryza* genetic resources. This list was developed in two different phases. During phase one, a list of highly discriminating Bioversity-IRRI descriptors for rice was defined and published in 2007 under the scientific direction of Dr Ruairaidh Sackville Hamilton of IRRI. For evaluation traits defined during phase two, a worldwide distribution of experts involved in an online survey was assured and the list was afterwards validated by a Core Advisory Group (see 'Contributors') led by Dr Edilberto D. Redoña of IRRI.

Biotic and abiotic stresses included in the list were chosen because of their wide geographic occurrence and significant economic impact at a global level.

Numbers in parentheses on the right-hand side are the corresponding descriptor numbers listed in the 2007 publication. Descriptors with numbers ending in 'letters' are new descriptors that were added during the development of the list below.

The code numbers of IRRI-SES for the corresponding characteristics are indicated beside the descriptor name between brackets [IS-] for easy reference. These codes belong to the Standard Evaluation System for Rice (INGER, Genetic Resources Centre, IRRI, July 1996).

To keep colour description simple, both Bioversity and IRRI colour codes have been listed. The IRRI colour coding system uses one code for one colour regardless of the descriptor.

## PLANT DATA

### Main heading

(7.2.3)

Date on which 80% of the plants are heading. It is specified either as the number of days from effective seeding date to main heading date or as effective seeding date and main heading date

**Auricle: colour**

(7.3.11)

Stage: late vegetative

(IRRI)

0	0	Absent (no auricles)
1	011	Whitish
2	062	Yellowish green
3	080	Purple
4	081	Light purple
5	084	Purple lines

**Flag leaf: attitude** (early observation)

(7.3.22)

Measured near the collar. Angle of attachment between the flag leaf blade and the main panicle axis. Record the average of five samples.

Stage: cultivated species at anthesis; wild species seven days after anthesis

1	Erect
3	Semi-erect (intermediate)
5	Horizontal
7	Descending

**Plant: height [cm] [IS-5]**

(7.3.25a)

Use actual measurement in cm from soil surface to the tip of the tallest panicle (awns excluded). For height measurements at other growth stages, specify the stage. Record in whole numbers (do not use decimals).

Stage: after flowering to maturity.

Alternatively, they can be coded as follows:

1	Semidwarf (lowland: less than 110 cm; upland: less than 90 cm)
5	Intermediate (lowland: 110–130 cm; upland: 90–125 cm)
9	Tall (lowland: more than 130 cm; upland: more than 125 cm)

**Culm: length [cm]**

(7.3.25)

Measured from ground level to the base of the panicle. Record the average of five actual measurements, to the nearest cm.

Stage: cultivated species after flowering to maturity; wild species seven days after anthesis.

Alternatively, cultivated species can be coded as follows:

1	Very short	(<50 cm)
2	Very short to short	(51–70 cm)
3	Short	(71–90 cm)
4	Short to intermediate	(91–105 cm)
5	Intermediate	(106–120 cm)
6	Intermediate to long	(121–140 cm)
7	Long	(141–155 cm)
8	Long to very long	(156–180 cm)
9	Very long	(>180 cm)

**Culm: anthocyanin colouration on nodes** (7.3.28)

The presence and distribution of purple colour from anthocyanin, observed on the outer surface of the nodes on the culm.

Stage: after flowering to near maturity

(IRRI)

0	0	Absent
1	080	Purple
2	081	Light purple
3	084	Purple lines

**Culm: underlying node colour** (7.3.29)

The underlying colour of the outer surface of the nodes on the culm, ignoring any anthocyanin colouration.

Stage: after flowering to near maturity

(IRRI)

0	0	No underlying colour visible due to anthocyanin
1	041	Light gold
2	060	Green

**Flag leaf: attitude (late observation)** (7.3.34)

(Cultivated species). Observed near the collar. Angle of attachment between the flag leaf blade and the main panicle axis. Record the average of five samples.

Stage: maturity

1	Erect
3	Semi-erect
5	Horizontal
7	Descending

**Stigma: colour** (7.4.2)

Observed at anthesis (between 0900 and 1400) using a hand lens

(IRRI)

1	010	White
2	061	Light green
3	030	Yellow
4	081	Light purple
5	080	Purple

### **Lemma: colour of apiculus** (early observation) (7.4.6)

Stage: cultivated species after anthesis to hard dough stage (pre-ripening stage); wild species at anthesis

	(IRRI)	
1	010	White
2	020	Straw
3	052	Brown (tawny)
4	060	Green
5	070	Red
6	071	Red apex
7	080	Purple
8	087	Purple apex
9	100	Black

### **Awns: distribution** (7.4.9)

(Cultivated species). The presence and distribution of awns along the panicle.

Stage: flowering to maturity

0	None (awnless)
1	Tip only
2	Upper quarter only
3	Upper half only
4	Upper three-quarters only
5	Whole length

### **Panicle: length [cm]** (7.4.18)

(Wild species). Length of main axis of panicle measured from the panicle base to the tip. Record the average of five representative plants.

Stage: seven days after anthesis or upon full panicle exertion

### **Panicle: attitude of main axis** (7.4.19)

Stage: near maturity

1	Upright
2	Semi-upright
3	Slightly drooping
4	Strongly drooping

### **Panicle: attitude of branches** (7.4.20)

The compactness of the panicle, classified according to its mode of branching, angle of primary branches, and spikelet density.

Stage: cultivated species near maturity; wild species seven days after anthesis

1	Erect (compact panicle)
3	Semi-erect (semi-compact panicle)
5	Spreading (open panicle)
7	Horizontal
9	Drooping

**Panicle: exertion**

(7.4.22)

Extent to which the panicle is exerted above the flag leaf sheath.

Stage: near maturity

- 1 Enclosed (panicle is partly or entirely enclosed within the leaf sheath of the flag leaf blade)
- 3 Partly exerted (panicle base is slightly beneath the collar of the flag leaf blade)
- 5 Just exerted (panicle base coincides with the collar of the flag leaf blade)
- 7 Moderately well exerted (panicle base is above the collar of the flag leaf blade)
- 9 Well exerted (panicle base appears well above the collar of the flag leaf blade)

**Lemma and palea: pubescence**

(7.5.4)

Visual assessment of the presence and distribution of mature grains using a hand lens

- 1 Glabrous
- 2 Hairs on lemma keel
- 3 Hairs on upper portion
- 4 Short hairs
- 5 Long hairs (velvety)

**Sterile lemma: length [mm]**

(7.5.10)

Record the average length of five spikelets. For spikelets with symmetrical sterile lemmas (i.e. sterile length the same on both sides), record the length here. For spikelets with asymmetrical sterile lemmas (i.e. sterile lemma on one side longer than that on the other), record here only the length of the shorter sterile lemma (see 7.5.11 for the longer sterile lemma).

May be coded as:

- 3 Short
- 5 Medium
- 7 Long
- 9 Extra long

**Longer sterile lemma: length [mm]**

(7.5.11)

(Only for spikelets with asymmetrical sterile lemmas). Record the average length of the longer sterile lemma on five spikelets.

May be coded as:

- 3 Short
- 5 Medium
- 7 Long
- 9 Extra long

**Sterile lemma: colour**

(7.5.13)

Observe five representative plants

(IRRI)

- 1 020 Straw
- 2 040 Gold
- 3 070 Red
- 4 080 Purple

**Caryopsis: length [mm]** (7.5.20)

**Caryopsis: shape** (7.5.22)

- 1 Round
- 2 Semi-round
- 3 Half spindle-shaped
- 4 Spindle-shaped
- 5 Long spindle-shaped

**Caryopsis: pericarp colour** (7.5.23)  
(IRRI)

- 1 010 White
- 2 051 Light brown
- 3 055 Speckled brown
- 4 050 Brown
- 5 070 Red
- 6 088 Variable purple
- 7 080 Purple

**Caryopsis: scent** (8.1.2)

From cooked kernel. Use freshly harvested grain. A molecular marker for fragrance is described in Section 12.3, **Fragrance** of 'Descriptors for wild and cultivated Rice (*Oryza* spp.)'

- 0 Non-scented
- 1 Lightly scented
- 2 Scented

**Endosperm amylose content [%]** (8.1.3)

Amylose content of all cultivars of low amylose and many of intermediate amylose is sensitive to high temperatures during grain-filling. Molecular markers for classifying amylose are listed in Section 12.1, **Amylose content** of 'Descriptors for wild and cultivated Rice (*Oryza* spp.)'

- 0 Waxy-glutinous (<3)
- 1 Very low (~9)
- 3 Low (~17)
- 5 Intermediate (~20)
- 7 High (~23)
- 9 Very high (>25)

**Phenotypic acceptability [IS-10]** (8.1.a)

Breeding objectives for each location vary. The score should reflect the overall acceptability of the variety in the location where it is being grown.

Stage: maturity

- 1 Excellent
- 3 Good
- 5 Fair
- 7 Poor
- 9 Unacceptable

## ABIOTIC STRESSES

<b>Cold</b> [IS-75]	(9.1)
<b>Heat</b> [IS-76]	(9.2)
<b>Drought</b> [IS-80]	(9.3)
<b>Alkali injury and salt injury</b> [IS-70-71]	(9.4)
<b>Flood or submergence</b> [IS-86]	(9.8)

## BIOTIC STRESSES

<b>Leaf blast</b> ( <i>Magnaporthe grisea</i> ) [IS-30]	(10.1.1)
<b>Brown spot</b> ( <i>Cochliobolus miyabeanus</i> ) [IS-32]	(10.1.3)
<b>Bacterial leaf streak</b> ( <i>Xanthomonas oryzae</i> pv. <i>oryzicola</i> ) [IS-33]	(10.1.5)
<b>Bacterial blight</b> ( <i>Xanthomonas oryzae</i> pv. <i>oryzae</i> ) [IS-35]	(10.1.7)
<b>Sheath blight</b> ( <i>Thanatephorus cucumeris</i> ) [IS-37]	(10.2.6)
<b>Rice tungro bacilliform virus</b> (RTBV) [IS-36]	(10.2.a)
<b>Brown planthopper</b> ( <i>Nilaparvata lugens</i> ) [IS-60]	(10.3.1)
<b>Stem borer</b> ( <i>Chilo suppressalis</i> ) [IS-63]	(10.3.5)

## CONTRIBUTORS

Bioversity is grateful to all the scientists and researchers who have contributed to the development of this strategic set of 'Key access and utilization descriptors for rice genetic resources', and in particular to Drs Ruairaidh Sackville Hamilton and Edilberto Redoña who provided scientific direction during the development of the characterization and evaluation categories, respectively. Adriana Alercia provided technical expertise and guided the entire production process.

## **CORE ADVISORY GROUP**

**Edilberto D. Redoña**, International Rice Research Institute (IRRI), Philippines  
**Teresita H. Borromeo**, University of the Philippines Los Baños, Philippines  
**Cesar P. Martinez**, Centro Internacional de Agricultura Tropical (CIAT), Colombia  
**Khin Than Nwe**, Department of Agricultural Research, Myanmar  
**Luis Salaices**, Oficina Española de Variedades Vegetales, Spain  
**N. Shobha Rani**, Directorate of Rice Research, India  
**Shengxiang Tang**, China National Rice Research Institute, China

## **REVIEWERS**

### **Australia**

**Laurie Lewin**, NSW Department of Primary Industries

### **Bangladesh**

**A. K. G. Md. Enamul Haque**, Bangladesh Rice Research Institute

### **Czech Republic**

**Iva Faberova**, Crop Research Institute, Prague

### **India**

**Narasimha Murthi Anishetty**

**Manish Kumar Pandey**, Directorate of Rice Research

**S. R. Pandravada**, National Bureau of Plant Genetic Resources (NBPGR), Regional Station, Rajendranagar

**J. C. Rana**, National Bureau of Plant Genetic Resources (NBPGR), Regional Station, Phagli, Shimla

**Natarajan Sivaraj**, National Bureau of Plant Genetic Resources (NBPGR)

### **Indonesia**

**Afif Nafisah**, Indonesian Center for Rice Research

**Untung Susanto**, Indonesian Center for Rice Research

### **Madagascar**

**Alain Ramanantsoanirina**, Centre National de la Recherche Applique au Développement Rural (FOFIFA/CENRADERU)

### **Malaysia**

**Site Noorzuraini Binti Abd Rahman**, Malaysian Agricultural Research and Development Institute (MARDI)

**Abdullah Md Zain**, University Malaysia Terengganu



**Philippines**

Ma. Socorro R. Almazan, International Rice Research Institute (IRRI)

Noel A. Catibog, Philippine Council for Agriculture, Forestry and Natural Resources Research and Development (PCARRD)

Thelma F. Padolina, Philippine Rice Research Institute (PhilRice)

Renato A. Reaño, Genetic Resources Center, International Rice Research Institute (TTC-GRC, IRRI)

**Republic of Korea**

Myung Chul Lee, National Agrobiodiversity Center

Sae-Jun Yang, National Institute of Crop Science (RDA)

**Russia**

Olga Romanova, N. I. Vavilov Institute of Plant Industry (VIR)

**Thailand**

Orapin Watanesek, Bureau of Rice Research and Development

**Vietnam**

Nguyen Thi Lang, Cuulong Delta Rice Research Institute