

# STAT\_757\_Assignment1

DUE 2/4/2018 11:59PM

*AG Schissler, modified from P. Hurtado*

*1/24/2018*

## i. DataCamp: Introduction to R [30 points]

Please complete the course an Introduction to R. You should have received an email with an invitation link. Please email me if you did not. If you already know R, please talk to me in class and follow up with an email to opt out.

## ii. Instructions for the rest of this assignment

The purpose of this portion of the assignment is to get a little experience making R Markdown documents as a way of nicely formatting output from R code while exploring the datasets from Sheather Ch.1 and learning to generate realizations of random variables (aka “fake data”). Modify this RMarkdown file (STAT\_757\_Assignment1.Rmd) and compile your document as a PDF (or Word document if you’re having LaTeX issue) and naming it according to the format SURNAME-FIRSTNAME-Assignment1.pdf, and emailing that PDF to the instructor by the due date listed above.

## 2. Reproduce the plots from Sheather Ch.1 [40 points]

Modify this file so that it reproduces all the output from the R script located at <http://www.stat.tamu.edu/~sheather/book/docs/rcode/Chapter1.R>. I’ve done the plots for the first dataset for you below. Remember that you will need to download each of the four data sets from [http://www.stat.tamu.edu/~sheather/book/data\\_sets.php](http://www.stat.tamu.edu/~sheather/book/data_sets.php), and set your working directory (under the “Session” menu in Rstudio) appropriately. (And yes, this really is as easy as copying the blocks of R code for each dataset into this document into the appropriate places!) Need help? First, see <http://rmarkdown.rstudio.com>. Especially the resources under Learning More (<http://rmarkdown.rstudio.com/#learning-more>).

Below are the plots that appear in Chapter 1 of the textbook. They were created from the R script <http://www.stat.tamu.edu/~sheather/book/docs/rcode/Chapter1.R> and the data files at [http://www.stat.tamu.edu/~sheather/book/data\\_sets.php](http://www.stat.tamu.edu/~sheather/book/data_sets.php).


### NFL Kicker Data

```
kicker <- read.csv("~/OneDrive - University of Nevada, Reno/Teaching/STAT_757/Sheather_data/Data/FieldG
## Sorry this line is too long, the data are labeled 'FieldGoals2003to2006.csv'

attach(kicker) ## THIS IS NOT USUALLY RECOMMENDED, ASK ME IN CLASS WHY NOT.

#Figure 1.1 on page 2
plot(kicker$FGtM1,kicker$FGt,
main="Unadjusted Correlation = -0.139",
xlab="Field Goal Percentage in Year t-1",ylab="Field Goal Percentage in Year t")
```

**Unadjusted Correlation = -0.139**



*#p-values on page 3*

```
fit.1 <- lm(FGt~FGtM1 +Name +FGtM1:Name,data=kicker)
anova(fit.1)
```

## Analysis of Variance Table

##

## Response: FGt

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
FGtM1	1	87	87.2	1.90	0.1760
Name	18	2252	125.1	2.73	0.0046
FGtM1:Name	18	418	23.2	0.51	0.9386
Residuals	38	1743	45.9		

*#slope and intercepts of lines in Figure 1.2 on page 3*

```
fit.2 <- lm(FGt ~ Name + FGtM1,data=kicker)
fit.2
```

##

## Call:

## lm(formula = FGt ~ Name + FGtM1, data = kicker)

##

## Coefficients:


	(Intercept)	NameDavid Akers
	126.687	-4.646
NameJason Elam		NameJason Hanson
	-3.017	2.117
NameJay Feely		NameJeff Reed
	-10.374	-8.296
NameJeff Wilkins		NameJohn Carney
	2.310	-5.977
NameJohn Hall		NameKris Brown

```
##           -8.486           -13.360
##      NameMatt Stover      NameMike Vanderjagt
##           8.736           4.896
##      NameNeil Rackers      NameOlindo Mare
##          -6.620          -13.036
##      NamePhil Dawson      NameRian Lindell
##           3.552           -4.867
##      NameRyan Longwell  NameSebastian Janikowski
##          -2.231          -3.976
##      NameShayne Graham      FGtM1
##           2.135           -0.504
```

*#Figure 1.2 on page 3*

```
plot(kicker$FGtM1,kicker$FGt,
main="Slope of each line = -0.504",
xlab="Field Goal Percentage in Year t-1",
ylab="Field Goal Percentage in Year t")
tt <- seq(60,100,length=1001)
slope.piece <- summary(fit.2)$coef[20]*tt
lines(tt,summary(fit.2)$coef[1]+slope.piece,lty=2)
for (i in 2:19)
{lines(tt,summary(fit.2)$coef[1]+summary(fit.2)$coef[i]+slope.piece,lty=2)}
```

**Slope of each line =  $-0.504$**



```
detach(kicker)
```

Newspaper circulation

and so on...

### 3. Generating fake data [30 points]

3.1 Generate 100 random variates from a normal distribution with mean 0 and standard deviation of 100. Summarize and plot the data. (Set a seed to make it reproducible).

3.2 Generate 1000 random variates from a beta distribution with the parameters  $\alpha$  and  $\beta$  both equal to 2. Summarize and plot the data. (Set a seed to make it reproducible).

3.3 Generate 10000 random variates from a binomial distribution with the parameters  $n = 10$  and  $p = 0.2$ . Summarize and plot the data. (Set a seed to make it reproducible).