

PowerPoint® Lectures for
University Physics, 14th Edition, Global Edition
– Hugh D. Young and Roger A. Freedman

Chapter 23

Electric Potential

Lectures by Gunsu Yun
(2018 Fall).

Contents modified by GS Yun,
CK Hong and JW Chung.

Goals for Chapter 23

- **Electric potential energy**
- **Electric potential**
- **Equipotential surfaces**
- **Electric potential \rightarrow electric field**

(chap.21) **Substance** (Charge) \rightarrow Force

\rightarrow **Field**, Field lines; Dipole

(chap.22) \rightarrow **Flux, Gauss's Law**

(chap.23) **Force (-Work)** \rightarrow Potential energy \rightarrow Potential;

Equipotential/Gradient \rightarrow **Field**

23.1 Electric potential energy

Electric potential energy in a uniform field


- In the figure, a uniform field exerts a downward force on a positive test charge.
- As the charge moves downward from point a to point b , **the work done by the field is independent of the path** the particle takes.

→ conservative force

→ potential energy

$$U = - \int_C \vec{F} \cdot d\vec{l} = - \int_C q_0 \vec{E} \cdot d\vec{l}$$

Point charge q_0 moving in a uniform electric field




The work done by the electric force is the same for any path from a to b :
 $W_{a \rightarrow b} = -\Delta U = q_0 E d$

A positive(negative) charge moving in a uniform field

- If the positive(negative) charge moves in the direction of the field, the field does positive(negative) work on the charge.
- The potential energy **decreases(increases)**.


Positive charge q_0 moves in the direction of \vec{E} :

- Field does *positive* work on charge.
- U decreases.



Negative charge q_0 moves in the direction of \vec{E} :

- Field does *negative* work on charge.
- U increases.




Electric potential energy of two point charges

- The work done by the electric field of one point charge on another does not depend on the path taken.
- Therefore, the electric potential energy **only depends on the distance** between the charges.

$$\begin{aligned}\Delta U &= U_b - U_a = -W_{a \rightarrow b} \\ &= -\int_a^b \vec{F} \cdot d\vec{l} \\ &= -\int_{r_a}^{r_b} \frac{1}{4\pi\epsilon_0} \frac{qq_0}{r^2} dr \\ &= \frac{qq_0}{4\pi\epsilon_0} \left(\frac{1}{r_b} - \frac{1}{r_a} \right)\end{aligned}$$

Test charge q_0 moves from a to b along an arbitrary path.



Electric potential energy of two point charges

Electric potential energy of two point charges



$$U = \frac{1}{4\pi\epsilon_0} \frac{qq_0}{r}$$

Values of two charges

Distance between two charges


Electric constant

- Potential energy is defined to be **zero** when the charges are infinitely far apart.



Electrical potential energy with several point charges

- The potential energy associated with q_0 depends on the other charges and their distances from q_0 .
- The electric potential energy is the **algebraic sum**:



Electric potential energy of point charge q_0 and collection of charges q_1, q_2, q_3, \dots

$$U = \frac{q_0}{4\pi\epsilon_0} \left(\frac{q_1}{r_1} + \frac{q_2}{r_2} + \frac{q_3}{r_3} + \dots \right) = \frac{q_0}{4\pi\epsilon_0} \sum_i \frac{q_i}{r_i}$$

Electric constant

Distances from q_0 to q_1, q_2, q_3, \dots


cf. Total potential energy U of several of point charges initially separated by infinite distances:

$$U_{\text{tot}} = \frac{1}{4\pi\epsilon_0} \sum_{i < j} \frac{q_i q_j}{r_{ij}}$$

23.2 Electric potential

- **Electric Potential(V)** is electric potential energy per unit charge.
- The electric potential of b with respect to a ($V_{ba} = V_b - V_a$) equals the minus of the work done by the electric force when a unit charge moves from b to a .
- SI unit of electric potential: **volt**, $V = \text{J/C}$

$$\begin{aligned} V_b - V_a &\equiv (U_b - U_a)/q_0 \\ &= - \int_a^b q_0^{-1} \vec{F} \cdot d\vec{l} \\ &= - \int_a^b \vec{E} \cdot d\vec{l} \end{aligned}$$



Electric potential of point charges

- The potential due to a single point charge is:

Electric potential due to a point charge

$$V = \frac{1}{4\pi\epsilon_0} \frac{q}{r}$$

Value of point charge

Distance from point charge to where potential is measured

Electric constant

Detailed description: This diagram shows the formula for the electric potential V due to a single point charge q. The formula is V = (1 / (4 * pi * epsilon_0)) * (q / r). Annotations include: 'Electric potential due to a point charge' pointing to V; 'Value of point charge' pointing to q; 'Distance from point charge to where potential is measured' pointing to r; and 'Electric constant' pointing to epsilon_0.

- Like electric field, potential is independent of the test charge q_0 that we use to define it.
- For a collection of point charges:

Electric potential due to a collection of point charges

$$V = \frac{1}{4\pi\epsilon_0} \sum_i \frac{q_i}{r_i}$$

Value of i th point charge

Distance from i th point charge to where potential is measured

Electric constant


Detailed description: This diagram shows the formula for the electric potential V due to a collection of point charges. The formula is V = (1 / (4 * pi * epsilon_0)) * sum_i (q_i / r_i). Annotations include: 'Electric potential due to a collection of point charges' pointing to V; 'Value of i-th point charge' pointing to q_i; 'Distance from i-th point charge to where potential is measured' pointing to r_i; and 'Electric constant' pointing to epsilon_0.

Typical electric potential differences

TABLE 17–1 Some Typical Potential Differences (Voltages)

Source	Voltage (approx.)
Thundercloud to ground	10^8 V
High-voltage power line	$10^5\text{--}10^6 \text{ V}$
Power supply for TV tube	10^4 V
Automobile ignition	10^4 V
Household outlet	10^2 V
Automobile battery	12 V
Flashlight battery	1.5 V
Resting potential across nerve membrane	10^{-1} V
Potential changes on skin (EKG and EEG)	10^{-4} V

Copyright © 2005 Pearson Prentice Hall, Inc.



The electron volt (a unit of energy)


- $\Delta U = q \Delta V$
- If a particle with the charge of e ($=1.602 \times 10^{-19}$ C) moves across a potential difference of 1 V, the change in **kinetic (or potential) energy** is defined as one **electron volt (eV)**:

$$1 \text{ eV} = 1.602 \times 10^{-19} \text{ J}$$

- Pohang Accelerator Laboratory: $\sim 1\text{--}10$ GeV
-

Moving through a potential difference

In Fig. 23.15 a dust particle with mass $m = 5.0 \times 10^{-9} \text{ kg}$ $= 5.0 \mu\text{g}$ and charge $q_0 = 2.0 \text{ nC}$ starts from rest and moves in a straight line from point a to point b . What is its speed v at point b ?



The mechanical energy $K+U$ is conserved since only conservative electric force acts on the particle.

$$0 + q_0 V_a = \frac{1}{2} m v^2 + q_0 V_b \quad \Rightarrow \quad v = \sqrt{\frac{2 q_0 (V_a - V_b)}{m}}$$

$$V_a = k \left(\frac{q}{r_{1a}} + \frac{-q}{r_{2a}} \right) = k q \left(\frac{1}{r_{1a}} - \frac{1}{r_{2a}} \right) = 1350 \text{ V}$$

$$V_b = -1350 \text{ V}$$



$$\Rightarrow v = \sqrt{\frac{2(2.0 \times 10^{-9} \text{ C})(2700 \text{ V})}{5.0 \times 10^{-9} \text{ kg}}} = 46 \text{ m/s}$$

23.3 Calculating electric potential

Electric potential and field of a charged conductor

- A solid conducting sphere of radius R has a total charge q .
- The electric field inside the sphere is zero.
- Outside, $\vec{E} = \hat{r} \frac{1}{4\pi\epsilon_0} \frac{q}{r^2}$

$$V = - \int_{\infty}^r \vec{E} \cdot d\vec{l}$$




Oppositely charged parallel plates

Find the potential at any height y between the two oppositely charged parallel plates discussed in Section 23.1 (Fig. 23.18).

$$V(y) = \frac{U(y)}{q_0} = \frac{yq_0E}{q_0} = yE$$

$$V_a - V_b = Ed \text{ and } E = \frac{V_a - V_b}{d} = \frac{V_{ab}}{d} \quad (\text{Volt/meter})$$



Line charge


Find the potential at a distance r from a very long line of charge with linear charge density (charge per unit length) λ .

$$\begin{aligned} V_2 - V_1 &= - \int_1^2 \vec{E} \cdot d\vec{l} = - \int_1^2 E_r dr = \frac{-\lambda}{2\pi\epsilon_0} \int_{r_1}^{r_2} \frac{dr}{r} \\ &= \frac{-\lambda}{2\pi\epsilon_0} \ln r \Big|_{r_1}^{r_2} = \frac{\lambda}{2\pi\epsilon_0} \ln \left(\frac{r_1}{r_2} \right) \end{aligned}$$

Taking $V = 0$ at the surface of the conducting wire of radius R ,

$$V(r) = \frac{\lambda}{2\pi\epsilon_0} \ln \frac{R}{r} \quad (\text{for } r \geq R)$$


How about $V(r)$ for $r < R$?



A ring of charge


Electric charge Q is distributed uniformly around a thin ring of radius a (Fig. 23.20). Find the potential at a point P on the ring axis at a distance x from the center of the ring.

$$V(x) = \frac{1}{4\pi\epsilon_0} \int \frac{dq}{r} = \frac{1}{4\pi\epsilon_0} \frac{1}{\sqrt{x^2 + a^2}} \int dq = \frac{1}{4\pi\epsilon_0} \frac{Q}{\sqrt{x^2 + a^2}}$$



Positive electric charge Q is distributed uniformly along a line of length $2a$ lying along the y -axis between $y = -a$ and $y = +a$ (Fig. 23.21). Find the electric potential at a point P on the x -axis at a distance x from the origin.

$$\begin{aligned} V(x) &= \frac{1}{4\pi\epsilon_0} \int_{y=-a}^{y=a} \frac{dQ}{r} = \frac{1}{4\pi\epsilon_0} \left(\frac{Q}{2a} \right) \int_{-a}^a \frac{dy}{\sqrt{y^2 + x^2}} \\ &= \frac{1}{4\pi\epsilon_0} \left(\frac{Q}{2a} \right) \ln \left(\sqrt{y^2 + x^2} + y \right) \Big|_{-a}^a \\ &= \frac{1}{4\pi\epsilon_0} \left(\frac{Q}{2a} \right) \ln \left(\frac{\sqrt{a^2 + x^2} + a}{\sqrt{a^2 + x^2} - a} \right) \end{aligned}$$




For $x \gg a$, $V(x) = \frac{1}{4\pi\epsilon_0} \frac{Q}{x}$.


23.4 Equipotential surfaces and field lines

- An **equipotential surface** is a surface on which the electric potential is the same.
- Field lines and equipotential surfaces are always **mutually perpendicular**.
- Examples of equipotential surfaces (blue lines) and electric field lines (red lines):


(a) A single positive charge



(b) An electric dipole




(c) Two equal positive charges




→ Electric field lines — Cross sections of equipotential surfaces

Equipotential and potential gradient


Topographic map:
contour lines (등고선)



Temperature map



Pressure map (기압도)




Equipotentials and conductors

- If the electric field had a **tangential component** at the surface of a conductor, a net amount of work would be done on a test charge by moving it around a loop as shown here—which is impossible because the electric force is conservative.


An impossible electric field

If the electric field just outside a conductor had a tangential component E_{\parallel} , a charge could move in a loop with net work done.



Equipotentials and conductors

- When all charges are **at rest** (*electro-statics*):
 - ✓ the surface of a conductor is always an **equipotential surface**.
 - ✓ the electric field just outside a conductor is always **perpendicular to the surface**.



- Cross sections of equipotential surfaces
- ➔ Electric field lines

23.5 Potential gradient

- The components of the electric field can be found by taking **partial derivatives** of the electric potential:

$$V = - \int \vec{E} \cdot d\vec{l} = - \left(\int E_x dx + \int E_y dy + \int E_z dz \right) \Rightarrow$$

Electric field
components found
from potential:

Each electric field component ...

$$E_x = - \frac{\partial V}{\partial x} \quad E_y = - \frac{\partial V}{\partial y} \quad E_z = - \frac{\partial V}{\partial z}$$

... equals the negative of the corresponding partial derivative of electric potential function V .

- Equivalently, the electric field is the negative **gradient** of the potential:

$$\vec{E} = -\nabla V = - \left(\hat{x} \frac{\partial V}{\partial x} + \hat{y} \frac{\partial V}{\partial y} + \hat{z} \frac{\partial V}{\partial z} \right)$$

Potential gradient

From Eq. (23.14) the potential at a radial distance r from a point charge q is $V = q/4\pi\epsilon_0 r$. Find the vector electric field from this expression for V .

$$\vec{E} = \hat{r}E_r \quad \text{by symmetry, where}$$

$$E_r = -\frac{\partial V}{\partial r} = -\frac{\partial}{\partial r} \left(\frac{q}{4\pi\epsilon_0 r} \right) = \frac{q}{4\pi\epsilon_0 r^2}$$

Component-wise:

Noting that $r = (x^2 + y^2 + z^2)^{1/2}$ and $\frac{\partial r}{\partial x} = \frac{x}{r}$,

$$E_x = -\frac{\partial V}{\partial x} = -\frac{\partial V}{\partial r} \frac{\partial r}{\partial x} = \frac{q}{4\pi\epsilon_0 r^2} \left(\frac{x}{r} \right)$$

Similarly,

$$E_y = -\frac{\partial V}{\partial y} = \frac{q}{4\pi\epsilon_0 r^2} \left(\frac{y}{r} \right)$$


$$E_z = -\frac{\partial V}{\partial z} = \frac{q}{4\pi\epsilon_0 r^2} \left(\frac{z}{r} \right)$$

$$\Rightarrow \vec{E} = \frac{q}{4\pi\epsilon_0 r^2} \left(\frac{\hat{x}x + \hat{y}y + \hat{z}z}{r} \right) = \frac{q}{4\pi\epsilon_0 r^2} \hat{r}$$

In **Example 23.11** (Section 23.3) we found that for a ring of charge with radius a and total charge Q , the potential at a point P on the ring's symmetry axis a distance x from the center is

$$V = \frac{Q}{4\pi\epsilon_0} \frac{1}{(x^2 + a^2)^{1/2}}$$

$$E_x = -\frac{\partial V}{\partial x} = \frac{Q}{4\pi\epsilon_0} \frac{x}{(x^2 + a^2)^{3/2}}$$



23 Summary

- **Electric potential energy**
 - **Electric potential**
 - **Equipotential surfaces**
 - **Finding electric field from electric potential**
-