

UNIVERSITY OF OSLO  
COMPUTATIONAL PHYSICS

---

**Project 2**

---



UiO : **University of Oslo**

---

Authors:

Birgitte Madsen

Magnus Isaksen

Soumya Chalakkal

---

Autumn 2015





**UiO • University of Oslo**

**Department of Physics**

**University of Oslo**

Sem Sælands vei 24

0371 Oslo, Norway

+47 22 85 64 28

<http://www.mn.uio.no/fysikk/english/>

**Course:**

Computational Physics

**Project number:**

2

**Link to GitHub folder:**

<https://??>

**Hand-in deadline:**

Monday, October 5, 2015

**Project Members:**

Birgitte Madsen

Magnus Isaksen

Soumya Chalakkal

**Copies:** 1

**Page count:** ??

**Appendices:** 0

**Completed:** ??

*The content of the report is freely available, but publication (with source) may only be made with the agreement of the authors.*





---

# TABLE OF CONTENTS

<b>Chapter 1</b>	<b>Introduction</b>	<b>1</b>
<b>Chapter 2</b>	<b>Method</b>	<b>3</b>
2.1	Nature of the problem . . . . .	3
2.2	Gauss-Legendre Method for Computing the Integral . . . . .	3
<b>Chapter 3</b>	<b>Conclusion</b>	<b>5</b>
<b>Appendix A</b>	<b>MatLab code for smt....</b>	<b>7</b>



# INTRODUCTION





---

# METHOD

## 2.1 Nature of the problem

## 2.2 Gauss-Legendre Method for Computing the Integral

To be able to use the Gauss-Legendre method to compute the integral in <sup>1</sup>, the limits of the integral must be made finite. Since the wave function

$$e^{-2\alpha x} \tag{2.1}$$

rapidly goes toward zero as  $x$  is increased (see fig), the integral can be approximated by the same integral with finite limits.

In this project, we <sup>2</sup> have accepted that  $10^{-9}$  is close enough to zero to neglect contributions from the part of the wave function when the wave function gives a value of this order. For  $x = 5$  the value of the wave function is  $e^{-10\alpha} \approx 2.1 \cdot 10^{-9}$ , when  $\alpha = 2$ . Hence, the considered integral that is to be solved by the Gauss-Legendre method is given by

$$\left\langle \frac{1}{|\mathbf{r}_1 - \mathbf{r}_2|} \right\rangle = \int_{-5}^5 \frac{e^{-2\alpha x}}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2 \tag{2.2}$$

---

<sup>1</sup>FiXme Note: ref. to problem eq

<sup>2</sup>FiXme Note: sorry about the "we"



---

## CONCLUSION

Conclude.... conclude.... conclude....



**A**

---

# MATLAB CODE FOR SMT....

This is how, we write MatLab code in the report

---

```
close all
clear all
clc
%I am a comment

filename = 'Results.xlsx';
sheet = 4;
xlRange = 'B3:C12';

[v,T,vT] = xlsread(filename, sheet, xlRange);
x10=v(:,1);y10=v(:,2);

figure
plot(??)
legend(??)

xlim(??)
ylim(??)

title(??)
xlabel('x')
ylabel('y')
```

---