DETERMINISTIC TWO-ZONE MODELING OF TRANSIENT LIBS PLASMA: CORRECTING THE LOCAL THERMODYNAMIC EQUILIBRIUM ILLUSION VIA COLLISIONAL-RADIATIVE MATRICES AND STATISTICAL INVERSION

BIRRUL WALIDAIN, NASRULLAH IDRIS, KHAIRUN SADDAMI

Abstract. The extraction of fundamental thermodynamic parameters, such as electron temperature (T) and density (n_e), in Laser-Induced Breakdown Spectroscopy (LIBS) frequently relies on the assumption of Local Thermodynamic Equilibrium (LTE) via the Saha equation. However, in rapidly expanding transient plasmas operating on nanosecond scales, the LTE criterion collapses, and intense spatial gradients induce severe optical self-absorption. This paper presents a deterministic computational architecture that replaces the homogeneous plasma assumption with a Two-Zone Model (Core and Shell). We strictly reject the Saha equilibrium assumption, instead solving the highly stiff Ordinary Differential Equations (ODEs) of a Collisional-Radiative (CR) model using an implicit solver with an analytical Jacobian matrix to preserve absolute thermodynamic conservation. Furthermore, rather than employing computationally expensive stochastic blind-search algorithms or black-box deep learning models, we utilize a statistical machine learning regression trained on deterministically synthesized spectra. This inverse mathematical solver effectively deconvolutes the self-absorbed macroscopic spectra to reveal the true microscopic kinetic parameters of the core, thereby correcting the McWhirter criterion and eliminating the LTE illusion.

Introduction

Laser-Induced Breakdown Spectroscopy (LIBS) has been widely established as a robust analytical technique for in-situ material characterization. Despite its empirical success, the theoretical extraction of plasma parameters relies heavily on the assumption that the plasma adheres to Local Thermodynamic Equilibrium (LTE). Under LTE, atomic state populations are assumed to follow the Boltzmann distribution, and ionization balances are governed by the Saha equation. However, laser-induced plasmas are inherently transient and chaotic, exhibiting extreme spatial gradients and rapid expansion velocities on nanosecond and femtosecond timescales. In such regimes, collisional rates often fail to equilibrate with radiative decay processes, causing a fundamental breakdown of the LTE condition.

Furthermore, the sharp temperature gradient between the hot, dense core and the cooler, expanding periphery induces profound optical self-absorption. Recent literature has attempted to address this self-reversal phenomenon by applying homogeneous plasma models or by utilizing global stochastic optimization algorithms (such as `CRS2-LM`) to fit experimental spectra. These

Date: March 23, 2026.

MSC2020: Primary 82D10 (Plasma and Plasmonics), Secondary 65L04 (Numerical Methods for Stiff ODE), 62J02 (General Nonlinear Regression).

approaches contain fatal thermodynamic flaws: they either ignore the spatial non-uniformity of the transient expansion or forcibly embed the Saha equilibrium assumption into mathematically partitioned spatial zones [**?**, **?**].

In rejecting black-box deep learning architectures and probabilistic curve-fitting that obscure physical realities, this study proposes a strictly deterministic computational framework. We implement a physically rigorous Two-Zone Model where the atomic population kinetics are governed purely by an exact Collisional-Radiative (CR) matrix, eliminating the Saha assumption entirely. By mapping the deterministic forward model to a rapid statistical machine learning regression for spectral inversion, this methodology provides a high-fidelity diagnostic tool capable of exposing the LTE illusion and validating the empirical limits of the McWhirter criteria [**?**].

Material and Methods

The computational architecture is strictly divided into a deterministic forward physical model—ensuring absolute conservation of mass and energy—and a statistical inverse solver for parameter extraction.

2.1**The Two-Zone Geometric Architecture** To capture the spatial heterogeneity of the self-absorption phenomenon without triggering the exponential computational cost of full 3D hydrodynamic simulations, the plasma is spatially discretized into two distinct regions: the emissive Core (V_1) and the absorptive Shell (V_p). The macroscopic intensity observed by the spectrometer (I_{obs}) is modeled via the 1D integration of the Radiative Transfer Equation (RTE):

$$I_{\text{obs}}(\lambda) = I_{\text{core}}(\lambda)\, \exp\!\big(-\kappa_{\text{shell}}(\lambda)\, d_{\text{shell}}\big) + I_{\text{shell}}(\lambda), \tag{1}$$

where I_{core} is the unattenuated emission from the core, κ_{shell} is the wavelength-dependent absorption coefficient of the periphery, and d_{shell} is the optical thickness of the absorbing layer [**?**].

2.2**Deterministic Population Kinetics (The Core)** The fundamental divergence of this methodology from established homogeneous models lies in the calculation of the energy state populations (n_i). We unequivocally reject the Saha equation. The temporal evolution of the fractional populations is solved via the Collisional-Radiative (CR) system of differential equations:

$$\frac{dn_i}{dt} = \sum_{j \neq i} n_j\, R_{ji} - n_i \sum_{j \neq i} R_{ij}, \tag{2}$$

where the rate matrix R_{ij} encapsulates all fundamental electron-impact collisional rates and radiative Einstein transition probabilities (A_{ij}, B_{ij}). Because the kinetic rates of collisional excitation and spontaneous emission are separated by multiple orders of magnitude (ranging from 10^{-14} to 10^{-3} seconds), the resulting Jacobian matrix is exceptionally stiff.

To prevent numerical divergence and unphysical negative particle populations—which routinely occur when using explicit methods such as standard Runge-Kutta—the temporal integration is executed using an implicit solver (Backward Differentiation Formula, BDF, or Radau). The computational bottleneck of inverting the $\mathcal{O}(N^3)$ matrix at each step is mitigated by

supplying the exact analytical Jacobian:

$$J_{ij} = \frac{\partial f_i}{\partial n_j}, \tag{3}$$

which is analytically identical to the CR rate matrix M for the linear ODE system $\mathbf{f}(n) = M\,n$. This guarantees that Newton iterations within the implicit solver converge quadratically, preserving strict thermodynamic conservation at every time step [**?**].

2.3**Socratic Boundaries and Synthetic Dataset Generation** The forward model computes the absorption coefficients (κ) by summing Voigt line profiles over all NIST-validated transitions, integrating both Stark broadening (pressure broadening, Lorentzian) and Doppler broadening (thermal motion, Gaussian):

$$\kappa(\lambda) = \sum_{\text{transitions}} n_i\,\sigma_{ij}(\lambda)\,\left(1 - \frac{n_k\,g_i}{n_i\,g_k}\right), \tag{4}$$

where the stimulated emission correction in parentheses ensures detailed balance is maintained [**?**].

By sweeping the parameter space ($T_e \in [4000, 20000]\,\mathrm{K}$ and $n_e \in [10^{14}, 10^{18}]\,\mathrm{cm}^{-3}$), we synthesize a comprehensive dataset of self-absorbed spectra. We apply Socratic ignorance regarding fundamental atomic inputs: transition probabilities A_{ki} and statistical weights g_i are strictly sourced from the NIST Atomic Spectra Database [**?**]. Transitions lacking validated empirical cross-sections are not synthetically approximated, ensuring the mathematical space does not exceed the boundaries of verified physical literature.

2.4**Inverse Modeling via Statistical Regression** To solve the inverse problem—extracting the pure core temperature (T_{core}) and electron density ($n_{e,\text{core}}$) from a convoluted experimental spectrum—we bypass computationally exhaustive and often divergent stochastic blind-search algorithms. Instead, intermediate statistical machine learning algorithms (e.g., Support Vector Regression or Ridge Regression) are trained on the deterministically generated synthetic dataset. Here, the algorithm acts purely as a high-speed mathematical inversion operator rather than a predictive physical model, translating the distorted line profiles back into their foundational thermodynamic parameters.

Expected Results and Discussion

By applying the trained regression model to experimental LIBS spectra, this computational architecture is expected to successfully deconvolute the optical self-absorption. Consequently, the extracted parameters (T and n_e) will reflect the true, unattenuated state of the plasma core.

When these corrected parameters are substituted back into the classical McWhirter criterion [**?**]:

$$N_e \geq 1.6 \times 10^{12}\,T^{1/2}\,(\Delta E)^3, \tag{5}$$

we hypothesize a significant deviation from the values derived via traditional homogeneous Boltzmann plot methods. This will quantitatively demonstrate that previous assertions of LTE in transient LIBS plasmas are largely mathematical illusions caused by the failure to isolate the absorptive shell from the emissive core. The stability of the implicit CR solver guarantees

that this correction is grounded in absolute thermodynamic conservation rather than statistical curve-fitting anomalies.

References

[1] Fujimoto, T. (2004). *Plasma Spectroscopy*. Clarendon Press, Oxford.

[2] Griem, H. R. (1997). *Principles of Plasma Spectroscopy*. Cambridge University Press, Cambridge.

[3] Hairer, E., & Wanner, G. (1996). *Solving Ordinary Differential Equations II: Stiff and Differential-Algebraic Problems* (2nd ed.). Springer, Berlin.

[4] McWhirter, R. W. P. (1965). Spectral intensities. In R. H. Huddlestone & S. L. Leonard (Eds.), *Plasma Diagnostic Techniques* (pp. 201–264). Academic Press, New York.

[5] Kramida, A., Ralchenko, Yu., Reader, J., & NIST ASD Team (2023). *NIST Atomic Spectra Database* (version 5.11). National Institute of Standards and Technology, Gaithersburg, MD. `https://physics.nist.gov/asd`

[6] Rybicki, G. B., & Lightman, A. P. (1979). *Radiative Processes in Astrophysics*. Wiley-Interscience, New York.

[7] van Regemorter, H. (1962). Rate of collisional excitation in stellar atmospheres. *The Astrophysical Journal*, **136**, 906–915.

Department of Physics, Faculty of Mathematics and Natural Sciences, Universitas Syiah Kuala, Banda Aceh, Indonesia

Email address: `birrul.walidain@mhs.usk.ac.id`