Doolittle Decomposition

From Compact LU Factorization to Partial Pivoting

Pengju Jin Siquan Fang Haoye He

Zhejiang University

October 6, 2026

Outline

1. Motivation and the Basic Method
2. Compact Storage
3. Shortcomings of the Basic Method
4. The Partial-Pivoting Algorithm
5. Why the Algorithm Is Correct
6. Numerical Example with Row Swaps
7. In-Place Implementation, Solving, and Cost
8. Conclusion

Why LU Factorization?

Solve $A\mathbf{x} = \mathbf{b}$ with $A \in \mathbb{R}^{n\times n}$ nonsingular, for many $\mathbf{b}$:

$$A = LU \implies \underbrace{L\mathbf{y} = \mathbf{b}}_{\text{forward}}, \quad \underbrace{U\mathbf{x} = \mathbf{y}}_{\text{back}}.$$

Decompose *once*; each new $\mathbf{b}$ then costs only $O(n^2)$.

Roadmap

Basic formulas & compact storage $\rightarrow$ shortcomings $\rightarrow$ partial pivoting ($PA = LU$) $\rightarrow$ correctness.

The Doolittle Convention

LU is not unique: $LU = (LD)(D^{-1}U)$ for any nonsingular diagonal D. Fix the scale by making L *unit* lower triangular ($\ell_{ii} = 1$):

$$L = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ \ell_{21} & 1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ \ell_{n1} & \cdots & \ell_{n,n-1} & 1 \end{bmatrix}, \qquad U = \begin{bmatrix} u_{11} & u_{12} & \cdots & u_{1n} \\ 0 & u_{22} & \cdots & u_{2n} \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & u_{nn} \end{bmatrix}.$$

For nonsingular A (when the factorization exists), this makes L, U **unique**.

Recursive Formulas

Since $a_{ij} = \sum_{k=1}^{\min(i,j)} \ell_{ik} u_{kj}$, with $\ell_{ii} = 1$:

Row i of U ($i \le j$):

$$u_{ij} = a_{ij} - \sum_{k=1}^{i-1} \ell_{ik} u_{kj}$$

Column i of L ($j > i$, $u_{ii} \neq 0$):

$$\ell_{ji} = \frac{a_{ji} - \sum_{k=1}^{i-1} \ell_{jk} u_{ki}}{u_{ii}}$$

Stage i: compute row i of U, then column i of L (empty sums are zero).

Basic Algorithm and Existence Condition

Set $\ell_{ii} = 1$. For $i = 1, \ldots, n$:

1. compute u_{ij}, $j = i, \ldots, n$;
2. if $u_{ii} = 0$, stop (no pivot);
3. compute ℓ_{ji}, $j = i + 1, \ldots, n$.

Existence (nonsingular A)

All pivots are nonzero **iff** every leading principal minor is nonzero:

$$\Delta_k = \det A_{1:k,1:k} \neq 0, \quad k = 1, \ldots, n, \qquad u_{kk} = \Delta_k / \Delta_{k-1}.$$

Worked Example (No Pivoting)

$$A = \begin{bmatrix} 2 & 1 & 1 \\ 4 & -6 & 0 \\ -2 & 7 & 2 \end{bmatrix}.$$

Stage 1: $u_{11} = 2,\ u_{12} = 1,\ u_{13} = 1; \quad \ell_{21} = 2,\ \ell_{31} = -1.$

Stage 2: $u_{22} = -6 - 2 \cdot 1 = -8, \quad u_{23} = -2, \quad \ell_{32} = (-8)/(-8) = -1.$

Stage 3: $u_{33} = 2 - (-1)(1) - (-1)(-2) = 1.$

$$L = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -1 & -1 & 1 \end{bmatrix}, \quad U = \begin{bmatrix} 2 & 1 & 1 \\ 0 & -8 & -2 \\ 0 & 0 & 1 \end{bmatrix}, \qquad LU = A. \checkmark$$

Works here — but not for every nonsingular matrix.

Why Is It Called the Compact Form?

Known structure is free (L unit lower triangular, U upper triangular). Store both factors in **one** $n \times n$ array:

$$c_{ij} = \begin{cases} u_{ij}, & i \le j, \\ \ell_{ij}, & i > j, \end{cases} \qquad L = I + \operatorname{tril}(C, -1), \quad U = \operatorname{triu}(C).$$

$$C = \begin{bmatrix} 2 & 1 & 1 \\ 2 & -8 & -2 \\ -1 & -1 & 1 \end{bmatrix}$$

Storage	Scalars
Two full matrices	$2n^2$
Two triangular arrays	$n(n+1)$
Doolittle compact	$\mathbf{n^2}$

- Elimination link: multipliers sit where entries were zeroed, so the factorization can **overwrite** A in place.

Shortcoming 1: Zero Pivot despite Nonsingularity

$$A_0 = \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}, \qquad \det A_0 = -1 \neq 0.$$

First pivot $u_{11} = 0$, so $\ell_{21} = 1/0$ is undefined. If $A_0 = LU$ with unit lower L, the first column needs $u_{11} = 0$ *and* $\ell_{21} u_{11} = 1$ — impossible. Swapping rows gives

$$PA_0 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \qquad P = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \qquad L = I,\ U = PA_0.$$

Key point

A zero pivot does **not** imply singularity — just a bad row order.

Shortcoming 2: Small Pivot Amplifies Error

$$A_\varepsilon = \begin{bmatrix} \varepsilon & 1 \\ 1 & 1 \end{bmatrix}, \quad \mathbf{b} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, \quad 0 < \varepsilon \ll 1.$$

Without swapping, $\ell_{21} = 1/\varepsilon$ is huge: $u_{22} = 1 - \frac{1}{\varepsilon}$, $y_2 = 2 - \frac{1}{\varepsilon}$.

With $\varepsilon = 10^{-5}$ and 4 significant digits both round to -1.000×10^5, so

$$\widehat{x}_2 = 1, \qquad \widehat{x}_1 = \frac{1 - \widehat{x}_2}{10^{-5}} = 0 \qquad (\text{exact } x_1, x_2 \approx 1).$$

Diagnosis

Not a divide-by-zero issue but *error propagation*; swapping makes the pivot 1.

The Fix: Partial Pivoting

At step k, pick the largest magnitude in the **current** column of $B^{(k)}$:

$$p \in \arg\max_{k \le i \le n} |b_{ik}^{(k)}|; \qquad \text{swap rows } k, p, \text{ then eliminate.}$$

- Uses the *updated* column, not the original column of A.
- A nonzero candidate gives a nonzero pivot; in exact arithmetic $|\ell_{ik}| \le 1$.

Limits

Bounding multipliers does not prevent element growth, nor cure ill-conditioning.

Partial-Pivoting Algorithm: $PA = LU$

P records row swaps; L unit lower triangular, U upper triangular. Left-multiplying by P only reorders *equations*.

Init $B \leftarrow A$, $L \leftarrow I_n$, $P \leftarrow I_n$. For $k = 1, \ldots, n$:

1. $p \in \arg\max_{k \le i \le n} |b_{ik}|$.
2. if $b_{pk} = 0$, stop (A singular).
3. if $p \neq k$: swap rows k, p of B and P; swap **only the first** $k-1$ **entries** of rows k, p of L.
4. for $i = k+1, \ldots, n$: $\ell_{ik} \leftarrow b_{ik}/b_{kk}$; $b_{ij} \leftarrow b_{ij} - \ell_{ik} b_{kj}$ $(j > k)$; $b_{ik} \leftarrow 0$.

Finally $U \leftarrow B$.

Why Swap the Already-Computed L?

- Columns $1..k-1$ of L record the multipliers each row used earlier; swapping rows of the working matrix swaps those histories.
- Mispaired multipliers **break** $PA = LU$.
- Swap **only** the finished first $k-1$ entries — swapping whole L rows would move the unit diagonal and break the structure.

Nonsingular A Always Has a Nonzero Pivot

Theorem

If A is nonsingular, in exact arithmetic every step of partial pivoting finds a nonzero pivot and terminates with $PA = LU$.

Proof.

Each step is a row swap (det ± 1) followed by unit lower-triangular elimination (det 1), so the working matrix keeps a nonzero determinant. Its leading part is upper triangular with the nonzero pivots on the diagonal, so in the block form $\begin{bmatrix} U_{11} & U_{12} \\ 0 & S_k \end{bmatrix}$ the trailing block satisfies $\det S_k \neq 0$. Hence S_k is nonsingular, its first column is not all zero, and the max-magnitude pivot is nonzero. Induction over k gives $PA = LU$. □

Conversely, an all-zero candidate column would give $\det S_k = 0 \Rightarrow \det A = 0$.

Example: Two Row Swaps

$$A = \begin{bmatrix} 2 & 1 & 1 \\ 4 & -6 & 0 \\ -2 & 7 & 2 \end{bmatrix}.$$

Step 1. Max in column 1 is 4: swap rows 1, 2; $\ell_{21} = \frac{1}{2},\ \ell_{31} = -\frac{1}{2}$:

$$B^{(2)} = \begin{bmatrix} 4 & -6 & 0 \\ 0 & 4 & 1 \\ 0 & 4 & 2 \end{bmatrix}, \quad L^{(2)} = \begin{bmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 1 & 0 \\ -\frac{1}{2} & 0 & 1 \end{bmatrix}.$$

Step 2. Column 2 ties at 4: choose row 3, swapping rows 2, 3 of B *and the finished first column of* L:

$$\widetilde{B} = \begin{bmatrix} 4 & -6 & 0 \\ 0 & 4 & 2 \\ 0 & 4 & 1 \end{bmatrix}, \quad \widetilde{L} = \begin{bmatrix} 1 & 0 & 0 \\ -\frac{1}{2} & 1 & 0 \\ \frac{1}{2} & 0 & 1 \end{bmatrix} \Rightarrow\ \ell_{32} = 1,\ b_{33} = -1.$$

Result and Verification

$$L = \begin{bmatrix} 1 & 0 & 0 \\ -\frac{1}{2} & 1 & 0 \\ \frac{1}{2} & 1 & 1 \end{bmatrix}, \quad U = \begin{bmatrix} 4 & -6 & 0 \\ 0 & 4 & 2 \\ 0 & 0 & -1 \end{bmatrix}, \quad P = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}, \quad LU = \begin{bmatrix} 4 & -6 & 0 \\ -2 & 7 & 2 \\ 2 & 1 & 1 \end{bmatrix} = PA. \checkmark$$

Skipping the L swap is fatal

Swapping only B gives $L_{\text{wrong}} = \begin{bmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 1 & 0 \\ -\frac{1}{2} & 1 & 1 \end{bmatrix}$, for which $(L_{\text{wrong}}U)_{21} = 2 \neq -2 = (PA)_{21}$.

In-Place Compact Implementation

Let $C \leftarrow A$. At step k: columns $1..k-1$ hold L, rows $1..k-1$ hold U, and the trailing block holds S_k. Swapping rows k, p swaps **entire rows** of C (finished U rows are untouched since $k, p \geq k$).

$C \leftarrow A$, $\pi \leftarrow (1, \ldots, n)$. For $k = 1, \ldots, n$:

1. $p \in \arg\max_{k \leq i \leq n} |c_{ik}|$; if $c_{pk} = 0$, stop.
2. swap rows k, p of C and $\pi_k \leftrightarrow \pi_p$.
3. for $i > k$: $c_{ik} \leftarrow c_{ik}/c_{kk}$; $c_{ij} \leftarrow c_{ij} - c_{ik}c_{kj}$, $j > k$.

Read L, U back by the compact rule; $P_{i,\pi_i} = 1$, so $(PA)_{ij} = a_{\pi_i, j}$.

$$C = \begin{bmatrix} 4 & -6 & 0 \\ -\frac{1}{2} & 4 & 2 \\ \frac{1}{2} & 1 & -1 \end{bmatrix}, \qquad \pi = (2, 3, 1).$$

Solving $A\mathbf{x} = \mathbf{b}$ with $PA = LU$

$$A\mathbf{x} = \mathbf{b} \Rightarrow PA\mathbf{x} = P\mathbf{b} \Rightarrow LU\mathbf{x} = P\mathbf{b} \Rightarrow \boxed{L\mathbf{y} = P\mathbf{b}, \quad U\mathbf{x} = \mathbf{y}.}$$

With $\widetilde{b}_i = b_{\pi_i}$:

$$y_i = \widetilde{b}_i - \sum_{j=1}^{i-1} \ell_{ij} y_j, \qquad x_i = \frac{y_i - \sum_{j=i+1}^{n} u_{ij} x_j}{u_{ii}}.$$

Example

$\mathbf{b} = (4, -2, 7)^{\mathsf{T}} \Rightarrow P\mathbf{b} = (-2, 7, 4)^{\mathsf{T}}$: $y = (-2, 6, -1)^{\mathsf{T}} \Rightarrow x = (1, 1, 1)^{\mathsf{T}}$.

Cost and Stability

Work

- Factorization: $\approx \frac{2}{3}n^3$.
- Pivot search + swaps: $O(n^2)$; overall still $O(n^3)$.
- Each new $\mathbf{b}$: $O(n^2)$.

Stability

- Exact: no zero-pivot failure, $|\ell_{ik}| \leq 1$.
- Floating point: usually better, growth still possible.
- Ill-conditioned A stays sensitive.

Two different questions

Algebraic correctness $\neq$ floating-point stability. A tiny pivot justifies a warning, not a claim of singularity.

Conclusion

- The unit-diagonal convention gives recursive formulas and a **compact** n^2 storage that works in place.
- Zero or tiny pivots make the basic method fail or lose accuracy.
- **Partial pivoting** yields $PA = LU$, keeping the computed multipliers consistent on each row swap.
- Nonsingular A always completes in exact arithmetic, terminating with $PA = LU$.

References

- G. H. Golub and C. F. Van Loan, *Matrix Computations*, 4th ed., Johns Hopkins University Press, 2013.
- N. J. Higham, *Accuracy and Stability of Numerical Algorithms*, 2nd ed., SIAM, 2002.
- R. L. Burden and J. D. Faires, *Numerical Analysis*, 9th ed., Brooks/Cole, 2011.

Thank you! Questions?